

Bis(2-amino-4-methylpyridinium) *trans*-diaquabis(pyrazine-2,3-dicarboxylato)-cuprate(II) hexahydrate

Hossein Eshtiagh-Hosseini,^{a*} Fabienne Gschwind,^b
Nafiseh Alfi^a and Masoud Mirzaei^a

^aDepartment of Chemistry, School of Sciences, Ferdowsi University of Mashhad, Mashhad 917791436, Iran, and ^bDepartment of Chemistry, University of Fribourg, Chemin Du Musée 9, 1700 Fribourg, Switzerland
Correspondence e-mail: heshtiagh@ferdowsi.um.ac.ir


Received 27 May 2010; accepted 15 June 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.036; wR factor = 0.092; data-to-parameter ratio = 17.5.

The title compound, $(\text{C}_6\text{H}_9\text{N}_2)_2[\text{Cu}(\text{C}_6\text{H}_2\text{N}_2\text{O}_4)_2(\text{H}_2\text{O})_2] \cdot 6\text{H}_2\text{O}$, consists of a mononuclear *trans*- $[\text{Cu}(\text{pzdc})_2(\text{H}_2\text{O})_2]^{2-}$ dianion (pzdc is pyrazine-2,3-dicarboxylate) and two [ampyH]⁺ cations (ampy is 2-amino-4-methylpyridine) with six water molecules of solvation. The Cu^{II} atom is hexacoordinated by two pzdc groups and two water molecules. The coordinated water molecules are in *trans*-diaxial positions and the pzdc dianion acts as a bidentate ligand through an O atom of the carboxylate group and the N atom of the pyrazine ring. There are diverse hydrogen-bonding interactions, such as N—H...O and O—H...O contacts, which lead to the formation of a three-dimensional supramolecular architecture.

Related literature

For the crystal structure of pyrazine-2,3-dicarboxylic acid (pzdcH₂), see: Takusagawa & Shimada (1973). For complexes of pzdcH₂ with manganese and zinc, see: Eshtiagh-Hosseini *et al.* (2010*a,b*). For the structure of bis(2,4,6-triamino-1,3,5-triazin-1-ium) pyrazine-2,3-dicarboxylate tetrahydrate, see: Eshtiagh-Hosseini *et al.* (2010*c*). For a review article on water cluster chemistry, see: Aghabozorg *et al.* (2010).



Experimental

Crystal data

$(\text{C}_6\text{H}_9\text{N}_2)_2[\text{Cu}(\text{C}_6\text{H}_2\text{N}_2\text{O}_4)_2(\text{H}_2\text{O})_2] \cdot 6\text{H}_2\text{O}$
 $M_r = 758.17$
Triclinic, $P\bar{1}$
 $a = 6.9075$ (14) Å
 $b = 8.4710$ (17) Å
 $c = 14.505$ (3) Å
 $\alpha = 78.28$ (3)°
 $\beta = 83.62$ (3)°
 $\gamma = 85.81$ (3)°
 $V = 824.8$ (3) Å³
 $Z = 1$
Mo $K\alpha$ radiation
 $\mu = 0.75$ mm⁻¹
 $T = 293$ K
 $0.3 \times 0.2 \times 0.1$ mm

Data collection

Stoe IPDS 2 diffractometer
Absorption correction: for a sphere [modified Dwiggins (1975) interpolation procedure]
 $T_{\min} = 0.743$, $T_{\max} = 0.745$
17015 measured reflections
4684 independent reflections
4282 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.092$
 $S = 1.04$
4684 reflections
268 parameters
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.36$ e Å⁻³
 $\Delta\rho_{\min} = -0.52$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N3}-\text{H13} \cdots \text{O4}^{\text{i}}$	0.82	1.91	2.7221 (19)	169
$\text{N4}-\text{H14A} \cdots \text{O8}^{\text{ii}}$	0.77	2.12	2.8879 (19)	177
$\text{N4}-\text{H14B} \cdots \text{O3}^{\text{i}}$	0.84 (2)	2.07 (2)	2.903 (2)	168 (2)
$\text{O5}-\text{H5B} \cdots \text{O7}^{\text{i}}$	0.79 (3)	1.92 (3)	2.703 (2)	173 (3)
$\text{O5}-\text{H5A} \cdots \text{O4}^{\text{i}}$	0.82 (3)	2.09 (3)	2.8556 (18)	157 (3)
$\text{O8}-\text{H8B} \cdots \text{O1}^{\text{iii}}$	0.76 (3)	2.03 (3)	2.7838 (18)	173 (3)
$\text{O6}-\text{H6B} \cdots \text{O4}^{\text{iii}}$	0.81 (4)	2.06 (4)	2.839 (2)	162 (3)
$\text{O7}-\text{H17B} \cdots \text{O8}^{\text{iv}}$	0.76 (4)	2.04 (4)	2.797 (2)	172 (4)

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+1, -y, -z$; (iii) $x+1, y, z$; (iv) $x, y+1, z$.

Data collection: *X-AREA* (Stoe & Cie, 2009); cell refinement: *X-RED* (Stoe & Cie, 2009); data reduction: *X-RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Crystal Impact, 2009); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Ferdowsi University of Mashhad is gratefully acknowledged for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2310).

References

- Aghabozorg, H., Eshtiagh-Hosseini, H., Salimi, A. R. & Mirzaei, M. (2010). *J. Iran. Chem. Soc.* **7**, 289–300.
Crystal Impact (2009). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Dwiggins, C. W. (1975). *Acta Cryst.* **A31**, 146–148.
Eshtiagh-Hosseini, H., Aghabozorg, H. & Mirzaei, M. (2010*a*). *Acta Cryst.* **E66**. Submitted. [XU2784]

- Eshtiagh-Hosseini, H., Hassanpoor, A., Alfi, N., Mirzaei, M., Fromm, K. M., Shokrollahi, A., Gschwind, F. & Karami, E. (2010*b*). *J. Coord. Chem.* In the press.
- Eshtiagh-Hosseini, H., Hassanpoor, A., Canadillas-Delgado, L. & Mirzaei, M. (2010*c*). *Acta Cryst.* **E66**, o1368–o1369.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (2009). *X-AREA* and *X-RED* Stoe & Cie, Darmstadt, Germany.
- Takusagawa, F. & Shimada, A. (1973). *Chem. Lett.* pp. 1121–1123.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.

supplementary materials

Acta Cryst. (2010). E66, m826-m827 [doi:10.1107/S1600536810023081]

Bis(2-amino-4-methylpyridinium) *trans*-diaquabis(pyrazine-2,3-dicarboxylato)cuprate(II) hexahydrate

H. Eshtiagh-Hosseini, F. Gschwind, N. Alfi and M. Mirzaei

Comment

Dicarboxylate ligands are widely used to assemble supramolecular network organized by coordination bonds, hydrogen bonds and π - π stacking interaction. Due to the manifold N- and O-donors of pyridine or pyrazine-(di)carboxylic ligands, metal pyridine- or pyrazine dicarboxylates can contrast versatile structural motifs, which finally aggregate to generate various supramolecular architectures with interesting properties. As ones of the dicarboxylate ligands, pzdcH₂ have drawn extensive attentions. For the first time, Takusagawa & Shimada (1973) by single crystal X-ray diffraction, determined the structure of pzdcH₂. Continuing with our previous works on synthesizing coordination compounds *via* proton transfer mechanism including zinc atom (Eshtiagh-Hosseini *et al.*, 2010a), manganese atom (Eshtiagh-Hosseini *et al.*, 2010b), Bis(2,4,6-triamino-1,3,5-triazin-1-ium) pyrazine-2,3-dicarboxylate tetrahydrate (Eshtiagh-Hosseini *et al.*, 2010c), herein, we planned the reaction between pzdcH₂, ampy, and copper(II) chloride which resulted in the formation of (ampy)₂[Cu(pzdc)₂(H₂O)₂].6H₂O (Fig. 1). Crystal packing diagram related to the title compound is also rendered in the Fig. 2. As you can see, the equatorial plane is occupied by two (pzdc)²⁻ ligands coordinating through the pyridine nitrogen and one oxygen of the deprotonated carboxylate groups. The two coordinated water molecules occupy axial plane. This compound consists of an anionic moiety, *trans*-[Cu(pzdc)₂(H₂O)₂]²⁻ complex, counter-ions, (ampy)⁺, and six uncoordinated water molecules. The Cu—O and Cu—N bond distances related to (pzdc)²⁻ ligand in herein presented compound are in the category of 1.9644 (13) Å, and 1.9840 (14) Å, respectively. These observed bond lengths are comparable with Zn(II) polymeric coordination compound which recently reported by our research group (Eshtiagh-Hosseini *et al.*, 2010a). In this polymeric compound which consist of only (pzdc)²⁻ coordinative ligand, {(C₃H₁₂N₂)₂[Zn(C₁₀H₂O₈)₂]}_n, Zn—O and Zn—N bond distances are 2.0317 (15) to 2.2437 (15) Å, and 2.0901 (18) Å, respectively. These data show in this polymeric compound Zn—O bond distance is longer than herein presented compound. The intermolecular forces between the anionic and cationic parts in the title compound consist of hydrogen bonding and ion pairing interactions. Indeed, six uncoordinated water molecules increase the number of hydrogen bonds in the crystalline network and lead to the formation of (H₂O)_n clusters throughout the crystalline network (see Review article by Aghabozorg *et al.* 2010).

Experimental

A solution of pzdcH₂ (0.18 mmol, 0.03 mg) in water (10 ml) refluxed for 1 hr, then a solution of CuCl₂.6H₂O (0.02 mmol, 0.01 g) was added dropwise and continued refluxing for 6 hrs at 60°C. The obtained blue solution gave blue block like crystals of title compound after slow evaporation of solvent at room temperature.

Refinement

The space group was confirmed by using PLATON software package. The structure was solved by direct methods using SHELXS-97 and refined using full matrix least-squares on F^2 with the SHELX-97 package. H-Atoms were constrained to the parent site using a rigid model. A final verification of possible voids was performed using the VOID routine on PLATON software.

Figures




Fig. 1. Molecular structure of $(ampy)_2[Cu(pzdc)_2(H_2O)_2] \cdot 6H_2O$. Ellipsoids are drawn at the 50% probability level. Hydrogen atoms are omitted for further clarity.





Fig. 2. Packing diagram of $(ampy)_2[Cu(pzdc)_2(H_2O)_2] \cdot 6H_2O$.

Bis(2-amino-4-methylpyridinium) *trans*-diaquabis(pyrazine-2,3-dicarboxylato)cuprate(II) hexahydrate

Crystal data



$M_r = 758.17$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 6.9075(14)\ \text{\AA}$

$b = 8.4710(17)\ \text{\AA}$

$c = 14.505(3)\ \text{\AA}$

$\alpha = 78.28(3)^\circ$

$\beta = 83.62(3)^\circ$

$\gamma = 85.81(3)^\circ$

$V = 824.8(3)\ \text{\AA}^3$

$Z = 1$

$F(000) = 395.0$

$D_x = 1.526\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 34097 reflections

$\theta = 3.8\text{--}59.7^\circ$

$\mu = 0.75\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Block, blue

$0.3 \times 0.2 \times 0.1\ \text{mm}$

Data collection

Stoe IPDS 2
diffractometer

Radiation source: fine-focus sealed tube
graphite

4684 independent reflections

4282 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

Detector resolution: 6.67 pixels mm⁻¹ $\theta_{\max} = 29.8^\circ$, $\theta_{\min} = 2.5^\circ$
 rotation method scans $h = -9 \rightarrow 9$
 Absorption correction: for a sphere $k = -11 \rightarrow 11$
 modified Dwiggin (1975) interpolation procedure
 $T_{\min} = 0.743$, $T_{\max} = 0.745$ $l = -20 \rightarrow 20$
 17015 measured reflections

Refinement

Refinement on F^2 Primary atom site location: structure-invariant direct methods
 Least-squares matrix: full Secondary atom site location: difference Fourier map
 $R[F^2 > 2\sigma(F^2)] = 0.036$ Hydrogen site location: inferred from neighbouring sites
 $wR(F^2) = 0.092$ H atoms treated by a mixture of independent and constrained refinement
 $S = 1.03$ $w = 1/[\sigma^2(F_o^2) + (0.0453P)^2 + 0.5453P]$
 4684 reflections where $P = (F_o^2 + 2F_c^2)/3$
 268 parameters $(\Delta/\sigma)_{\max} = 0.001$
 0 restraints $\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.52 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. Absorption correction: Interpolation using Int. Tab. Vol. C (1992) p. 523, Tab. 6.3.3.3 for values of μ_R in the range 0-2.5, and Int. Tab. Vol. II (1959) p. 302; Table 5.3.6 B for μ_R in the range 2.6-10.0. The interpolation procedure of C. W. Dwiggin Jr is used with some modification.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.0000	0.0000	0.5000	0.01658 (8)
O3	0.30782 (17)	0.49841 (13)	0.17617 (8)	0.0195 (2)
O4	0.11843 (16)	0.64677 (13)	0.26546 (8)	0.0180 (2)
O1	-0.09643 (16)	0.33706 (13)	0.27415 (8)	0.0190 (2)
O5	0.1806 (2)	-0.17345 (17)	0.40519 (9)	0.0258 (3)
O2	-0.12649 (16)	0.11778 (13)	0.38957 (8)	0.0189 (2)
N1	0.19550 (18)	0.16670 (15)	0.45776 (8)	0.0147 (2)
N2	0.43093 (18)	0.41844 (16)	0.37769 (9)	0.0173 (2)

supplementary materials

C1	-0.0415 (2)	0.24286 (17)	0.34432 (10)	0.0144 (2)
C2	0.1458 (2)	0.27423 (16)	0.38093 (9)	0.0134 (2)
C5	0.2651 (2)	0.40007 (17)	0.34084 (10)	0.0139 (2)
C4	0.4760 (2)	0.31081 (19)	0.45398 (11)	0.0187 (3)
H4	0.5897	0.3212	0.4806	0.022*
C3	0.3583 (2)	0.18338 (18)	0.49501 (10)	0.0171 (3)
H3	0.3936	0.1103	0.5483	0.021*
C6	0.2244 (2)	0.52402 (17)	0.25287 (10)	0.0148 (2)
N3	0.17729 (19)	-0.08982 (15)	0.12469 (9)	0.0175 (2)
H13	0.1634	-0.1765	0.1614	0.021*
N4	0.2985 (2)	-0.23942 (16)	0.01280 (10)	0.0200 (3)
H14A	0.3217	-0.2450	-0.0394	0.024*
C10	0.2467 (2)	-0.09571 (18)	0.03481 (10)	0.0160 (3)
C9	0.2613 (2)	0.05143 (19)	-0.03132 (11)	0.0184 (3)
C8	0.2056 (2)	0.19545 (18)	-0.00370 (11)	0.0197 (3)
C11	0.1206 (2)	0.05082 (19)	0.15352 (11)	0.0212 (3)
H11	0.0728	0.0489	0.2162	0.025*
C12	0.1332 (3)	0.19396 (19)	0.09169 (12)	0.0223 (3)
C7	0.2186 (3)	0.3528 (2)	-0.07308 (14)	0.0288 (4)
H7A	0.2698	0.4314	-0.0443	0.043*
H7B	0.0911	0.3898	-0.0912	0.043*
H7C	0.3034	0.3379	-0.1281	0.043*
O6	0.7121 (2)	0.67499 (18)	0.32078 (13)	0.0381 (4)
O7	0.4817 (2)	0.9530 (2)	0.28417 (12)	0.0385 (3)
O8	0.60159 (18)	0.25375 (14)	0.18411 (8)	0.0203 (2)
H12	0.097 (4)	0.287 (3)	0.1154 (19)	0.040 (7)*
H14B	0.289 (3)	-0.322 (3)	0.0565 (17)	0.024 (5)*
H9	0.311 (3)	0.045 (2)	-0.0939 (15)	0.017 (5)*
H5B	0.267 (4)	-0.130 (3)	0.372 (2)	0.040 (7)*
H5A	0.138 (4)	-0.236 (3)	0.377 (2)	0.040 (7)*
H8A	0.516 (4)	0.326 (3)	0.1824 (19)	0.039 (7)*
H8B	0.677 (4)	0.279 (3)	0.2116 (19)	0.034 (6)*
H6B	0.823 (5)	0.645 (4)	0.306 (2)	0.057 (9)*
H6A	0.632 (6)	0.606 (5)	0.333 (3)	0.076 (11)*
H17A	0.570 (5)	0.885 (4)	0.288 (2)	0.049 (8)*
H17B	0.524 (5)	1.032 (4)	0.259 (3)	0.062 (10)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.01841 (13)	0.01479 (13)	0.01464 (12)	-0.00519 (9)	-0.00398 (9)	0.00436 (9)
O3	0.0237 (5)	0.0173 (5)	0.0150 (5)	0.0014 (4)	0.0007 (4)	0.0003 (4)
O4	0.0203 (5)	0.0131 (5)	0.0190 (5)	0.0006 (4)	-0.0017 (4)	-0.0002 (4)
O1	0.0208 (5)	0.0176 (5)	0.0172 (5)	-0.0032 (4)	-0.0062 (4)	0.0032 (4)
O5	0.0265 (6)	0.0295 (6)	0.0239 (6)	-0.0017 (5)	-0.0016 (5)	-0.0116 (5)
O2	0.0193 (5)	0.0175 (5)	0.0185 (5)	-0.0068 (4)	-0.0056 (4)	0.0037 (4)
N1	0.0172 (5)	0.0131 (5)	0.0129 (5)	-0.0012 (4)	-0.0023 (4)	0.0003 (4)
N2	0.0158 (5)	0.0174 (6)	0.0179 (6)	-0.0020 (4)	-0.0023 (4)	-0.0007 (5)

C1	0.0154 (6)	0.0140 (6)	0.0134 (6)	-0.0015 (5)	-0.0013 (5)	-0.0014 (5)
C2	0.0149 (6)	0.0124 (6)	0.0121 (6)	-0.0004 (5)	-0.0013 (5)	-0.0006 (5)
C5	0.0149 (6)	0.0125 (6)	0.0133 (6)	0.0006 (5)	-0.0005 (5)	-0.0014 (5)
C4	0.0165 (6)	0.0196 (7)	0.0196 (7)	-0.0020 (5)	-0.0051 (5)	-0.0009 (5)
C3	0.0193 (6)	0.0165 (6)	0.0148 (6)	0.0005 (5)	-0.0044 (5)	-0.0006 (5)
C6	0.0154 (6)	0.0127 (6)	0.0154 (6)	-0.0035 (5)	-0.0015 (5)	0.0004 (5)
N3	0.0221 (6)	0.0146 (5)	0.0143 (5)	-0.0001 (4)	-0.0021 (4)	0.0007 (4)
N4	0.0262 (6)	0.0161 (6)	0.0162 (6)	0.0006 (5)	-0.0010 (5)	-0.0009 (5)
C10	0.0155 (6)	0.0175 (6)	0.0149 (6)	-0.0022 (5)	-0.0036 (5)	-0.0010 (5)
C9	0.0185 (6)	0.0192 (7)	0.0156 (6)	-0.0020 (5)	-0.0021 (5)	0.0014 (5)
C8	0.0194 (7)	0.0159 (7)	0.0223 (7)	-0.0034 (5)	-0.0057 (5)	0.0022 (5)
C11	0.0277 (8)	0.0199 (7)	0.0161 (6)	0.0016 (6)	-0.0033 (6)	-0.0043 (6)
C12	0.0282 (8)	0.0164 (7)	0.0232 (7)	-0.0002 (6)	-0.0067 (6)	-0.0044 (6)
C7	0.0340 (9)	0.0173 (7)	0.0308 (9)	-0.0031 (6)	-0.0035 (7)	0.0058 (6)
O6	0.0263 (7)	0.0240 (7)	0.0620 (10)	-0.0045 (5)	0.0142 (7)	-0.0127 (7)
O7	0.0273 (7)	0.0304 (7)	0.0498 (9)	-0.0033 (6)	-0.0024 (6)	0.0108 (7)
O8	0.0208 (5)	0.0204 (5)	0.0203 (5)	0.0013 (4)	-0.0044 (4)	-0.0051 (4)

Geometric parameters (Å, °)

Cu1—O2	1.9644 (13)	N3—C11	1.358 (2)
Cu1—O2 ⁱ	1.9644 (12)	N3—H13	0.8202
Cu1—N1	1.9840 (14)	N4—C10	1.335 (2)
Cu1—N1 ⁱ	1.9840 (14)	N4—H14A	0.7669
Cu1—O5	2.4038 (15)	N4—H14B	0.84 (2)
Cu1—O5 ⁱ	2.4038 (15)	C10—C9	1.412 (2)
O3—C6	1.2467 (18)	C9—C8	1.376 (2)
O4—C6	1.2597 (18)	C9—H9	0.95 (2)
O1—C1	1.2364 (18)	C8—C12	1.416 (2)
O5—H5B	0.79 (3)	C8—C7	1.500 (2)
O5—H5A	0.82 (3)	C11—C12	1.356 (2)
O2—C1	1.2732 (18)	C11—H11	0.9300
N1—C3	1.3291 (19)	C12—H12	0.93 (3)
N1—C2	1.3477 (18)	C7—H7A	0.9600
N2—C4	1.333 (2)	C7—H7B	0.9600
N2—C5	1.3486 (18)	C7—H7C	0.9600
C1—C2	1.5119 (19)	O6—H6B	0.81 (4)
C2—C5	1.387 (2)	O6—H6A	0.81 (4)
C5—C6	1.517 (2)	O7—H17A	0.80 (3)
C4—C3	1.393 (2)	O7—H17B	0.76 (4)
C4—H4	0.9300	O8—H8A	0.81 (3)
C3—H3	0.9300	O8—H8B	0.76 (3)
N3—C10	1.3475 (19)		
O2—Cu1—O2 ⁱ	180.00 (4)	N1—C3—H3	120.1
O2—Cu1—N1	83.31 (5)	C4—C3—H3	120.1
O2 ⁱ —Cu1—N1	96.69 (5)	O3—C6—O4	126.59 (14)
O2—Cu1—N1 ⁱ	96.69 (5)	O3—C6—C5	116.73 (13)

supplementary materials

O2 ⁱ —Cu1—N1 ⁱ	83.31 (5)	O4—C6—C5	116.56 (13)
N1—Cu1—N1 ⁱ	180.00 (7)	C10—N3—C11	122.73 (14)
O2—Cu1—O5	90.70 (5)	C10—N3—H13	116.8
O2 ⁱ —Cu1—O5	89.30 (5)	C11—N3—H13	120.3
N1—Cu1—O5	90.80 (6)	C10—N4—H14A	119.0
N1 ⁱ —Cu1—O5	89.20 (6)	C10—N4—H14B	117.9 (15)
O2—Cu1—O5 ⁱ	89.30 (5)	H14A—N4—H14B	122.6
O2 ⁱ —Cu1—O5 ⁱ	90.70 (5)	N4—C10—N3	118.61 (14)
N1—Cu1—O5 ⁱ	89.20 (6)	N4—C10—C9	123.36 (14)
N1 ⁱ —Cu1—O5 ⁱ	90.80 (6)	N3—C10—C9	118.02 (14)
O5—Cu1—O5 ⁱ	180.00 (5)	C8—C9—C10	120.30 (14)
Cu1—O5—H5B	113 (2)	C8—C9—H9	123.0 (12)
Cu1—O5—H5A	127.8 (19)	C10—C9—H9	116.7 (12)
H5B—O5—H5A	107 (3)	C9—C8—C12	119.11 (14)
C1—O2—Cu1	114.87 (9)	C9—C8—C7	121.06 (15)
C3—N1—C2	119.43 (13)	C12—C8—C7	119.83 (15)
C3—N1—Cu1	129.00 (10)	C12—C11—N3	120.60 (15)
C2—N1—Cu1	111.57 (10)	C12—C11—H11	119.7
C4—N2—C5	117.45 (13)	N3—C11—H11	119.7
O1—C1—O2	126.27 (13)	C11—C12—C8	119.24 (15)
O1—C1—C2	118.40 (13)	C11—C12—H12	117.2 (17)
O2—C1—C2	115.33 (12)	C8—C12—H12	123.5 (17)
N1—C2—C5	120.04 (13)	C8—C7—H7A	109.5
N1—C2—C1	114.86 (12)	C8—C7—H7B	109.5
C5—C2—C1	125.09 (12)	H7A—C7—H7B	109.5
N2—C5—C2	121.23 (13)	C8—C7—H7C	109.5
N2—C5—C6	114.89 (13)	H7A—C7—H7C	109.5
C2—C5—C6	123.87 (13)	H7B—C7—H7C	109.5
N2—C4—C3	122.10 (14)	H6B—O6—H6A	117 (3)
N2—C4—H4	118.9	H17A—O7—H17B	107 (3)
C3—C4—H4	118.9	H8A—O8—H8B	105 (3)
N1—C3—C4	119.74 (14)		

Symmetry codes: (i) $-x, -y, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H13 \cdots O4 ⁱⁱ	0.82	1.91	2.7221 (19)	169
N4—H14A \cdots O8 ⁱⁱⁱ	0.77	2.12	2.8879 (19)	177
N4—H14B \cdots O3 ⁱⁱ	0.84 (2)	2.07 (2)	2.903 (2)	168 (2)
O5—H5B \cdots O7 ⁱⁱ	0.79 (3)	1.92 (3)	2.703 (2)	173 (3)
O5—H5A \cdots O4 ⁱⁱ	0.82 (3)	2.09 (3)	2.8556 (18)	157 (3)
O8—H8B \cdots O1 ^{iv}	0.76 (3)	2.03 (3)	2.7838 (18)	173 (3)
O6—H6B \cdots O4 ^{iv}	0.81 (4)	2.06 (4)	2.839 (2)	162 (3)
O7—H17B \cdots O8 ^v	0.76 (4)	2.04 (4)	2.797 (2)	172 (4)

Symmetry codes: (ii) $x, y-1, z$; (iii) $-x+1, -y, -z$; (iv) $x+1, y, z$; (v) $x, y+1, z$.

Fig. 1




Fig. 2

