

Bis{2-[2-(2-benzoylhydrazin-1-ylidene)-methyl]-6-methoxyphenolato}iron(III) chloride monohydrate

Li-Fei Zou, Yu-Qin Ma, Gui-Miao Yu, Feng-Jiao Gan and Yun-Hui Li*

School of Chemistry and Environmental Engineering, Changchun University of Science and Technology, Changchun 130022, People's Republic of China
Correspondence e-mail: liyh@cust.edu.cn


Received 9 June 2010; accepted 16 June 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.051; wR factor = 0.153; data-to-parameter ratio = 13.1.

In the title mononuclear iron(III) complex, $[\text{Fe}(\text{C}_{15}\text{H}_{13}\text{N}_2\text{O}_3)_2]\text{Cl}\cdot\text{H}_2\text{O}$, the Fe^{III} atom has a distorted octahedral geometry and is six-coordinated by four O atoms and two N atoms from two ligands. In the crystal structure, the complex cations, Cl^- anions and water molecules are connected into a chain along [100] through $\text{N}-\text{H}\cdots\text{O}$, $\text{O}-\text{H}\cdots\text{Cl}$ and $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds. Two adjacent chains are linked by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For the applications of metal-Schiff base compounds, see: Dilworth (1976); Merchant & Clothia (1970); Pickart *et al.* (1983). For the ligand synthesis, see: Pouralimardan *et al.* (2007); Sacconi (1954). For related structures, see: Gao *et al.* (1998); Monfared *et al.* (2007); Yu *et al.* (2010).



Experimental

Crystal data

$[\text{Fe}(\text{C}_{15}\text{H}_{13}\text{N}_2\text{O}_3)_2]\text{Cl}\cdot\text{H}_2\text{O}$

$M_r = 647.86$

Monoclinic, $P2_1/c$

$a = 12.7778(10)\text{ \AA}$

$b = 22.7113(18)\text{ \AA}$

$c = 10.0604(7)\text{ \AA}$

$\beta = 94.542(1)^\circ$

$V = 2910.4(4)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.67\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.24 \times 0.18 \times 0.15\text{ mm}$

Data collection

Bruker SMART APEX CCD

diffractometer

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.857$, $T_{\max} = 0.907$

14540 measured reflections

5098 independent reflections

3508 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.052$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$

$wR(F^2) = 0.153$

$S = 0.98$

5098 reflections

390 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.95\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.48\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

Fe1—O1	2.070 (3)	Fe1—O5	1.901 (3)
Fe1—O2	1.904 (3)	Fe1—N2	2.106 (3)
Fe1—O4	2.062 (3)	Fe1—N4	2.124 (3)

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

D—H \cdots A	D—H	H \cdots A	D \cdots A	D—H \cdots A
N1—H1B \cdots Cl ⁱ	0.86	2.25	3.087 (3)	163
N3—H3B \cdots O1W	0.86	1.92	2.759 (4)	164
O1W—H1WA \cdots O5 ⁱⁱ	0.85	2.39	3.045 (4)	134
O1W—H1WB \cdots Cl1	0.85	2.37	3.198 (3)	163

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 1, -y + 1, -z + 2$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT-Plus* (Bruker, 2007); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *SHELXL97* and *publCIF* (Westrip, 2010).

We thank the Jilin Environmental Protection Bureau Foundation of China (2007-28) and Changchun University of Science and Technology for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2322).

References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Bruker (2007). *SMART* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
Dilworth, J.-R. (1976). *Coord. Chem. Rev.* **21**, 29–62.
Gao, S., Weng, Z.-Q. & Liu, S.-X. (1998). *Polyhedron*, **17**, 3595–3606.
Merchant, J. R. & Clothia, D. S. (1970). *J. Med. Chem.* **13**, 335–336.
Monfared, H. H., Sadighian, S., Kamyabi, M. A. & Mayer, P. (2007). *J. Mol. Catal. A*, **304**, 139–146.
Pickart, L., Goodwin, W. H., Burgua, W., Murphy, T. B. & Johnson, D. K. (1983). *Biochem. Pharmacol.* **32**, 3868–3871.
Pouralimardan, O., Chamayou, A. C., Janiak, C. & Monfared, H. H. (2007). *Inorg. Chim. Acta*, **360**, 1599–1608.
Sacconi, L. (1954). *Z. Anorg. Allg. Chem.* **275**, 249–256.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A*, **64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.
Yu, G.-M., Li, Y.-H., Zou, L.-F., Zhu, J.-W. & Liu, X.-Q. (2010). *Acta Cryst. E*, **66**, m693–m694.

supplementary materials

Acta Cryst. (2010). E66, m828 [doi:10.1107/S1600536810023226]

Bis{2-[(2-benzoylhydrazin-1-ylidene)methyl]-6-methoxyphenolato}iron(III) chloride monohydrate

L.-F. Zou, Y.-Q. Ma, G.-M. Yu, F.-J. Gan and Y.-H. Li

Comment

Studies of acylhydrazone Schiff base and the dependence of their chelation mode with transition metal ions have been of significant interest. On one hand, their metal compounds have been reported to act as enzyme inhibitors (Dilworth, 1976) and are useful due to their pharmacological applications (Merchant & Clothia, 1970). On the other hand, it seems to be a good candidate for catalytic oxidation studies because of their stability to resist oxidation (Pickart *et al.*, 1983). These findings have triggered the exploration of new molecular clusters based on acylhydrazone Schiff base. During the last several years, the crystal structures of metal compounds with 3-methoxysalicylaldehyde benzoylhydrazide have been attracted tremendous interest (Gao *et al.*, 1998; Monfared *et al.*, 2007; Yu *et al.*, 2010). As a continuation of our effort in this system, the preparation and crystal structure of the title Schiff base iron(III) compound are reported here.

The molecular structure of the title compound is illustrated in Fig. 1, which consists of one mononuclear $[\text{Fe}(\text{C}_{15}\text{H}_{13}\text{N}_2\text{O}_3)_2]^+$ cation, one Cl^- anion and one water molecule. The Fe^{III} atom has a distorted octahedral geometry and is six-coordinated by four O atoms and two N atoms from two ligands (Table 1). In one ligand, the strained angle of $\text{O}1-\text{Fe}1-\text{N}2$ [74.54 (11) $^\circ$] correlates with the bite angle for the five-membered chelate ring $\text{Fe}1-\text{O}1-\text{C}7-\text{N}1-\text{N}2$, and the loose angle of $\text{O}2-\text{Fe}1-\text{N}2$ [84.74 (11) $^\circ$] correlates with the six-membered ring $\text{Fe}1-\text{N}2-\text{C}8-\text{C}9-\text{C}10-\text{O}2$. The axial angle $\text{N}2-\text{Fe}1-\text{N}4$ [159.46 (12) $^\circ$] deviates significantly from the ideal 180 $^\circ$. Similar case occurs for another ligand. In the crystal structure, the complex cations, Cl^- anions and water molecules are connected into a chain through $\text{N}-\text{H}\cdots\text{O}$, $\text{O}-\text{H}\cdots\text{Cl}$ and $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds. Two adjacent chains are linked by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. (Fig. 2 and Table 2).

Experimental

The 3-methoxysalicylaldehyde benzoylhydrazide ligand (H_2L) was prepared in a similar manner according to the reported procedures (Pouralimardan *et al.*, 2007; Sacconi, 1954). The title compound was synthesized by adding $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (27.0 mg, 0.1 mmol) to a solution of H_2L (27.3 mg, 0.10 mmol) in methanol (15 ml). The resulting mixture was stirred for 3 h at room temperature to afford a dark brown solution and then filtered. The filtrate was allowed to stand at room temperature for about three weeks and black crystals were produced at the bottom of the vessel on slow evaporation of methanol.

Refinement

All H atoms were placed in calculated positions and refined using a riding model, with $\text{C}-\text{H} = 0.93$ (aromatic), 0.96 (methyl) Å and $\text{N}-\text{H} = 0.86$ Å and with $U_{\text{iso}}(\text{H}) = 1.2(1.5$ for methyl) $U_{\text{eq}}(\text{C}, \text{N})$. Water H atoms were located in a difference Fourier map and refined as riding, with $\text{O}-\text{H} = 0.85$ Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$.

supplementary materials

Figures




Fig. 1. Molecular structure of the title compound. H atoms are omitted for clarity. Displacement ellipsoids are drawn at the 30% probability level.



Fig. 2. One-dimensional chain structure in the title compound. Hydrogen bonds are shown as green dashed lines.

Bis{2-[(2-benzoylhydrazin-1-ylidene)methyl]-6-methoxyphenolato}iron(III) chloride monohydrate

Crystal data

[Fe(C₁₅H₁₃N₂O₃)₂]Cl·H₂O

$F(000) = 1340$

$M_r = 647.86$

$D_x = 1.479 \text{ Mg m}^{-3}$

Monoclinic, $P2_1/c$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Hall symbol: -P 2ybc

Cell parameters from 4767 reflections

$a = 12.7778 (10) \text{ \AA}$

$\theta = 4.8\text{--}51.7^\circ$

$b = 22.7113 (18) \text{ \AA}$

$\mu = 0.67 \text{ mm}^{-1}$

$c = 10.0604 (7) \text{ \AA}$

$T = 296 \text{ K}$

$\beta = 94.542 (1)^\circ$

Block, black

$V = 2910.4 (4) \text{ \AA}^3$

$0.24 \times 0.18 \times 0.15 \text{ mm}$

$Z = 4$

Data collection

Bruker SMART APEX CCD diffractometer

5098 independent reflections

Radiation source: fine-focus sealed tube

3508 reflections with $I > 2\sigma(I)$

graphite

$R_{\text{int}} = 0.052$

φ and ω scans

$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 1.8^\circ$

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$h = -15 \rightarrow 15$

$T_{\text{min}} = 0.857$, $T_{\text{max}} = 0.907$

$k = -27 \rightarrow 23$

14540 measured reflections

$l = -11 \rightarrow 10$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.051$

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.153$

H-atom parameters constrained

$S = 0.98$
 $w = 1/[\sigma^2(F_o^2) + (0.0933P)^2 + 0.1872P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 5098 reflections
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 390 parameters
 $\Delta\rho_{\text{max}} = 0.95 \text{ e \AA}^{-3}$
 0 restraints
 $\Delta\rho_{\text{min}} = -0.47 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.28270 (4)	0.56160 (2)	0.90592 (5)	0.02612 (19)
Cl1	0.87327 (8)	0.66511 (5)	0.74420 (12)	0.0455 (3)
C1	0.0861 (3)	0.69236 (18)	0.5203 (4)	0.0376 (10)
H1A	0.0396	0.7024	0.5832	0.045*
C2	0.0804 (4)	0.7195 (2)	0.3979 (5)	0.0449 (12)
H2A	0.0305	0.7487	0.3787	0.054*
C3	0.1472 (4)	0.7040 (2)	0.3036 (5)	0.0511 (13)
H3A	0.1416	0.7227	0.2211	0.061*
C4	0.2219 (4)	0.6615 (2)	0.3292 (4)	0.0433 (11)
H4A	0.2659	0.6508	0.2640	0.052*
C5	0.2313 (3)	0.63511 (19)	0.4515 (4)	0.0350 (10)
H5A	0.2837	0.6073	0.4703	0.042*
C6	0.1631 (3)	0.64933 (18)	0.5489 (4)	0.0307 (9)
C7	0.1786 (3)	0.61984 (17)	0.6799 (4)	0.0275 (9)
C8	0.0525 (3)	0.58652 (18)	0.9627 (4)	0.0310 (9)
H8A	-0.0129	0.6023	0.9359	0.037*
C9	0.0651 (3)	0.55879 (18)	1.0885 (4)	0.0294 (9)
C10	0.1608 (3)	0.53399 (17)	1.1403 (4)	0.0290 (9)
C11	0.1644 (3)	0.50693 (18)	1.2666 (4)	0.0319 (10)
C12	0.0756 (3)	0.5034 (2)	1.3356 (4)	0.0397 (11)
H12A	0.0786	0.4844	1.4177	0.048*
C13	-0.0175 (3)	0.5278 (2)	1.2842 (4)	0.0440 (12)
H13A	-0.0766	0.5256	1.3322	0.053*
C14	-0.0234 (3)	0.5551 (2)	1.1631 (4)	0.0405 (11)
H14A	-0.0867	0.5716	1.1294	0.049*
C15	0.2726 (4)	0.4590 (2)	1.4393 (4)	0.0500 (13)
H15A	0.3446	0.4478	1.4594	0.075*
H15B	0.2288	0.4246	1.4392	0.075*
H15C	0.2522	0.4862	1.5056	0.075*
C16	0.5865 (3)	0.71820 (18)	1.0512 (4)	0.0353 (10)
H16A	0.6324	0.6914	1.0169	0.042*
C17	0.6243 (4)	0.76907 (19)	1.1109 (4)	0.0400 (11)
H17A	0.6961	0.7766	1.1178	0.048*
C18	0.5563 (4)	0.8090 (2)	1.1605 (4)	0.0435 (12)
H18A	0.5822	0.8436	1.2003	0.052*
C19	0.4499 (4)	0.7979 (2)	1.1514 (4)	0.0444 (12)
H19A	0.4043	0.8250	1.1852	0.053*
C20	0.4108 (3)	0.74682 (18)	1.0923 (4)	0.0356 (10)
H20A	0.3389	0.7395	1.0863	0.043*

supplementary materials

C21	0.4784 (3)	0.70675 (17)	1.0423 (4)	0.0294 (9)
C22	0.4344 (3)	0.65140 (17)	0.9850 (4)	0.0257 (9)
C23	0.5028 (3)	0.52154 (17)	0.8316 (4)	0.0257 (9)
H23A	0.5735	0.5297	0.8244	0.031*
C24	0.4618 (3)	0.46776 (17)	0.7770 (4)	0.0267 (9)
C25	0.3558 (3)	0.45109 (17)	0.7842 (4)	0.0269 (9)
C26	0.3226 (3)	0.39603 (18)	0.7277 (4)	0.0303 (9)
C27	0.3928 (3)	0.36044 (19)	0.6688 (4)	0.0351 (10)
H27A	0.3703	0.3246	0.6319	0.042*
C28	0.4968 (3)	0.37752 (19)	0.6638 (4)	0.0366 (10)
H28A	0.5434	0.3527	0.6246	0.044*
C29	0.5312 (3)	0.42949 (18)	0.7147 (4)	0.0327 (10)
H29A	0.6008	0.4404	0.7091	0.039*
C30	0.1794 (4)	0.3313 (2)	0.6782 (6)	0.0551 (14)
H30A	0.1063	0.3280	0.6928	0.083*
H30B	0.2167	0.2981	0.7175	0.083*
H30C	0.1874	0.3321	0.5842	0.083*
N1	0.1015 (3)	0.61943 (14)	0.7608 (3)	0.0307 (8)
H1B	0.0412	0.6352	0.7399	0.037*
N2	0.1262 (2)	0.59143 (14)	0.8822 (3)	0.0260 (7)
N3	0.4973 (2)	0.61077 (13)	0.9384 (3)	0.0277 (8)
H3B	0.5640	0.6158	0.9378	0.033*
N4	0.4475 (2)	0.55953 (13)	0.8906 (3)	0.0231 (7)
O1	0.2638 (2)	0.59602 (12)	0.7150 (3)	0.0307 (6)
O1W	0.7122 (2)	0.60423 (13)	0.9294 (3)	0.0421 (8)
H1WA	0.7368	0.5719	0.9611	0.050*
H1WB	0.7434	0.6240	0.8725	0.050*
O2	0.2468 (2)	0.53545 (13)	1.0763 (3)	0.0346 (7)
O3	0.2611 (2)	0.48627 (13)	1.3112 (3)	0.0396 (7)
O4	0.3375 (2)	0.64162 (12)	0.9791 (3)	0.0318 (7)
O5	0.2873 (2)	0.48311 (12)	0.8409 (3)	0.0323 (7)
O6	0.2204 (2)	0.38391 (13)	0.7375 (3)	0.0423 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0196 (3)	0.0345 (3)	0.0245 (3)	0.0041 (2)	0.0039 (2)	0.0024 (3)
Cl1	0.0289 (6)	0.0524 (7)	0.0550 (8)	0.0074 (5)	0.0009 (5)	0.0078 (6)
C1	0.036 (3)	0.040 (3)	0.036 (3)	0.001 (2)	-0.0045 (19)	0.007 (2)
C2	0.045 (3)	0.045 (3)	0.043 (3)	0.001 (2)	-0.008 (2)	0.013 (2)
C3	0.055 (3)	0.062 (3)	0.034 (3)	-0.011 (3)	-0.010 (2)	0.024 (2)
C4	0.046 (3)	0.055 (3)	0.029 (3)	-0.001 (2)	0.001 (2)	0.008 (2)
C5	0.036 (2)	0.042 (3)	0.027 (2)	0.001 (2)	0.0000 (19)	0.002 (2)
C6	0.029 (2)	0.036 (2)	0.027 (2)	-0.0047 (18)	-0.0044 (18)	0.0020 (18)
C7	0.028 (2)	0.028 (2)	0.026 (2)	-0.0035 (17)	0.0018 (17)	-0.0018 (17)
C8	0.026 (2)	0.041 (2)	0.026 (2)	0.0031 (18)	0.0046 (18)	0.0017 (19)
C9	0.024 (2)	0.038 (2)	0.026 (2)	0.0022 (18)	0.0032 (17)	-0.0015 (19)
C10	0.028 (2)	0.032 (2)	0.028 (2)	0.0005 (17)	0.0064 (17)	-0.0040 (18)

C11	0.036 (2)	0.035 (2)	0.025 (2)	-0.0016 (19)	0.0011 (18)	-0.0016 (19)
C12	0.037 (3)	0.056 (3)	0.026 (2)	-0.003 (2)	0.0081 (19)	0.001 (2)
C13	0.034 (3)	0.068 (3)	0.032 (3)	-0.002 (2)	0.013 (2)	0.005 (2)
C14	0.030 (2)	0.058 (3)	0.034 (3)	0.006 (2)	0.0046 (19)	0.007 (2)
C15	0.056 (3)	0.068 (3)	0.026 (2)	0.006 (3)	0.003 (2)	0.017 (2)
C16	0.033 (2)	0.036 (2)	0.037 (3)	0.0015 (19)	0.0023 (19)	-0.001 (2)
C17	0.043 (3)	0.039 (3)	0.036 (3)	-0.005 (2)	-0.005 (2)	0.002 (2)
C18	0.062 (3)	0.039 (3)	0.028 (2)	-0.006 (2)	-0.003 (2)	-0.009 (2)
C19	0.052 (3)	0.041 (3)	0.040 (3)	0.010 (2)	0.003 (2)	-0.009 (2)
C20	0.040 (3)	0.038 (2)	0.029 (2)	0.006 (2)	0.0011 (19)	-0.0023 (19)
C21	0.032 (2)	0.033 (2)	0.022 (2)	-0.0003 (18)	-0.0045 (17)	0.0024 (18)
C22	0.023 (2)	0.033 (2)	0.021 (2)	0.0052 (17)	-0.0003 (16)	0.0050 (17)
C23	0.020 (2)	0.037 (2)	0.020 (2)	0.0047 (17)	0.0027 (16)	0.0028 (17)
C24	0.022 (2)	0.037 (2)	0.020 (2)	0.0021 (17)	-0.0017 (16)	0.0021 (18)
C25	0.028 (2)	0.033 (2)	0.019 (2)	0.0039 (17)	0.0007 (17)	0.0034 (17)
C26	0.027 (2)	0.037 (2)	0.027 (2)	-0.0001 (18)	-0.0022 (17)	0.0054 (18)
C27	0.043 (3)	0.035 (2)	0.027 (2)	0.002 (2)	-0.0001 (19)	-0.0036 (19)
C28	0.033 (2)	0.043 (3)	0.034 (2)	0.008 (2)	0.0037 (19)	-0.010 (2)
C29	0.025 (2)	0.041 (3)	0.032 (2)	0.0027 (18)	0.0034 (18)	-0.0056 (19)
C30	0.041 (3)	0.047 (3)	0.077 (4)	-0.013 (2)	-0.001 (3)	-0.012 (3)
N1	0.0245 (18)	0.040 (2)	0.0274 (19)	0.0083 (15)	0.0023 (14)	0.0098 (15)
N2	0.0231 (17)	0.0344 (19)	0.0201 (17)	0.0038 (14)	-0.0001 (14)	0.0063 (14)
N3	0.0231 (17)	0.0315 (18)	0.0281 (19)	-0.0025 (14)	0.0002 (14)	-0.0003 (15)
N4	0.0208 (16)	0.0286 (17)	0.0197 (16)	0.0011 (14)	-0.0008 (13)	-0.0002 (14)
O1	0.0250 (15)	0.0424 (17)	0.0251 (15)	0.0053 (13)	0.0053 (12)	0.0069 (13)
O1W	0.0332 (17)	0.0456 (19)	0.0476 (19)	0.0050 (14)	0.0044 (14)	0.0046 (15)
O2	0.0249 (15)	0.0553 (19)	0.0242 (15)	0.0088 (13)	0.0043 (12)	0.0116 (13)
O3	0.0346 (17)	0.059 (2)	0.0248 (16)	0.0076 (15)	0.0022 (13)	0.0145 (14)
O4	0.0279 (16)	0.0359 (16)	0.0318 (16)	0.0046 (12)	0.0033 (12)	-0.0048 (13)
O5	0.0235 (15)	0.0342 (16)	0.0399 (17)	0.0001 (12)	0.0073 (12)	-0.0046 (13)
O6	0.0303 (17)	0.0440 (18)	0.053 (2)	-0.0081 (14)	0.0050 (14)	-0.0108 (15)

Geometric parameters (\AA , $^\circ$)

Fe1—O1	2.070 (3)	C16—C17	1.372 (6)
Fe1—O2	1.904 (3)	C16—C21	1.402 (6)
Fe1—O4	2.062 (3)	C16—H16A	0.9300
Fe1—O5	1.901 (3)	C17—C18	1.377 (6)
Fe1—N2	2.106 (3)	C17—H17A	0.9300
Fe1—N4	2.124 (3)	C18—C19	1.379 (7)
C1—C2	1.373 (6)	C18—H18A	0.9300
C1—C6	1.401 (6)	C19—C20	1.379 (6)
C1—H1A	0.9300	C19—H19A	0.9300
C2—C3	1.372 (7)	C20—C21	1.377 (5)
C2—H2A	0.9300	C20—H20A	0.9300
C3—C4	1.367 (6)	C21—C22	1.476 (5)
C3—H3A	0.9300	C22—O4	1.254 (4)
C4—C5	1.365 (6)	C22—N3	1.334 (5)
C4—H4A	0.9300	C23—N4	1.289 (5)

supplementary materials

C5—C6	1.399 (6)	C23—C24	1.422 (5)
C5—H5A	0.9300	C23—H23A	0.9300
C6—C7	1.478 (5)	C24—C25	1.413 (5)
C7—O1	1.241 (5)	C24—C29	1.422 (5)
C7—N1	1.328 (5)	C25—O5	1.303 (4)
C8—N2	1.294 (4)	C25—C26	1.425 (5)
C8—C9	1.411 (5)	C26—O6	1.346 (5)
C8—H8A	0.9300	C26—C27	1.375 (6)
C9—C10	1.408 (5)	C27—C28	1.389 (6)
C9—C14	1.409 (5)	C27—H27A	0.9300
C10—O2	1.317 (4)	C28—C29	1.347 (6)
C10—C11	1.409 (5)	C28—H28A	0.9300
C11—O3	1.364 (5)	C29—H29A	0.9300
C11—C12	1.379 (5)	C30—O6	1.417 (5)
C12—C13	1.375 (6)	C30—H30A	0.9600
C12—H12A	0.9300	C30—H30B	0.9600
C13—C14	1.364 (6)	C30—H30C	0.9600
C13—H13A	0.9300	N1—N2	1.391 (4)
C14—H14A	0.9300	N1—H1B	0.8600
C15—O3	1.428 (5)	N3—N4	1.394 (4)
C15—H15A	0.9600	N3—H3B	0.8600
C15—H15B	0.9600	O1W—H1WA	0.8500
C15—H15C	0.9600	O1W—H1WB	0.8502
O5—Fe1—O2	91.94 (12)	C21—C16—H16A	120.1
O5—Fe1—O4	158.43 (11)	C16—C17—C18	120.2 (4)
O2—Fe1—O4	93.00 (12)	C16—C17—H17A	119.9
O5—Fe1—O1	92.26 (11)	C18—C17—H17A	119.9
O2—Fe1—O1	159.12 (11)	C17—C18—C19	120.2 (4)
O4—Fe1—O1	90.57 (11)	C17—C18—H18A	119.9
O5—Fe1—N2	108.53 (12)	C19—C18—H18A	119.9
O2—Fe1—N2	84.74 (11)	C18—C19—C20	120.3 (4)
O4—Fe1—N2	92.83 (11)	C18—C19—H19A	119.9
O1—Fe1—N2	74.54 (11)	C20—C19—H19A	119.9
O5—Fe1—N4	84.04 (11)	C21—C20—C19	119.8 (4)
O2—Fe1—N4	111.65 (11)	C21—C20—H20A	120.1
O4—Fe1—N4	74.63 (11)	C19—C20—H20A	120.1
O1—Fe1—N4	89.13 (11)	C20—C21—C16	119.9 (4)
N2—Fe1—N4	159.46 (12)	C20—C21—C22	118.4 (4)
C2—C1—C6	118.9 (4)	C16—C21—C22	121.7 (4)
C2—C1—H1A	120.5	O4—C22—N3	118.8 (4)
C6—C1—H1A	120.5	O4—C22—C21	120.9 (3)
C3—C2—C1	120.9 (4)	N3—C22—C21	120.3 (3)
C3—C2—H2A	119.6	N4—C23—C24	123.6 (3)
C1—C2—H2A	119.6	N4—C23—H23A	118.2
C4—C3—C2	121.0 (4)	C24—C23—H23A	118.2
C4—C3—H3A	119.5	C25—C24—C29	119.5 (4)
C2—C3—H3A	119.5	C25—C24—C23	122.4 (3)
C5—C4—C3	119.3 (4)	C29—C24—C23	118.1 (3)
C5—C4—H4A	120.3	O5—C25—C24	123.5 (4)

C3—C4—H4A	120.3	O5—C25—C26	118.4 (3)
C4—C5—C6	120.9 (4)	C24—C25—C26	118.1 (3)
C4—C5—H5A	119.5	O6—C26—C27	125.5 (4)
C6—C5—H5A	119.5	O6—C26—C25	114.2 (3)
C5—C6—C1	118.9 (4)	C27—C26—C25	120.3 (4)
C5—C6—C7	118.2 (4)	C26—C27—C28	120.6 (4)
C1—C6—C7	122.8 (4)	C26—C27—H27A	119.7
O1—C7—N1	120.0 (4)	C28—C27—H27A	119.7
O1—C7—C6	120.3 (3)	C29—C28—C27	121.0 (4)
N1—C7—C6	119.8 (4)	C29—C28—H28A	119.5
N2—C8—C9	124.2 (4)	C27—C28—H28A	119.5
N2—C8—H8A	117.9	C28—C29—C24	120.5 (4)
C9—C8—H8A	117.9	C28—C29—H29A	119.8
C10—C9—C14	119.3 (4)	C24—C29—H29A	119.8
C10—C9—C8	123.0 (3)	O6—C30—H30A	109.5
C14—C9—C8	117.7 (4)	O6—C30—H30B	109.5
O2—C10—C9	122.9 (3)	H30A—C30—H30B	109.5
O2—C10—C11	118.8 (4)	O6—C30—H30C	109.5
C9—C10—C11	118.3 (3)	H30A—C30—H30C	109.5
O3—C11—C12	125.1 (4)	H30B—C30—H30C	109.5
O3—C11—C10	114.3 (3)	C7—N1—N2	114.4 (3)
C12—C11—C10	120.6 (4)	C7—N1—H1B	122.8
C13—C12—C11	120.6 (4)	N2—N1—H1B	122.8
C13—C12—H12A	119.7	C8—N2—N1	117.5 (3)
C11—C12—H12A	119.7	C8—N2—Fe1	129.2 (3)
C14—C13—C12	120.3 (4)	N1—N2—Fe1	113.2 (2)
C14—C13—H13A	119.8	C22—N3—N4	115.3 (3)
C12—C13—H13A	119.8	C22—N3—H3B	122.4
C13—C14—C9	120.8 (4)	N4—N3—H3B	122.4
C13—C14—H14A	119.6	C23—N4—N3	117.7 (3)
C9—C14—H14A	119.6	C23—N4—Fe1	129.1 (3)
O3—C15—H15A	109.5	N3—N4—Fe1	112.6 (2)
O3—C15—H15B	109.5	C7—O1—Fe1	117.7 (2)
H15A—C15—H15B	109.5	H1WA—O1W—H1WB	121.9
O3—C15—H15C	109.5	C10—O2—Fe1	135.8 (3)
H15A—C15—H15C	109.5	C11—O3—C15	118.2 (3)
H15B—C15—H15C	109.5	C22—O4—Fe1	118.6 (2)
C17—C16—C21	119.7 (4)	C25—O5—Fe1	135.6 (2)
C17—C16—H16A	120.1	C26—O6—C30	118.0 (3)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1B···C11 ⁱ	0.86	2.25	3.087 (3)	163
N3—H3B···O1W	0.86	1.92	2.759 (4)	164
O1W—H1WA···O5 ⁱⁱ	0.85	2.39	3.045 (4)	134
O1W—H1WB···C11	0.85	2.37	3.198 (3)	163

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1, -y+1, -z+2$.

supplementary materials

Fig. 1




Fig. 2

