

1-Cyanomethyl-4-aza-1-azoniabicyclo-[2.2.2]octane bromide dihydrate

Ying Cai

Ordered Matter Science Research Center, Southeast University, Nanjing 211189, People's Republic of China
Correspondence e-mail: cyik@163.com


Received 25 May 2010; accepted 26 May 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; disorder in main residue; R factor = 0.041; wR factor = 0.077; data-to-parameter ratio = 19.4.

In the crystal structure of the title compound, $\text{C}_8\text{H}_{14}\text{N}_3^+\cdot\text{Br}^-\cdot 2\text{H}_2\text{O}$, intermolecular $\text{O}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{Br}$ hydrogen bonding occurs. The water molecules are connected into chains extending in the a -axis direction. The bromide anions are connected to the water molecules, forming 10-membered rings. The cations are connected to the anions via weak $\text{C}-\text{H}\cdots\text{Br}$ interactions. Two carbon atoms of the cation are disordered and were refined using a split model (occupancy ratio 0.70:0.3).

Related literature

For uses of DABCO (1,4-biazabicyclo[2.2.2]octane) and its derivatives, see: Basaviah *et al.* (2003); Chen *et al.* (2010).



Experimental

Crystal data

$\text{C}_8\text{H}_{14}\text{N}_3^+\cdot\text{Br}^-\cdot 2\text{H}_2\text{O}$
 $M_r = 268.16$
Orthorhombic, $P2_12_12_1$
 $a = 7.461 (5)\text{ \AA}$
 $b = 12.008 (7)\text{ \AA}$
 $c = 13.236 (8)\text{ \AA}$

$V = 1185.8 (13)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 3.45\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.20 \times 0.20 \times 0.20\text{ mm}$

Data collection

Rigaku Mercury CCD diffractometer
Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.701$, $T_{\max} = 1.000$

13047 measured reflections
2711 independent reflections
2219 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.073$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.077$
 $S = 1.01$
2711 reflections
140 parameters
101 restraints

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.29\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
1134 Friedel pairs
Flack parameter: 0.033 (14)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1O1 \cdots Br1	0.82	2.58	3.354 (3)	159
O2—H1O2 \cdots O1	0.82	1.98	2.791 (4)	170
O1—H2O1 \cdots O2 ⁱ	0.82	1.99	2.788 (5)	164
O2—H2O2 \cdots Br1 ⁱ	0.82	2.50	3.314 (3)	172
C7—H7A \cdots Br1	0.97	2.81	3.740 (5)	161
C7—H7B \cdots Br1 ⁱⁱ	0.97	2.92	3.792 (8)	151

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, -z + 1$; (ii) $-x + \frac{1}{2}, -y, z + \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NC2186).

References

- Basaviah, D., Rao, A. J. & Satyanarayana, T. (2003). *Chem. Rev.* **103**, 811–891.
- Chen, L. Z., Huang, Y., Xiong, R. G. & Hu, H. W. (2010). *J. Mol. Struct.* **963**, 16–21.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, o1527 [doi:10.1107/S1600536810019926]

1-Cyanomethyl-4-aza-1-azoniabicyclo[2.2.2]octane bromide dihydrate

Y. Cai

Comment

1,4-Diazabicyclo[2.2.2]octane (DABCO) is used as a organocatalyst for a large number of reactions because of its nucleophilicity (Basaviah *et al.*, 2003) and some of its derivatives are ferroelectrics (Chen *et al.*, 2010). The structure determination of the title compound was performed within a project on the electric properties of 1,4-Diazabicyclo[2.2.2]octane derivatives. Within this project the crystals of the title compound were obtained by accident.

In the crystal stucture of the title compound two C atoms of the cation are disordered (Fig. 1). The cations and anions are connected by weak intermolecular C—H···Br interactions. The bromide anions are additionally linked to the water molecules *via* intermolecular O—H···Br hydrogen bonding and the water molecules are connected into chains that elongate in the direction of the α axis (Fig. 2). Each water molecule act as hydrogen bond donor and acceptor. The bromide anions and the water molecules forming ten-membered rings.

Experimental

1,4-Diaza-bicyclo[2.2.2]octane (dabco) (0.05?mol, 5.6?g) and bromoacetonitrile (0.1?mol, 12.00?g) were dissolved in CH₃CN(40?ml). The mixture was stirred for 1?h leading to a white precipitate of the title compound which was filtered off, washed with acetonitrile and dried. Yield: 80%. Afterwards a mixture of 1-(cyanomethyl)-4-aza-1-azoniabicyclo[2.2.2]octane bromide (0.01?mol 2.32?g) and tetrafluoro-borate sodium (0.01?mol 1.10?g) in H₂O (20?ml) was stirred until a clear solution was obtained. On slow evaporation of the solvent colourless plate crystals of the title compand suitable for X-ray analysis were obtained accidentally.

The dielectric constant of the title compound as a function of temperature goes smoothly between 93 and 363?K and there is no dielectric anomaly observed within the measured temperature range.

Refinement

The C—H H atoms were positioned with idealized geometry and refined using a riding model ($U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$). The O—H H atoms were located in difference map, their bond lengths set to ideal values and finally they were refined using a riding model ($U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$). Two carbon atoms are disordered and were refined using a split model and sof of 0.7 and 0.3. The C atoms with lower occupancy were refined only isotropic. The absolute structure was determined on the basis of 1134 Friedel-pairs.

supplementary materials

Figures




Fig. 1. Crystal structure of the title compound with labelling and displacement ellipsoids drawn at the 30% probability level. Disordering is shown with full and open bonds.




Fig. 2. Crystal structure of the title compound with view along the a axis. Disordered C and H atoms are omitted and intermolecular hydrogen bonding is shown as dashed lines.

1-Cyanomethyl-4-aza-1-azoniabicyclo[2.2.2]octane bromide dihydrate

Crystal data

$C_8H_{14}N_3^+\cdot Br^- \cdot 2H_2O$	$F(000) = 552$
$M_r = 268.16$	$D_x = 1.502 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 3350 reflections
$a = 7.461 (5) \text{ \AA}$	$\theta = 6.3\text{--}55.2^\circ$
$b = 12.008 (7) \text{ \AA}$	$\mu = 3.45 \text{ mm}^{-1}$
$c = 13.236 (8) \text{ \AA}$	$T = 293 \text{ K}$
$V = 1185.8 (13) \text{ \AA}^3$	Prism, colourless
$Z = 4$	$0.20 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Rigaku Mercury CCD diffractometer	2711 independent reflections
Radiation source: fine-focus sealed tube graphite	2219 reflections with $I > 2\sigma(I)$
Detector resolution: 13.6620 pixels mm^{-1}	$R_{\text{int}} = 0.073$
ω scans	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.701, T_{\text{max}} = 1.000$	$k = -15 \rightarrow 15$
13047 measured reflections	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.041$	$w = 1/[\sigma^2(F_o^2) + (0.0275P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.077$	$(\Delta/\sigma)_{\max} < 0.001$
$S = 1.01$	$\Delta\rho_{\max} = 0.29 \text{ e \AA}^{-3}$
2711 reflections	$\Delta\rho_{\min} = -0.29 \text{ e \AA}^{-3}$
140 parameters	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$
101 restraints	Extinction coefficient: 0.0055 (11)
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 1134 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: 0.033 (14)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
N1	0.3080 (4)	0.2169 (2)	0.79551 (19)	0.0332 (7)	
C1	0.3169 (6)	0.2800 (3)	0.8942 (3)	0.0470 (10)	
H1A	0.3725	0.2343	0.9458	0.056*	
H1B	0.1972	0.2997	0.9165	0.056*	
C2	0.4287 (6)	0.3857 (3)	0.8761 (3)	0.0623 (11)	
H2A	0.3617	0.4499	0.8995	0.075*	
H2B	0.5381	0.3812	0.9154	0.075*	
C3	0.2110 (6)	0.2873 (3)	0.7178 (3)	0.0640 (13)	
H3A	0.0839	0.2899	0.7330	0.077*	0.70
H3B	0.2262	0.2556	0.6509	0.077*	0.70
H3C	0.1046	0.3208	0.7470	0.077*	0.30
H3D	0.1755	0.2422	0.6604	0.077*	0.30
C4	0.2926 (10)	0.4068 (6)	0.7216 (5)	0.053 (2)	0.70
H4A	0.3034	0.4364	0.6536	0.063*	0.70
H4B	0.2148	0.4557	0.7600	0.063*	0.70

supplementary materials

C4'	0.349 (2)	0.3801 (15)	0.6842 (12)	0.078 (8)	0.30
H4C	0.4151	0.3555	0.6253	0.093*	0.30
H4D	0.2852	0.4480	0.6669	0.093*	0.30
C5	0.4948 (6)	0.1953 (3)	0.7609 (4)	0.0645 (12)	
H5A	0.4948	0.1434	0.7048	0.077*	0.70
H5B	0.5654	0.1638	0.8154	0.077*	0.70
H5C	0.4924	0.1661	0.6926	0.077*	0.30
H5D	0.5497	0.1398	0.8042	0.077*	0.30
C6	0.5746 (11)	0.3096 (5)	0.7274 (8)	0.0645 (12)	0.70
H6A	0.6987	0.3144	0.7489	0.077*	0.70
H6B	0.5718	0.3148	0.6543	0.077*	0.70
C6'	0.6115 (17)	0.3066 (10)	0.7634 (12)	0.078 (8)	0.30
H6C	0.6908	0.3071	0.8215	0.094*	0.30
H6D	0.6830	0.3135	0.7025	0.094*	0.30
N2	0.4744 (4)	0.4012 (3)	0.7705 (3)	0.0493 (8)	
C7	0.2164 (5)	0.1068 (3)	0.8091 (3)	0.0435 (9)	
H7A	0.2139	0.0676	0.7450	0.052*	
H7B	0.2836	0.0619	0.8568	0.052*	
C8	0.0335 (6)	0.1213 (3)	0.8459 (3)	0.0488 (10)	
N3	-0.1082 (6)	0.1335 (3)	0.8753 (3)	0.0724 (11)	
Br1	0.08721 (5)	-0.01932 (3)	0.56114 (3)	0.05169 (15)	
O1	0.4108 (4)	0.1366 (2)	0.4565 (2)	0.0779 (9)	
H1O1	0.3519	0.0980	0.4953	0.117*	
H2O1	0.4970	0.1418	0.4946	0.117*	
O2	0.2440 (4)	0.3445 (2)	0.4497 (2)	0.0736 (9)	
H1O2	0.2887	0.2823	0.4448	0.110*	
H2O2	0.3207	0.3932	0.4458	0.110*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0371 (15)	0.0258 (15)	0.0368 (16)	0.0016 (13)	-0.0012 (12)	-0.0057 (12)
C1	0.061 (3)	0.042 (2)	0.038 (2)	-0.0100 (19)	0.0000 (18)	-0.0081 (16)
C2	0.063 (3)	0.049 (2)	0.074 (3)	-0.022 (3)	0.012 (3)	-0.017 (2)
C3	0.079 (3)	0.056 (3)	0.057 (2)	-0.013 (3)	-0.027 (2)	0.021 (2)
C4	0.070 (4)	0.044 (4)	0.044 (4)	0.010 (3)	0.003 (3)	0.012 (3)
C4'	0.12 (2)	0.036 (10)	0.081 (15)	-0.034 (12)	-0.028 (13)	0.018 (10)
C5	0.051 (2)	0.041 (2)	0.101 (3)	0.0023 (19)	0.032 (2)	-0.009 (2)
C6	0.051 (2)	0.041 (2)	0.101 (3)	0.0023 (19)	0.032 (2)	-0.009 (2)
C6'	0.076 (13)	0.085 (13)	0.072 (13)	-0.067 (12)	0.050 (10)	-0.046 (10)
N2	0.0519 (19)	0.0353 (17)	0.061 (2)	-0.0021 (15)	0.0030 (16)	0.0087 (16)
C7	0.051 (2)	0.0298 (19)	0.049 (2)	-0.0041 (18)	0.0025 (19)	0.0010 (16)
C8	0.052 (3)	0.042 (2)	0.052 (2)	-0.0124 (19)	-0.0037 (19)	-0.0013 (18)
N3	0.058 (3)	0.072 (3)	0.087 (3)	-0.014 (2)	0.004 (2)	-0.012 (2)
Br1	0.0441 (2)	0.0516 (2)	0.0595 (2)	0.00308 (19)	-0.00010 (19)	-0.00776 (18)
O1	0.0656 (19)	0.0689 (19)	0.099 (2)	0.0058 (19)	-0.008 (2)	0.0238 (17)
O2	0.0561 (18)	0.0486 (17)	0.116 (3)	-0.0053 (15)	-0.009 (2)	0.0014 (18)

Geometric parameters (Å, °)

N1—C5	1.490 (5)	C4'—H4D	0.9700
N1—C7	1.499 (4)	C5—C6	1.560 (8)
N1—C1	1.511 (4)	C5—C6'	1.595 (12)
N1—C3	1.515 (5)	C5—H5A	0.9700
C1—C2	1.538 (5)	C5—H5B	0.9700
C1—H1A	0.9700	C5—H5C	0.9700
C1—H1B	0.9700	C5—H5D	0.9700
C2—N2	1.451 (5)	C6—N2	1.448 (7)
C2—H2A	0.9700	C6—H6A	0.9700
C2—H2B	0.9700	C6—H6B	0.9700
C3—C4	1.560 (8)	C6'—N2	1.532 (13)
C3—C4'	1.579 (14)	C6'—H6C	0.9700
C3—H3A	0.9700	C6'—H6D	0.9700
C3—H3B	0.9700	C7—C8	1.459 (6)
C3—H3C	0.9700	C7—H7A	0.9700
C3—H3D	0.9700	C7—H7B	0.9700
C4—N2	1.504 (8)	C8—N3	1.136 (5)
C4—H4A	0.9700	O1—H1O1	0.8201
C4—H4B	0.9700	O1—H2O1	0.8200
C4'—N2	1.500 (14)	O2—H1O2	0.8201
C4'—H4C	0.9700	O2—H2O2	0.8200
C5—N1—C7	108.0 (3)	N1—C5—C6'	111.0 (5)
C5—N1—C1	108.2 (3)	C6—C5—C6'	20.1 (9)
C7—N1—C1	111.0 (3)	N1—C5—H5A	110.3
C5—N1—C3	109.6 (3)	C6—C5—H5A	110.3
C7—N1—C3	110.8 (3)	C6'—C5—H5A	123.7
C1—N1—C3	109.2 (3)	N1—C5—H5B	110.3
N1—C1—C2	107.6 (3)	C6—C5—H5B	110.3
N1—C1—H1A	110.2	C6'—C5—H5B	90.9
C2—C1—H1A	110.2	H5A—C5—H5B	108.6
N1—C1—H1B	110.2	N1—C5—H5C	109.4
C2—C1—H1B	110.2	C6—C5—H5C	93.5
H1A—C1—H1B	108.5	C6'—C5—H5C	109.4
N2—C2—C1	112.6 (3)	H5A—C5—H5C	18.8
N2—C2—H2A	109.1	H5B—C5—H5C	124.1
C1—C2—H2A	109.1	N1—C5—H5D	109.4
N2—C2—H2B	109.1	C6—C5—H5D	127.7
C1—C2—H2B	109.1	C6'—C5—H5D	109.4
H2A—C2—H2B	107.8	H5A—C5—H5D	90.6
N1—C3—C4	107.7 (4)	H5B—C5—H5D	20.4
N1—C3—C4'	105.9 (7)	H5C—C5—H5D	108.0
C4—C3—C4'	26.6 (8)	N2—C6—C5	111.1 (5)
N1—C3—H3A	110.2	N2—C6—H6A	109.4
C4—C3—H3A	110.2	C5—C6—H6A	109.4
C4'—C3—H3A	132.1	N2—C6—H6B	109.4
N1—C3—H3B	110.2	C5—C6—H6B	109.4

supplementary materials


C4—C3—H3B	110.2	H6A—C6—H6B	108.0
C4'—C3—H3B	86.8	N2—C6'—C5	104.9 (8)
H3A—C3—H3B	108.5	N2—C6'—H6C	110.8
N1—C3—H3C	110.6	C5—C6'—H6C	110.8
C4—C3—H3C	85.7	N2—C6'—H6D	110.8
C4'—C3—H3C	110.6	C5—C6'—H6D	110.8
H3A—C3—H3C	26.3	H6C—C6'—H6D	108.8
H3B—C3—H3C	128.5	C6—N2—C2	113.8 (5)
N1—C3—H3D	110.6	C6—N2—C4'	84.0 (10)
C4—C3—H3D	130.2	C2—N2—C4'	124.4 (7)
C4'—C3—H3D	110.6	C6—N2—C4	109.2 (5)
H3A—C3—H3D	85.1	C2—N2—C4	102.0 (4)
H3B—C3—H3D	25.6	C4'—N2—C4	27.8 (9)
H3C—C3—H3D	108.7	C6—N2—C6'	21.1 (8)
N2—C4—C3	109.0 (5)	C2—N2—C6'	96.9 (6)
N2—C4—H4A	109.9	C4'—N2—C6'	104.3 (10)
C3—C4—H4A	109.9	C4—N2—C6'	127.5 (6)
N2—C4—H4B	109.9	C8—C7—N1	111.2 (3)
C3—C4—H4B	109.9	C8—C7—H7A	109.4
H4A—C4—H4B	108.3	N1—C7—H7A	109.4
N2—C4'—C3	108.2 (9)	C8—C7—H7B	109.4
N2—C4'—H4C	110.1	N1—C7—H7B	109.4
C3—C4'—H4C	110.1	H7A—C7—H7B	108.0
N2—C4'—H4D	110.1	N3—C8—C7	179.2 (5)
C3—C4'—H4D	110.1	H1O1—O1—H2O1	94.5
H4C—C4'—H4D	108.4	H1O2—O2—H2O2	111.1
N1—C5—C6	106.9 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H1O1···Br1	0.82	2.58	3.354 (3)	159.
O2—H1O2···O1	0.82	1.98	2.791 (4)	170.
O1—H2O1···O2 ⁱ	0.82	1.99	2.788 (5)	164.
O2—H2O2···Br1 ⁱ	0.82	2.50	3.314 (3)	172.
C7—H7A···Br1	0.97	2.81	3.740 (5)	161
C7—H7B···Br1 ⁱⁱ	0.97	2.92	3.792 (8)	151

Symmetry codes: (i) $x+1/2, -y+1/2, -z+1$; (ii) $-x+1/2, -y, z+1/2$.

Fig. 1



supplementary materials

Fig. 2

