

9-Benzyl-10-methylacridinium trifluoromethanesulfonate

Damian Trzybiński, Beata Zadykiewicz, Karol Krzywiński, Artur Sikorski and Jerzy Błażejowski*

Faculty of Chemistry, University of Gdańsk, J. Sobieskiego 18, 80-952 Gdańsk, Poland

Correspondence e-mail: bla@chem.univ.gda.pl


Received 18 May 2010; accepted 25 May 2010

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.052; wR factor = 0.153; data-to-parameter ratio = 13.0.

In the crystal structure of the title compound, $\text{C}_{21}\text{H}_{18}\text{N}^+\cdot\text{CF}_3\text{OS}_3^-$, the cations form inversion dimers through $\pi-\pi$ interactions between the acridine ring systems. These dimers are further linked by $\text{C}-\text{H}\cdots\pi$ interactions. The cations and anions are connected by $\text{C}-\text{H}\cdots\text{O}$, $\text{C}-\text{F}\cdots\pi$ and $\text{S}-\text{O}\cdots\pi$ interactions. The acridine and benzene ring systems are oriented at a dihedral angle of $76.8(1)^\circ$ with respect to each other. The acridine moieties are either parallel or inclined at an angle of $62.4(1)^\circ$ in the crystal structure.

Related literature

For general background to acridinium derivatives, see: King *et al.* (2007); Roda *et al.* (2003); Wróblewska *et al.* (2004); Trzybiński *et al.* (2010); Zomer & Jacquemijns (2001). For related structures, see: Sikorski *et al.* (2007); Trzybiński *et al.* (2010). For intermolecular interactions, see: Bianchi *et al.* (2004); Dorn *et al.* (2005); Hunter *et al.* (2001); Novoa *et al.* (2006); Takahashi *et al.* (2001). For the synthesis, see: Huntress & Shaw (1948); Sikorski *et al.* (2007); Trzybiński *et al.* (2010).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{18}\text{N}^+\cdot\text{CF}_3\text{O}_3\text{S}^-$
 $M_r = 433.44$
 Monoclinic, $P2_1/n$
 $a = 14.6211(7)$ Å
 $b = 8.2514(2)$ Å

$c = 17.2900(8)$ Å
 $\beta = 107.707(5)^\circ$
 $V = 1987.12(15)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.22$ mm⁻¹
 $T = 295$ K

$0.41 \times 0.25 \times 0.08$ mm

Data collection

Oxford Diffraction Gemini R Ultra Ruby CCD diffractometer
 Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2008)
 $T_{\min} = 0.953$, $T_{\max} = 0.988$

16520 measured reflections
 3528 independent reflections
 2191 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.153$
 $S = 1.06$
 3528 reflections

272 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.37$ e Å⁻³
 $\Delta\rho_{\min} = -0.28$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg4 is the centroid of the C16–C21 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C2}-\text{H2}\cdots\text{O26}^{\text{i}}$	0.93	2.49	3.398 (5)	167
$\text{C3}-\text{H3}\cdots\text{Cg4}^{\text{ii}}$	0.93	2.74	3.630 (5)	161
$\text{C15}-\text{H15B}\cdots\text{O25}^{\text{iii}}$	0.97	2.49	3.423 (4)	160
$\text{C22}-\text{H22B}\cdots\text{O25}^{\text{iv}}$	0.96	2.56	3.386 (5)	144
$\text{C22}-\text{H22C}\cdots\text{O24}$	0.96	2.56	3.361 (5)	141

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x + 2, -y, -z + 1$; (iii) $-x + 2, -y + 1, -z + 1$; (iv) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Table 2

$\text{C}-\text{F}\cdots\pi$ and $\text{S}-\text{O}\cdots\pi$ interactions (Å, °).

Cg1 and Cg3 are the centroids of the C9/N10/C11–C14 and C5–C8/C13/C14 rings, respectively.

X	I	J	$I\cdots J$	$X\cdots J$	$X-I\cdots J$
C27	F30	Cg3^{v}	3.115 (3)	4.233 (3)	143.0 (2)
S23	O26	Cg1^{v}	3.085 (3)	4.167 (2)	131.4 (2)

Symmetry code: (v) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Table 3

$\pi-\pi$ interactions (Å, °).

Cg1 and Cg2 are the centroids of the C9/N10/C11–C14 and C1–C4/C11/C12 rings, respectively. $\text{CgI}\cdots\text{CgJ}$ is the distance between ring centroids. The dihedral angle is that between the planes of the rings I and J . CgI_{Perp} is the perpendicular distance of CgI from ring J . $\text{CgI}_{\text{Offset}}$ is the distance between CgI and perpendicular projection of CgJ on ring I .

I	J	$\text{CgI}\cdots\text{CgJ}$	Dihedral angle	CgI_{Perp}	$\text{CgI}_{\text{Offset}}$
1	2 ⁱⁱⁱ	3.806 (2)	2.11 (15)	3.575 (2)	1.306 (2)
2	1 ⁱⁱⁱ	3.806 (2)	2.11 (15)	3.530 (2)	1.423 (2)
2	2 ⁱⁱⁱ	3.886 (2)	0.02 (15)	3.563 (2)	1.551 (2)

Symmetry code: (iii) $-x + 2, -y + 1, -z + 1$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2008); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2008); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

This study was financed by the State Funds for Scientific Research (grant DS No. 8220-4-0087-9).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2343).

References

- Bianchi, R., Forni, A. & Pilati, T. (2004). *Acta Cryst.* **B60**, 559–568.
- Dorn, T., Janiak, C. & Abu-Shandi, K. (2005). *CrystEngComm*, **7**, 633–641.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hunter, C. A., Lawson, K. R., Perkins, J. & Urch, C. J. (2001). *J. Chem. Soc. Perkin Trans. 2*, pp. 651–669.
- Huntress, E. H. & Shaw, E. N. (1948). *J. Org. Chem.* **13**, 674–675.
- King, D. W., Cooper, W. J., Rusak, S. A., Peake, B. M., Kiddle, J. J., O'Sullivan, D. W., Melamed, M. L., Morgan, C. R. & Theberge, S. M. (2007). *Anal. Chem.* **79**, 4169–4176.
- Novoa, J. J., Mota, F. & D'Oria, E. (2006). *Hydrogen Bonding – New Insights*, edited by S. Grabowski, pp. 193–244. The Netherlands: Springer.
- Oxford Diffraction (2008). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Roda, A., Guardigli, M., Michelini, E., Mirasoli, M. & Pasini, P. (2003). *Anal. Chem.* **A75**, 462–470.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sikorski, A., Kowalska, K., Krzyński, K. & Błażejowski, J. (2007). *Acta Cryst.* **E63**, o2670–o2672.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Takahashi, O., Kohno, Y., Iwasaki, S., Saito, K., Iwaoka, M., Tomada, S., Umezawa, Y., Tsuboyama, S. & Nishio, M. (2001). *Bull. Chem. Soc. Jpn*, **74**, 2421–2430.
- Trzybiński, D., Krzyński, K., Sikorski, A., Malecha, P. & Błażejowski, J. (2010). *Acta Cryst.* **E66**, o826–o827.
- Wróblewska, A., Huta, O. M., Midyanyj, S. V., Patsay, I. O., Rak, J. & Błażejowski, J. (2004). *J. Org. Chem.* **69**, 1607–1614.
- Zomer, G. & Jacquemijns, M. (2001). *Chemiluminescence in Analytical Chemistry*, edited by A. M. Garcia-Campana & W. R. G. Baeyens, pp. 529–549. New York: Marcel Dekker.

supplementary materials

Acta Cryst. (2010). E66, o1548-o1549 [doi:10.1107/S160053681001963X]

9-Benzyl-10-methylacridinium trifluoromethanesulfonate

D. Trzybinski, B. Zadykowicz, K. Krzyminski, A. Sikorski and J. Blazejowski

Comment

Quaternary *N*-methylacridinium cations substituted in position 9 undergo oxidation with H₂O₂ or other oxidants in alkaline media accompanied by chemiluminescence (Zomer & Jacquemijns, 2001). The emission that originates from electronically excited 10-methyl-9-acridinone, the oxidation product, is affected by the features of the substituent in position 9. For these reasons various acridine derivatives of the above type have been synthesized and investigated from the point of view of their chemiluminogenic ability and applicability in immunological, biological, chemical and environmental analyses (Zomer & Jacquemijns, 2001; Roda *et al.*, 2003; King *et al.*, 2007). Continuing the search for 9-substituted acridinium derivatives with chemiluminogenic potential (Wróblewska *et al.*, 2004; Trzybiński *et al.*, 2010), we synthesized 9-benzyl-10-methylacridinium trifluoromethanesulfonate whose crystal structure is presented here.

In the crystal structure, the inversely oriented cations form dimers through multidirectional π - π interactions involving acridine moieties (Table 3, Fig. 2). These dimers are linked by C-H \cdots O (Table 1, Figs. 1 and 2), C-F \cdots π (acridine) (Table 2, Fig. 2) and S-O \cdots π (acridine) (Table 2, Fig. 2) interactions with adjacent anions, and by C-H \cdots π (phenyl) (Table 1, Fig. 2) interactions with neighboring cations. The C-H \cdots O interactions are of the hydrogen bond type (Bianchi *et al.*, 2004; Novoa *et al.*, 2006). The C-H \cdots π interactions should be of an attractive nature (Takahashi *et al.*, 2001), like the C-F \cdots π (Dorn *et al.*, 2005), S-O \cdots π (Dorn *et al.*, 2005) and the π - π (Hunter *et al.*, 2001) interactions. The crystal structure is stabilized by a network of these short-range specific interactions and by long-range electrostatic interactions between ions.

In the cation of the title compound (Fig. 1), the bond lengths and angles characterizing the geometry of the acridinium moiety are typical of acridine-based derivatives (Sikorski *et al.*, 2007; Trzybiński *et al.*, 2010). With respective average deviations from planarity of 0.0427 (3) Å and 0.0066 (3) Å, the acridine and benzene ring systems are oriented at 76.8 (1)°. The acridine moieties in pairs are parallel (remain at an angle of 0.0 (1)°), while in adjacent pairs they are inclined at an angle of 62.4 (1)°. The mutual arrangement of the acridine and benzene ring systems, as well as the acridine skeletons in the crystal lattice is similar in the compound investigated and its precursor – 9-benzylacridine (Sikorski *et al.*, 2007).

Experimental

9-Benzylacridine was prepared by treating *N*-phenylaniline with an equimolar amount of phenylacetic acid, both dispersed in molten zinc chloride (Huntress & Shaw, 1948; Sikorski *et al.*, 2007). The crude product was purified chromatographically (SiO₂, cyclohexane-ethyl acetate, 5:2 v/v). The compound thus obtained was quaternarized with a five-fold molar excess of methyltrifluoromethanesulfonate dissolved in anhydrous dichloromethane (Trzybiński *et al.*, 2010). The crude 9-benzyl-10-methylacridinium trifluoromethanesulfonate was dissolved in a small amount of ethanol, filtered, and precipitated with a 25 v/v excess of diethyl ether. Light-orange crystals suitable for X-Ray investigations were grown from absolute ethanol solution (m.p. 478–480 K).

Refinement

H atoms were positioned geometrically, with C—H = 0.93 Å, 0.96 Å and 0.97 Å for the aromatic, methyl and methylene H atoms, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.2$ for the aromatic and $x = 1.5$ for the aliphatic H atoms.

Figures




Fig. 1. The molecular structure of the title compound showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 25% probability level and H atoms are shown as small spheres of arbitrary radius. Cg1, Cg2, Cg3 and Cg4 denote the ring centroids. The C—H \cdots O hydrogen bond is represented by a dashed line.




Fig. 2. The arrangement of the ions in the crystal structure. The C—H \cdots O interactions are represented by dashed lines, the C—H \cdots π , C—F \cdots π , S—O \cdots π and π — π contacts by dotted lines. H atoms not involved in interactions have been omitted. [Symmetry codes: (i) $x + 1/2, -y + 1/2, z + 1/2$; (ii) $-x + 2, -y, -z + 1$; (iii) $-x + 2, -y + 1, -z + 1$; (iv) $-x + 3/2, y - 1/2, -z + 1/2$; (v) $-x + 3/2, y + 1/2, -z + 1/2$.]

9-Benzyl-10-methylacridinium trifluoromethanesulfonate

Crystal data

$\text{C}_{21}\text{H}_{18}\text{N}^+ \cdot \text{CF}_3\text{O}_3\text{S}^-$

$M_r = 433.44$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 14.6211(7)$ Å

$b = 8.2514(2)$ Å

$c = 17.2900(8)$ Å

$\beta = 107.707(5)^\circ$

$V = 1987.12(15)$ Å³

$Z = 4$

$F(000) = 896$

$D_x = 1.449$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5587 reflections

$\theta = 3.2\text{--}29.2^\circ$

$\mu = 0.22$ mm⁻¹

$T = 295$ K

Plate, light-orange

$0.41 \times 0.25 \times 0.08$ mm

Data collection

Oxford Diffraction Gemini R Ultra Ruby CCD diffractometer	3528 independent reflections
Radiation source: Enhanced (Mo) X-ray Source graphite	2191 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.048$
Detector resolution: 10.4002 pixels mm^{-1}	$\theta_{\text{max}} = 25.1^\circ$, $\theta_{\text{min}} = 3.3^\circ$
ω scans	$h = -17 \rightarrow 12$
Absorption correction: multi-scan (<i>Crys.Alis RED</i> ; Oxford Diffraction, 2008)	$k = -9 \rightarrow 9$
$T_{\text{min}} = 0.953$, $T_{\text{max}} = 0.988$	$l = -20 \rightarrow 20$
16520 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.052$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.153$	H-atom parameters constrained
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0814P)^2 + 0.218P]$
3528 reflections	where $P = (F_o^2 + 2F_c^2)/3$
272 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.37 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.0142 (2)	0.2164 (4)	0.5705 (2)	0.0680 (10)
H1	1.0341	0.1838	0.6245	0.082*
C2	1.0692 (3)	0.1812 (4)	0.5215 (3)	0.0870 (13)
H2	1.1267	0.1251	0.5419	0.104*
C3	1.0384 (4)	0.2302 (5)	0.4410 (3)	0.0919 (14)
H3	1.0760	0.2037	0.4081	0.110*
C4	0.9568 (3)	0.3141 (4)	0.4080 (3)	0.0793 (11)
H4	0.9395	0.3451	0.3538	0.095*
C5	0.6717 (3)	0.5685 (3)	0.4389 (2)	0.0616 (9)
H5	0.6566	0.6103	0.3866	0.074*

supplementary materials

C6	0.6130 (3)	0.5955 (4)	0.4840 (3)	0.0703 (10)
H6	0.5572	0.6554	0.4620	0.084*
C7	0.6331 (2)	0.5369 (4)	0.5617 (2)	0.0641 (9)
H7	0.5903	0.5551	0.5911	0.077*
C8	0.7146 (2)	0.4531 (3)	0.59546 (18)	0.0515 (8)
H8	0.7275	0.4155	0.6484	0.062*
C9	0.8675 (2)	0.3374 (3)	0.58718 (16)	0.0416 (7)
N10	0.8163 (2)	0.4426 (3)	0.42625 (14)	0.0545 (7)
C11	0.9266 (2)	0.3027 (3)	0.53895 (18)	0.0490 (8)
C12	0.8982 (2)	0.3545 (3)	0.45600 (19)	0.0536 (8)
C13	0.7815 (2)	0.4203 (3)	0.55233 (15)	0.0411 (7)
C14	0.7575 (2)	0.4760 (3)	0.47112 (18)	0.0486 (8)
C15	0.8974 (2)	0.2879 (3)	0.67532 (17)	0.0514 (8)
H15A	0.8641	0.3559	0.7039	0.062*
H15B	0.9657	0.3076	0.6987	0.062*
C16	0.8769 (2)	0.1104 (3)	0.68912 (16)	0.0450 (7)
C17	0.7896 (2)	0.0396 (3)	0.64921 (19)	0.0565 (8)
H17	0.7431	0.0991	0.6111	0.068*
C18	0.7708 (2)	-0.1185 (4)	0.6653 (2)	0.0623 (9)
H18	0.7117	-0.1645	0.6383	0.075*
C19	0.8388 (3)	-0.2080 (4)	0.72059 (19)	0.0661 (10)
H19	0.8256	-0.3141	0.7320	0.079*
C20	0.9259 (3)	-0.1408 (4)	0.7590 (2)	0.0739 (11)
H20	0.9730	-0.2022	0.7954	0.089*
C21	0.9445 (3)	0.0179 (4)	0.74392 (18)	0.0627 (9)
H21	1.0038	0.0631	0.7713	0.075*
C22	0.7881 (3)	0.4991 (6)	0.3401 (2)	0.0921 (13)
H22A	0.7789	0.6144	0.3383	0.138*
H22B	0.7294	0.4470	0.3098	0.138*
H22C	0.8379	0.4721	0.3167	0.138*
S23	0.84575 (6)	0.55745 (8)	0.14455 (5)	0.0509 (3)
O24	0.8608 (2)	0.4008 (3)	0.17948 (15)	0.0869 (8)
O25	0.87049 (18)	0.6881 (3)	0.20097 (14)	0.0771 (7)
O26	0.75711 (18)	0.5802 (3)	0.08205 (16)	0.0925 (8)
C27	0.9326 (2)	0.5715 (3)	0.09040 (19)	0.0553 (8)
F28	1.01928 (18)	0.5412 (4)	0.13525 (16)	0.1322 (11)
F29	0.9156 (2)	0.4660 (3)	0.03015 (15)	0.1090 (9)
F30	0.9335 (2)	0.7119 (3)	0.05685 (18)	0.1184 (9)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.051 (2)	0.0523 (17)	0.102 (3)	-0.0032 (16)	0.025 (2)	-0.0051 (18)
C2	0.062 (3)	0.061 (2)	0.147 (4)	-0.0002 (18)	0.045 (3)	-0.014 (3)
C3	0.098 (4)	0.074 (3)	0.133 (4)	-0.020 (2)	0.078 (3)	-0.035 (3)
C4	0.094 (3)	0.070 (2)	0.092 (3)	-0.021 (2)	0.055 (3)	-0.021 (2)
C5	0.063 (2)	0.0494 (17)	0.058 (2)	-0.0072 (16)	-0.0032 (18)	0.0132 (15)
C6	0.058 (2)	0.0511 (18)	0.092 (3)	0.0070 (16)	0.007 (2)	-0.0005 (19)

C7	0.055 (2)	0.0556 (18)	0.082 (3)	0.0044 (16)	0.0208 (19)	-0.0152 (18)
C8	0.057 (2)	0.0479 (16)	0.0494 (17)	-0.0042 (15)	0.0155 (16)	-0.0066 (13)
C9	0.0406 (17)	0.0361 (14)	0.0437 (16)	-0.0092 (12)	0.0063 (14)	-0.0046 (12)
N10	0.0601 (18)	0.0573 (14)	0.0459 (14)	-0.0176 (13)	0.0158 (13)	0.0010 (12)
C11	0.0440 (18)	0.0397 (14)	0.065 (2)	-0.0079 (13)	0.0191 (16)	-0.0050 (14)
C12	0.061 (2)	0.0460 (16)	0.064 (2)	-0.0236 (15)	0.0350 (18)	-0.0155 (15)
C13	0.0463 (17)	0.0345 (13)	0.0397 (15)	-0.0065 (12)	0.0091 (13)	-0.0031 (12)
C14	0.054 (2)	0.0409 (15)	0.0485 (18)	-0.0132 (13)	0.0121 (16)	0.0005 (13)
C15	0.0536 (19)	0.0475 (16)	0.0480 (17)	-0.0013 (13)	0.0078 (15)	0.0024 (13)
C16	0.0483 (18)	0.0450 (15)	0.0387 (15)	-0.0019 (13)	0.0088 (14)	-0.0037 (12)
C17	0.0489 (19)	0.0520 (17)	0.0626 (19)	-0.0003 (14)	0.0081 (16)	-0.0030 (15)
C18	0.061 (2)	0.0570 (19)	0.069 (2)	-0.0097 (16)	0.0209 (18)	-0.0068 (17)
C19	0.101 (3)	0.0482 (17)	0.055 (2)	-0.0102 (19)	0.032 (2)	-0.0009 (16)
C20	0.095 (3)	0.058 (2)	0.053 (2)	0.004 (2)	-0.0001 (19)	0.0128 (16)
C21	0.068 (2)	0.0598 (19)	0.0486 (18)	-0.0066 (16)	0.0000 (17)	0.0059 (15)
C22	0.095 (3)	0.134 (3)	0.046 (2)	-0.025 (3)	0.020 (2)	0.017 (2)
S23	0.0509 (5)	0.0466 (4)	0.0560 (5)	-0.0018 (3)	0.0174 (4)	-0.0017 (3)
O24	0.131 (2)	0.0614 (14)	0.0900 (17)	0.0137 (14)	0.0658 (17)	0.0192 (13)
O25	0.0813 (18)	0.0706 (14)	0.0855 (16)	-0.0086 (12)	0.0343 (14)	-0.0294 (13)
O26	0.0492 (15)	0.114 (2)	0.1010 (19)	-0.0060 (14)	0.0025 (14)	-0.0007 (16)
C27	0.058 (2)	0.0492 (17)	0.060 (2)	0.0005 (15)	0.0213 (17)	0.0023 (15)
F28	0.0559 (15)	0.234 (3)	0.1091 (19)	0.0300 (17)	0.0292 (14)	0.015 (2)
F29	0.159 (3)	0.0953 (15)	0.1012 (17)	-0.0266 (15)	0.0826 (17)	-0.0321 (13)
F30	0.156 (2)	0.0668 (13)	0.172 (2)	0.0008 (14)	0.108 (2)	0.0329 (14)

Geometric parameters (Å, °)

C1—C2	1.364 (5)	C13—C14	1.416 (4)
C1—C11	1.421 (4)	C15—C16	1.528 (4)
C1—H1	0.9300	C15—H15A	0.9700
C2—C3	1.387 (6)	C15—H15B	0.9700
C2—H2	0.9300	C16—C21	1.374 (4)
C3—C4	1.346 (6)	C16—C17	1.382 (4)
C3—H3	0.9300	C17—C18	1.378 (4)
C4—C12	1.403 (5)	C17—H17	0.9300
C4—H4	0.9300	C18—C19	1.367 (4)
C5—C6	1.342 (5)	C18—H18	0.9300
C5—C14	1.428 (4)	C19—C20	1.363 (5)
C5—H5	0.9300	C19—H19	0.9300
C6—C7	1.372 (5)	C20—C21	1.378 (4)
C6—H6	0.9300	C20—H20	0.9300
C7—C8	1.348 (4)	C21—H21	0.9300
C7—H7	0.9300	C22—H22A	0.9600
C8—C13	1.425 (4)	C22—H22B	0.9600
C8—H8	0.9300	C22—H22C	0.9600
C9—C13	1.396 (4)	S23—O24	1.415 (2)
C9—C11	1.402 (4)	S23—O25	1.425 (2)
C9—C15	1.508 (4)	S23—O26	1.425 (2)
N10—C14	1.350 (4)	S23—C27	1.796 (3)

supplementary materials

N10—C12	1.361 (4)	C27—F28	1.293 (4)
N10—C22	1.495 (4)	C27—F30	1.297 (3)
C11—C12	1.432 (4)	C27—F29	1.322 (3)
C2—C1—C11	120.1 (4)	C9—C15—C16	114.0 (2)
C2—C1—H1	120.0	C9—C15—H15A	108.8
C11—C1—H1	120.0	C16—C15—H15A	108.8
C1—C2—C3	119.2 (4)	C9—C15—H15B	108.8
C1—C2—H2	120.4	C16—C15—H15B	108.8
C3—C2—H2	120.4	H15A—C15—H15B	107.7
C4—C3—C2	123.4 (4)	C21—C16—C17	118.1 (3)
C4—C3—H3	118.3	C21—C16—C15	120.4 (3)
C2—C3—H3	118.3	C17—C16—C15	121.5 (2)
C3—C4—C12	119.6 (4)	C18—C17—C16	120.6 (3)
C3—C4—H4	120.2	C18—C17—H17	119.7
C12—C4—H4	120.2	C16—C17—H17	119.7
C6—C5—C14	120.2 (3)	C19—C18—C17	120.4 (3)
C6—C5—H5	119.9	C19—C18—H18	119.8
C14—C5—H5	119.9	C17—C18—H18	119.8
C5—C6—C7	121.7 (3)	C20—C19—C18	119.6 (3)
C5—C6—H6	119.2	C20—C19—H19	120.2
C7—C6—H6	119.2	C18—C19—H19	120.2
C8—C7—C6	120.2 (4)	C19—C20—C21	120.3 (3)
C8—C7—H7	119.9	C19—C20—H20	119.9
C6—C7—H7	119.9	C21—C20—H20	119.9
C7—C8—C13	121.9 (3)	C16—C21—C20	121.1 (3)
C7—C8—H8	119.1	C16—C21—H21	119.5
C13—C8—H8	119.1	C20—C21—H21	119.5
C13—C9—C11	118.7 (3)	N10—C22—H22A	109.5
C13—C9—C15	121.0 (3)	N10—C22—H22B	109.5
C11—C9—C15	120.2 (3)	H22A—C22—H22B	109.5
C14—N10—C12	122.2 (3)	N10—C22—H22C	109.5
C14—N10—C22	118.6 (3)	H22A—C22—H22C	109.5
C12—N10—C22	119.2 (3)	H22B—C22—H22C	109.5
C9—C11—C1	121.4 (3)	O24—S23—O25	115.13 (15)
C9—C11—C12	119.4 (3)	O24—S23—O26	115.57 (17)
C1—C11—C12	119.2 (3)	O25—S23—O26	113.74 (15)
N10—C12—C4	122.0 (3)	O24—S23—C27	103.79 (14)
N10—C12—C11	119.5 (3)	O25—S23—C27	103.62 (15)
C4—C12—C11	118.5 (3)	O26—S23—C27	102.73 (16)
C9—C13—C14	120.4 (3)	F28—C27—F30	107.4 (3)
C9—C13—C8	122.6 (2)	F28—C27—F29	105.1 (3)
C14—C13—C8	117.0 (3)	F30—C27—F29	105.1 (3)
N10—C14—C13	119.6 (3)	F28—C27—S23	113.2 (2)
N10—C14—C5	121.5 (3)	F30—C27—S23	113.3 (2)
C13—C14—C5	118.9 (3)	F29—C27—S23	112.0 (2)
C11—C1—C2—C3	-0.2 (5)	C12—N10—C14—C5	-179.9 (2)
C1—C2—C3—C4	1.2 (6)	C22—N10—C14—C5	-2.4 (4)
C2—C3—C4—C12	-0.6 (6)	C9—C13—C14—N10	2.8 (4)

C14—C5—C6—C7	0.4 (5)	C8—C13—C14—N10	-177.3 (2)
C5—C6—C7—C8	1.6 (5)	C9—C13—C14—C5	-176.4 (2)
C6—C7—C8—C13	-0.9 (4)	C8—C13—C14—C5	3.5 (4)
C13—C9—C11—C1	-179.0 (2)	C6—C5—C14—N10	177.8 (3)
C15—C9—C11—C1	2.0 (4)	C6—C5—C14—C13	-3.0 (4)
C13—C9—C11—C12	0.7 (4)	C13—C9—C15—C16	99.8 (3)
C15—C9—C11—C12	-178.3 (2)	C11—C9—C15—C16	-81.1 (3)
C2—C1—C11—C9	178.5 (3)	C9—C15—C16—C21	136.1 (3)
C2—C1—C11—C12	-1.2 (4)	C9—C15—C16—C17	-45.8 (4)
C14—N10—C12—C4	177.2 (3)	C21—C16—C17—C18	1.1 (5)
C22—N10—C12—C4	-0.4 (4)	C15—C16—C17—C18	-177.0 (3)
C14—N10—C12—C11	-3.7 (4)	C16—C17—C18—C19	-0.5 (5)
C22—N10—C12—C11	178.8 (3)	C17—C18—C19—C20	-1.0 (5)
C3—C4—C12—N10	178.3 (3)	C18—C19—C20—C21	1.9 (5)
C3—C4—C12—C11	-0.9 (5)	C17—C16—C21—C20	-0.2 (5)
C9—C11—C12—N10	2.8 (4)	C15—C16—C21—C20	177.9 (3)
C1—C11—C12—N10	-177.5 (2)	C19—C20—C21—C16	-1.3 (5)
C9—C11—C12—C4	-178.0 (3)	O24—S23—C27—F28	-54.1 (3)
C1—C11—C12—C4	1.7 (4)	O25—S23—C27—F28	66.5 (3)
C11—C9—C13—C14	-3.5 (4)	O26—S23—C27—F28	-174.8 (3)
C15—C9—C13—C14	175.5 (2)	O24—S23—C27—F30	-176.8 (3)
C11—C9—C13—C8	176.6 (2)	O25—S23—C27—F30	-56.1 (3)
C15—C9—C13—C8	-4.4 (4)	O26—S23—C27—F30	62.5 (3)
C7—C8—C13—C9	178.3 (2)	O24—S23—C27—F29	64.6 (3)
C7—C8—C13—C14	-1.7 (4)	O25—S23—C27—F29	-174.8 (2)
C12—N10—C14—C13	0.9 (4)	O26—S23—C27—F29	-56.2 (2)
C22—N10—C14—C13	178.5 (3)		

Hydrogen-bond geometry (Å, °)

Cg4 is the centroid of the C16—C21 ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C2—H2...O26 ⁱ	0.93	2.49	3.398 (5)	167
C3—H3...Cg4 ⁱⁱ	0.93	2.74	3.630 (5)	161
C15—H15B...O25 ⁱⁱⁱ	0.97	2.49	3.423 (4)	160
C22—H22B...O25 ^{iv}	0.96	2.56	3.386 (5)	144
C22—H22C...O24	0.96	2.56	3.361 (5)	141

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (ii) $-x+2, -y, -z+1$; (iii) $-x+2, -y+1, -z+1$; (iv) $-x+3/2, y-1/2, -z+1/2$.

Table 2

C—F... π and *S*—O... π interactions (Å, °)

Cg1 and Cg3 are the centroids of the C9/N10/C11—C14 and C5—C8/C13/C14 rings, respectively.

<i>X</i>	<i>I</i>	<i>J</i>	<i>I</i> ... <i>J</i>	<i>X</i> ... <i>J</i>	<i>X</i> — <i>I</i> ... <i>J</i>
C27	F30	Cg3 ^v	3.115 (3)	4.233 (3)	143.0 (2)
S23	O26	Cg1 ^v	3.085 (3)	4.167 (2)	131.4 (2)

Symmetry code: (v) $-x + 3/2, y + 1/2, -z + 1/2$.

Table 3

π - π interactions (\AA , $^\circ$)

$Cg1$ and $Cg2$ are the centroids of the C9/N10/C11–C14 and C1–C4/C11/C12 rings, respectively. $CgI\cdots CgJ$ is the distance between ring centroids. The dihedral angle is that between the planes of the rings I and J . CgI_Perp is the perpendicular distance of CgI from ring J . CgI_Offset is the distance between CgI and perpendicular projection of CgJ on ring I .

I	J	$CgI\cdots CgJ$	Dihedral angle	CgI_Perp	CgI_Offset
1	2 ⁱⁱⁱ	3.806 (2)	2.11 (15)	3.575 (2)	1.306 (2)
2	1 ⁱⁱⁱ	3.806 (2)	2.11 (15)	3.530 (2)	1.423 (2)
2	2 ⁱⁱⁱ	3.886 (2)	0.02 (15)	3.563 (2)	1.551 (2)

Symmetry code: (iii) $-x + 2, -y + 1, -z + 1$.

Fig. 1




Fig. 2

