

2-(3-Bromo-4-methoxyphenyl)acetic acid

Ilia A. Guzei,* Alan R. Gunderson and Nicholas J. Hill

Department of Chemistry, University of Wisconsin—Madison, 1101 University Ave, Madison, WI 53706, USA

Correspondence e-mail: iguzei@chem.wisc.edu


Received 14 May 2010; accepted 27 May 2010

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.026; wR factor = 0.073; data-to-parameter ratio = 14.4.

The title compound $\text{C}_9\text{H}_9\text{BrO}_3$, was synthesized by the regioselective bromination of 4-methoxyphenylacetic acid using bromine in acetic acid in a 84% yield. In the molecular structure, the methoxy group is almost coplanar with the phenyl ring within 0.06 Å; the acetic acid substituent is tilted by 78.15 (7)° relative to the ring. The C—C—C angles at the OMe, acetyl and Br substituents are 118.2 (2), 118.4 (2) and 121.5 (2)°, respectively, indicating that the Br atom is electron-withdrawing, whereas the other substituents possess electron-donating properties. In the crystal, the molecules form centrosymmetric strongly O—H...O hydrogen-bonded dimers of the type $R_2^2(8)$.

Related literature

For the use of the title compound in the synthesis of natural products such as *Combretastatin A-4*, see: Zou *et al.* (2008); for *Verongamine*, see: Wasserman & Wang (1998) and for model *Vancomycin*-type systems, see: Ghosh *et al.* (2009). The iodanalogue featured in the synthesis of (+)-*Phleichrome* and (+)-*Calphostin D*, see: Morgan *et al.* (2010). For the synthesis of the title compound, see: Coutts *et al.*, (1970); Morgan *et al.*, (2007); Zou *et al.* (2008); Ghosh *et al.* (2009). For background for our program to introduce natural product synthesis, crystal growing techniques and single crystal X-ray diffraction data analysis into the undergraduate curriculum, see: Findlater *et al.*, (2010); Guzei *et al.*, (2010a). For a discussion of hydrogen-bonding motif assignment, see: Guzei *et al.* (2010b). Outlier reflections were omitted based on the statistics test described by Prince & Nicholson (1983) and Rollett (1988), and implemented in *FCF_filter* (Guzei, 2007).



Experimental

Crystal data

$\text{C}_9\text{H}_9\text{BrO}_3$	$V = 930.67$ (5) Å ³
$M_r = 245.06$	$Z = 4$
Monoclinic, $P2_1/c$	Cu $K\alpha$ radiation
$a = 12.5022$ (4) Å	$\mu = 5.81$ mm ⁻¹
$b = 8.2690$ (2) Å	$T = 120$ K
$c = 9.0199$ (3) Å	$0.46 \times 0.37 \times 0.19$ mm
$\beta = 93.573$ (1)°	

Data collection

Bruker SMART APEXII area-detector diffractometer	12598 measured reflections
Absorption correction: analytical (<i>SADABS</i> ; Bruker, 2007)	1725 independent reflections
$T_{\min} = 0.177$, $T_{\max} = 0.398$	1708 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$	120 parameters
$wR(F^2) = 0.073$	H-atom parameters constrained
$S = 1.08$	$\Delta\rho_{\text{max}} = 0.86$ e Å ⁻³
1725 reflections	$\Delta\rho_{\text{min}} = -0.47$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O3}-\text{H3}\cdots\text{O2}^i$	0.84	1.82	2.661 (2)	179

Symmetry code: (i) $-x + 1, -y - 1, -z$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT-Plus* (Bruker, 2007); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *publCIF* (Westrip, 2010) and *modiCIFer* (Guzei, 2007).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2206).

References

- Bruker (2007). *APEX2*, *SADABS* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Coutts, I. G. C., Durbin, A. K. & Schofield, K. (1970). *Aust. J. Chem.* **23**, 791–800.
- Findlater, M., Hill, N. J. & Cowley, A. H. (2010). *J. Chem. Crystallogr.* **40**, 64–66.

- Ghosh, S., Kumar, A. S., Mehta, G. N. & Soundararajan, R. (2009). *Synthesis*, pp. 3322–3326.
- Guzei, I. A. (2007). In-house Crystallographic Programs: *FCF_filter*, *INSerter* and *modiCIFer*. Molecular Structure Laboratory, University of Wisconsin–Madison, Madison, Wisconsin, USA.
- Guzei, I. A., Hill, N. J. & Van Hout, M. R. (2010a). *Acta Cryst.* **E66**, o40–o41.
- Guzei, I. A., Spencer, L. C., Ainooson, M. K. & Darkwa, J. (2010b). *Acta Cryst.* **C66**, m89–m96.
- Morgan, B. J., Mulrooney, C. A., O'Brien, E. M. & Kozlowski, M. C. (2010). *J. Org. Chem.* **75**, 30–43.
- Morgan, B. J., Xie, X., Phuan, P.-W. & Kozlowski, M. C. (2007). *J. Org. Chem.* **72**, 6171–6182.
- Prince, E. & Nicholson, W. L. (1983). *Acta Cryst.* **A39**, 407–410.
- Rollett, J. S. (1988). *Crystallographic Computing*, edited by N. W. Isaacs & M. R. Taylor, Vol. 4, pp. 149–166. Oxford University Press.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wasserman, H. H. & Wang, J. (1998). *J. Org. Chem.* **63**, 5581–5586.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.
- Zou, Y., Xiao, C.-F., Zhong, R.-Q., Wei, W., Hunag, W.-M. & He, S.-J. (2008). *J. Chem. Res.* pp. 354–356.

supplementary materials

Acta Cryst. (2010). E66, o1555-o1556 [doi:10.1107/S1600536810020143]

2-(3-Bromo-4-methoxyphenyl)acetic acid

I. A. Guzei, A. R. Gunderson and N. J. Hill

Comment

Recently, we have been pursuing simple organic and organometallic compounds as candidates for the introduction of (a) natural product synthesis into the undergraduate teaching laboratory and (b) crystal growing techniques and single crystal X-ray diffraction data analysis into the undergraduate curriculum (Findlater *et al.*, 2010; Guzei *et al.*, 2010a). The 3-bromo-4-methoxyphenylacetic acid **I** has been employed in the synthesis of natural products such as *Combretastatin A-4*, (Zou *et al.*, 2008), *Verongamine* (Wasserman & Wang, 1998) and model *Vancomycin*-type systems (Ghosh *et al.*, 2009). The iodo-analogue features in the synthesis of the perylenequinones (+)-*Phleichrome* and (+)-*Calphostin D*, (Morgan *et al.*, 2010). Our interest in **I** stems from its role in the synthesis of the antimetabolic compound *Combretastatin A-4* via a simple Perkin condensation/decarboxylation sequence (Zou *et al.*, 2008). This concise route, employing commercially available starting materials, followed by the facile purification of **I** to furnish high quality crystals makes it ideal in both regards. Compound **I** is readily synthesized by the regioselective bromination of 4-methoxyphenylacetic acid using bromine in acetic acid (Coutts *et al.*, 1970; Morgan *et al.*, 2007; Zou *et al.*, 2008; Ghosh *et al.*, 2009). Compound **I** was isolated and characterized by NMR, mp, and single-crystal X-ray analysis. There are three main structural aspects students should identify. First, the positions of the alkyl substituents on the phenyl ring. The methoxy-group is almost coplanar with the ring, torsion angle C7—O1—C1—C6 is 1.2 (3)°, whereas the acetic acid terminus is nearly perpendicular to the ring with the dihedral angle between the planes defined by atoms C1—C6 and atoms C4,C8,C9,O2,O3 spanning 78.15 (7)°. Secondly, the distortions of the C—C—C angles from 120° at the substituents of the phenyl ring reflect their electronic properties. The stronger the electron-withdrawing power of a substituent, the larger the C—C—C angle. The angles at OMe, Ac and Br are 118.2 (2), 118.4 (2), and 121.5 (2)°, respectively, indicating that the Br atom is electron-withdrawing, whereas the other substituents possess electron-donating properties. Of course, the magnitude of the values is affected by the neighbouring substituents. Thirdly, the molecules of **I** form centrosymmetric strongly hydrogen-bonded dimers in the lattice. The hydrogen bonding motif is $R_2^2(8)$. A topical discussion of hydrogen bonding motif assignment was published by Guzei *et al.*, 2010b.

Experimental

To a stirred solution of 4-methoxyphenylacetic acid (10 g, 60.2 mmol) in acetic acid (60 ml) was added a solution of bromine (9.62 g, 3.1 ml, 60.2 mmol) in acetic acid (30 ml) slowly dropwise over 30 min. The mixture was stirred at room temperature (60 min) and then poured into 500 ml ice-water. The resultant pale yellow, turbid mixture was stirred (10 min), filtered, rinsed with ice-water (3×10 ml), air-dried (20 min) and recrystallized from hot xylene to give a white crystalline powder. Yield 12.41 g, 84 %. M.p. 386.3–387.2 K.

^1H NMR (CDCl_3): δ 3.56 (2H, s, CH_2); 3.89 (3H, s, OCH_3), 6.86 (1H, d), 7.19 (1H, dd), 7.48 (1H, d). ^{13}C (^1H) NMR (CDCl_3): δ 39.9; 56.5; 111.9; 112.2; 127.0; 129.6; 134.6; 155.5; 178.0.

Refinement

All H-atoms were placed in idealized locations. The C—H distances were 0.98 Å for the methyl group, 0.99 Å the methylene group, 0.95 Å for the sp^2 -hybridized atoms; the O—H distance was fixed at 0.84 Å. All H atoms were refined as riding with thermal displacement coefficients $U_{iso}(H)$ set to $1.5U_{eq}(C, O)$ for the methyl- and hydroxyl-groups and to $1.2U_{eq}(C)$ for the CH- and CH_2 -groups.

The outlier reflections were omitted based on the statistics test described by Prince & Nicholson, (1983) and Rollett, (1988), and implemented in program FCF_filter (Guzei, 2007). The number of omitted outliers is 4.

Figures




Fig. 1. Molecular structure of **I** with the atom numbering scheme. The displacement ellipsoids are shown at 50% probability level.

2-(3-Bromo-4-methoxyphenyl)acetic acid

Crystal data

$C_9H_9BrO_3$

$M_r = 245.06$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 12.5022(4)$ Å

$b = 8.2690(2)$ Å

$c = 9.0199(3)$ Å

$\beta = 93.573(1)^\circ$

$V = 930.67(5)$ Å³

$Z = 4$

$F(000) = 488$

$D_x = 1.749$ Mg m⁻³

Melting point = 386.3–387.2 K

Cu $K\alpha$ radiation, $\lambda = 1.54178$ Å

Cell parameters from 9563 reflections

$\theta = 3.5$ – 69.5°

$\mu = 5.81$ mm⁻¹

$T = 120$ K

Block, colourless

$0.46 \times 0.37 \times 0.19$ mm

Data collection

Bruker SMART APEXII area-detector
diffractometer

1725 independent reflections

Radiation source: fine-focus sealed tube graphite	1708 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.030$
$0.5^\circ \omega$ and $0.5^\circ \varphi$ scans	$\theta_{\text{max}} = 69.5^\circ$, $\theta_{\text{min}} = 3.5^\circ$
Absorption correction: analytical (SADABS; Bruker, 2007)	$h = -14 \rightarrow 15$
$T_{\text{min}} = 0.177$, $T_{\text{max}} = 0.398$	$k = -9 \rightarrow 10$
12598 measured reflections	$l = -10 \rightarrow 10$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.026$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.073$	H-atom parameters constrained
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.0442P)^2 + 1.1574P]$
1725 reflections	where $P = (F_o^2 + 2F_c^2)/3$
120 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.86 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.47 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.253622 (18)	0.20274 (3)	0.38506 (3)	0.02281 (12)
O1	0.05071 (12)	0.0202 (2)	0.32075 (17)	0.0186 (3)
O2	0.40281 (14)	-0.3902 (2)	0.05910 (19)	0.0234 (4)
O3	0.50409 (14)	-0.3283 (2)	-0.12886 (19)	0.0229 (4)
H3	0.5326	-0.4178	-0.1071	0.034*
C1	0.12214 (17)	-0.0244 (3)	0.2209 (2)	0.0151 (4)
C2	0.22340 (18)	0.0476 (3)	0.2339 (2)	0.0148 (4)
C3	0.30189 (17)	0.0087 (3)	0.1381 (2)	0.0160 (4)
H3A	0.3701	0.0593	0.1493	0.019*
C4	0.28118 (18)	-0.1041 (3)	0.0255 (2)	0.0172 (5)
C5	0.1810 (2)	-0.1765 (3)	0.0129 (3)	0.0201 (5)

supplementary materials

H5	0.1661	-0.2542	-0.0631	0.024*
C6	0.10183 (18)	-0.1383 (3)	0.1087 (3)	0.0181 (5)
H6	0.0339	-0.1898	0.0977	0.022*
C7	-0.05187 (18)	-0.0591 (3)	0.3100 (3)	0.0220 (5)
H7A	-0.0887	-0.0346	0.2135	0.033*
H7C	-0.0952	-0.0204	0.3896	0.033*
H7B	-0.0416	-0.1762	0.3195	0.033*
C8	0.3654 (2)	-0.1464 (3)	-0.0810 (3)	0.0209 (5)
H8A	0.4175	-0.0564	-0.0831	0.025*
H8B	0.3306	-0.1576	-0.1821	0.025*
C9	0.42502 (19)	-0.3007 (3)	-0.0408 (3)	0.0170 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.02043 (17)	0.02308 (18)	0.02445 (17)	-0.00116 (9)	-0.00227 (11)	-0.00996 (9)
O1	0.0164 (7)	0.0206 (8)	0.0193 (8)	-0.0018 (6)	0.0047 (6)	-0.0044 (6)
O2	0.0309 (9)	0.0194 (8)	0.0214 (9)	0.0053 (7)	0.0119 (7)	0.0032 (7)
O3	0.0222 (9)	0.0241 (9)	0.0236 (9)	0.0068 (7)	0.0096 (7)	0.0059 (7)
C1	0.0169 (10)	0.0131 (10)	0.0152 (10)	0.0020 (8)	0.0009 (8)	0.0024 (8)
C2	0.0193 (10)	0.0110 (10)	0.0137 (10)	0.0010 (8)	-0.0029 (8)	-0.0009 (8)
C3	0.0163 (10)	0.0137 (10)	0.0180 (10)	0.0004 (8)	0.0005 (8)	0.0035 (8)
C4	0.0210 (11)	0.0152 (11)	0.0157 (11)	0.0040 (9)	0.0041 (8)	0.0040 (8)
C5	0.0252 (12)	0.0181 (11)	0.0169 (11)	0.0004 (9)	0.0013 (9)	-0.0035 (9)
C6	0.0189 (11)	0.0165 (11)	0.0188 (11)	-0.0030 (9)	0.0005 (9)	-0.0020 (9)
C7	0.0160 (11)	0.0241 (12)	0.0262 (12)	-0.0026 (9)	0.0028 (9)	-0.0010 (10)
C8	0.0246 (12)	0.0211 (12)	0.0176 (11)	0.0032 (10)	0.0072 (9)	0.0016 (9)
C9	0.0178 (11)	0.0188 (12)	0.0145 (11)	-0.0014 (8)	0.0019 (9)	-0.0039 (8)

Geometric parameters (\AA , $^\circ$)

Br1—C2	1.893 (2)	C4—C5	1.386 (3)
O1—C1	1.359 (3)	C4—C8	1.510 (3)
O1—C7	1.438 (3)	C5—C6	1.390 (3)
O2—C9	1.212 (3)	C5—H5	0.9500
O3—C9	1.326 (3)	C6—H6	0.9500
O3—H3	0.8400	C7—H7A	0.9800
C1—C6	1.394 (3)	C7—H7C	0.9800
C1—C2	1.397 (3)	C7—H7B	0.9800
C2—C3	1.385 (3)	C8—C9	1.510 (3)
C3—C4	1.392 (3)	C8—H8A	0.9900
C3—H3A	0.9500	C8—H8B	0.9900
C1—O1—C7	116.85 (18)	C5—C6—H6	120.0
C9—O3—H3	109.5	C1—C6—H6	120.0
O1—C1—C6	124.6 (2)	O1—C7—H7A	109.5
O1—C1—C2	117.3 (2)	O1—C7—H7C	109.5
C6—C1—C2	118.2 (2)	H7A—C7—H7C	109.5
C3—C2—C1	121.5 (2)	O1—C7—H7B	109.5

C3—C2—Br1	119.28 (17)	H7A—C7—H7B	109.5
C1—C2—Br1	119.23 (17)	H7C—C7—H7B	109.5
C2—C3—C4	120.3 (2)	C4—C8—C9	113.34 (19)
C2—C3—H3A	119.9	C4—C8—H8A	108.9
C4—C3—H3A	119.9	C9—C8—H8A	108.9
C5—C4—C3	118.4 (2)	C4—C8—H8B	108.9
C5—C4—C8	120.7 (2)	C9—C8—H8B	108.9
C3—C4—C8	120.9 (2)	H8A—C8—H8B	107.7
C4—C5—C6	121.7 (2)	O2—C9—O3	123.7 (2)
C4—C5—H5	119.2	O2—C9—C8	124.2 (2)
C6—C5—H5	119.2	O3—C9—C8	112.16 (19)
C5—C6—C1	120.0 (2)		
C7—O1—C1—C6	1.2 (3)	C3—C4—C5—C6	0.4 (3)
C7—O1—C1—C2	-177.70 (19)	C8—C4—C5—C6	-179.3 (2)
O1—C1—C2—C3	179.39 (19)	C4—C5—C6—C1	0.1 (4)
C6—C1—C2—C3	0.4 (3)	O1—C1—C6—C5	-179.4 (2)
O1—C1—C2—Br1	-1.0 (3)	C2—C1—C6—C5	-0.5 (3)
C6—C1—C2—Br1	180.00 (17)	C5—C4—C8—C9	-81.5 (3)
C1—C2—C3—C4	0.1 (3)	C3—C4—C8—C9	98.8 (3)
Br1—C2—C3—C4	-179.53 (17)	C4—C8—C9—O2	5.8 (3)
C2—C3—C4—C5	-0.5 (3)	C4—C8—C9—O3	-175.0 (2)
C2—C3—C4—C8	179.2 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O3—H3 \cdots O2 ⁱ	0.84	1.82	2.661 (2)	179.

Symmetry codes: (i) $-x+1, -y-1, -z$.

Fig. 1

