

4-[[*E*-(3,5-Dimethyl-1-phenyl-1*H*-pyrazol-4-yl)methylidene]amino]-1,5-dimethyl-2-phenyl-1*H*-pyrazol-3(2*H*)-one

Hoong-Kun Fun,^{a*‡} Madhukar Hemamalini,^a Abdullah M. Asiri^{b§} and Salman A. Khan^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Chemistry, Faculty of Science, King Abdu Aziz University, Jeddah, Saudi Arabia
Correspondence e-mail: hkfun@usm.my


Received 31 May 2010; accepted 3 June 2010

Key indicators: single-crystal X-ray study; *T* = 296 K; mean $\sigma(\text{C}-\text{C}) = 0.003 \text{ \AA}$; *R* factor = 0.059; *wR* factor = 0.161; data-to-parameter ratio = 19.3.

The title Schiff base compound, $\text{C}_{23}\text{H}_{23}\text{N}_5\text{O}$, was synthesized by the reaction of 4-aminophenazone and 3,5-dimethyl-1-phenylpyrazole-4-carboxaldehyde. The molecule adopts an *E* configuration about the central $\text{C}=\text{N}$ double bond. A weak intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond generates an *S*(6) ring motif. The dihedral angle between the pyrazole rings is $24.72(10)^\circ$ and the dihedral angles between the pyrazole rings and the adjacent phenyl rings are $58.67(10)$ and $46.58(11)^\circ$. The crystal structure is stabilized by weak $\text{C}-\text{H}\cdots\pi$ interactions involving the pyrazolone and phenyl rings.

Related literature

For background to and applications of heterocyclic Schiff bases, see: Nawaz *et al.* (2009); Li *et al.* (1999); Urena *et al.* (2003); Geronikaki *et al.* (2003); Shanker *et al.* (2009); Pandeya *et al.* (1999); Sridhar *et al.* (2002); Nawrocka *et al.* (2004). For related structures, see: Eryigit & Kendi (1998); Manikandan *et al.* (2000). For details of hydrogen-bond motifs, see: Bernstein *et al.* (1995).



[‡] Thomson Reuters ResearcherID: A-3561-2009.

[§] On secondment to: The Center of Excellence for Advanced Materials Research, King Abdu Aziz University, Jeddah 21589, Saudi Arabia.

Experimental

Crystal data

$\text{C}_{23}\text{H}_{23}\text{N}_5\text{O}$
M_r = 385.46
Monoclinic, $P2_1/c$
a = 15.2985 (2) Å
b = 7.6827 (1) Å
c = 19.6737 (3) Å
 β = 116.905 (1) $^\circ$

V = 2062.03 (5) Å^3
Z = 4
Mo $K\alpha$ radiation
 μ = 0.08 mm^{-1}
T = 296 K
0.45 × 0.21 × 0.10 mm

Data collection

Bruker SMART APEXII CCD
area-detector diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)
T_{min} = 0.965, *T_{max}* = 0.992

23014 measured reflections
5993 independent reflections
2881 reflections with *I* > 2σ(*I*)
R_{int} = 0.048

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.161$
S = 1.03
5993 reflections
311 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.18 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å , $^\circ$).

*Cg*1 and *Cg*2 are the centroids of the N4/N5/C11–C13 and C1–C6 rings, respectively.

<i>D</i> — <i>H</i> ⋯ <i>A</i>	<i>D</i> — <i>H</i>	<i>H</i> ⋯ <i>A</i>	<i>D</i> ⋯ <i>A</i>	<i>D</i> — <i>H</i> ⋯ <i>A</i>
C10—H10A⋯O1	0.986 (18)	2.40 (2)	3.052 (3)	123.3 (14)
C19—H19A⋯ <i>Cg</i> 2 ⁱ	0.990 (19)	2.656 (19)	3.452 (2)	137.4 (17)
C20—H20C⋯ <i>Cg</i> 1 ⁱⁱ	0.96	2.85 (3)	3.720 (3)	149 (1)
C22—H22B⋯ <i>Cg</i> 2 ⁱⁱⁱ	0.96	2.82 (3)	3.585 (3)	135 (1)

Symmetry codes: (i) $-x + 1, -y - 1, -z - 1$; (ii) $-x, -y - 2, -z - 1$; (iii) $-x + 1, y - \frac{1}{2}, -z - \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and MH thank the Malaysian Government and Universiti Sains Malaysia (USM) for the Research University Golden Goose grant No. 1001/PFIZIK/811012. MH also thanks USM for a post-doctoral research fellowship. AMA and SAK thank the Chemistry Department, King Abdul Aziz University, Jeddah, for providing research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5060).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Eryigit, R. & Kendi, E. (1998). *J. Chem. Crystallogr.* **28**, 145–147.
Geronikaki, J. M. A., Litina, D. H. & Amourgianou, M. (2003). *Farmaco*, **58**, 489–495.

- Li, X., Schofield, B. H., Huang, C., Kleiner, G. I., Hugh, A. & Sampson, H. A. (1999). *J. Allergy Clin. Immunol.* **103**, 206–214.
- Manikandan, P., Justin Thomas, K. R. & Manoharan, P. T. (2000). *Acta Cryst. C* **56**, 308–309.
- Nawaz, H., Akhter, Z., Yameen, S., Siddiqi, H. M., Mirza, B. & Rifat, A. (2009). *J. Organomet. Chem.* **694**, 2198–2203.
- Nawrocka, W., Sztuba, B., Kowalska, M. W., Liszkiewicz, H., Wietrzyk, J., Nasulewicz, A., Peczyńska, M. & Opolski, A. (2004). *Il Farmaco*, **59**, 83–91.
- Pandeya, S. N., Sriram, D., Nath, G. & Clercq, E. D. (1999). *Pharm. Acta Helv.* **74**, 11–17.
- Shanker, K., Rohini, R., Ravinder, V., Reddy, P. M. & Ho, Y. (2009). *Spectrochim. Acta Part A*, **73**, 205–211.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Sridhar, S. K., Pandeya, S. N., Stables, J. P. & Ramesh, A. (2002). *Eur. J. Pharm. Sci.* **16**, 129–132.
- Urena, F. H., Illn-Cabeza, N. A., Moreno-Carretero, M. N. & José, M. (2003). *J. Inorg. Biochem.* **94**, 326–334.

supplementary materials

Acta Cryst. (2010). E66, o1602-o1603 [doi:10.1107/S1600536810021173]

4-[(*E*)-(3,5-Dimethyl-1-phenyl-1*H*-pyrazol-4-yl)methylidene]amino}-1,5-dimethyl-2-phenyl-1*H*-pyrazol-3(2*H*)-one

H.-K. Fun, M. Hemamalini, A. M. Asiri and S. A. Khan

Comment

Heterocyclic Schiff bases have attracted continuing interest over the years because of their varied biological activities (Nawaz *et al.*, 2009). Recently they have found application in drug development for the treatment of allergies (Li *et al.*, 1999), hypertension (Urena *et al.*, 2003), inflammation (Geronikaki *et al.*, 20003), bacterial (Shanker *et al.*, 2009), HIV infections (Pandeya *et al.*, 1999) and hypnotics (Sridhar *et al.*, 2002). More recently they have also been used for the treatment of pain acting as fibrinogen receptor antagonists with antithrombotic activity (Nawrocka *et al.*, 2004). Due to wide application of pyrazoline-containing Schiff bases, we have synthesized a novel pyrazoline-containing Schiff base from 4-aminophenazone.

In the title compound (Fig. 1), the rings A (C14–C19), B (N4/N5/C11–C13), C (N1/N2/C7–C9) and D (C1–C6) are essentially planar. The dihedral angle between the best planes of the rings are A/B = 58.67 (10)°, A/C = 83.07 (10)°, A/D = 79.53 (12)°, B/C = 24.72 (10)°, B/D = 44.68 (11)° and C/D = 46.58 (11)°. The molecule adopts a *trans* configuration about the central C10=N3 double bond. The C–N bond lengths of N1–C6 = 1.423 (2) Å; N2–C8 = 1.375 (2) Å; N3–C9 = 1.400 (2) Å; N5–C13 = 1.357 (2) Å; N1–C7 = 1.404 (2) Å; N2–C20 = 1.467 (2) Å; N4–C12 = 1.325 (2) Å and N5–C14 = 1.429 (2) are normal for C–N single-bond distances. The distance between C10–N3 (1.287 (2) Å) is typical for a C=N double-bond distance. These bonds are comparable with those in *N*-(1*H*-benzimidazol-2-ylmethyl)-*N*-(2,6-dichlorophenyl) amine (Eryigit & Kendi, 1998). The N1–N2 and N4–N5 (1.4082 (19) Å & 1.3702 (19) Å) single-bond lengths are comparable with those in 2,6-bis(3,5-dimethylpyrazol-1-ylmethyl) pyridine (Manikandan *et al.*, 2000). An weak intramolecular C10—H10A⋯O1 hydrogen bond interaction generates an *S*(6) ring motif (Bernstein *et al.*, 1995).

In the crystal structure (Fig. 2) there are no classical hydrogen bonds but stabilization is provided by weak C20—H20C⋯Cg1ⁱ, C19—H19A⋯Cg2ⁱⁱ and C22—H22B⋯Cg2ⁱⁱⁱ interactions (see Table 1 for symmetry codes). Cg1 and Cg2 are the centroids of rings (N4/N5/C11–C13) and (C1–C6) rings respectively.

Experimental

A mixture of 4-aminophenazone (0.50g, 0.0025 mol) and 3,5-dimethyl-1-phenylpyrazole-4-carboxaldehyde (0.50 g, 0.0025 mol) in methanol (15 mL) was heated for 3 h to give a yellow precipitate. It was then filtered and washed with methanol to give the pure schiff bases (I). Colourless crystals of (I) are recrystallized from methanol. Yield: 68%; m. p. 206°C. IR (KBr) ν_{\max} cm⁻¹: 2980 (C–H aromatic), 1642 (HC=N), 1607 (C=O), 1134 (C–N). ¹H-NMR (600 MHz, CDCl₃) δ : 9.75 (CH=N, s), 7.49–7.29 (CH aromatic, m), 3.1(N–CH₃, s), 2.84 (N–CH₃, s), 2.15 (CH₃, s), 1.70 (CH₃, s).

Refinement

Atoms H1A, H2A, H3A, H4A, H5A, H10A, H15A, H16A, H17A, H18A and H19A were located from a difference Fourier maps and refined freely. The methyl H atoms were positioned geometrically [C–H = 0.96 Å] and were refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$. A rotating group model was used for the methyl group.

Figures




Fig. 1. The asymmetric unit of the title compound, showing 50% probability displacement ellipsoids and the atom-numbering scheme. An intramolecular hydrogen bond is shown as dashed line.




Fig. 2. The crystal packing of the title compound showing the molecules stacked along *b*-axis.

4-[[*(E)*-(3,5-Dimethyl-1-phenyl-1*H*-pyrazol-4-yl)methylidene]amino]-1,5-dimethyl-2-phenyl-1*H*-pyrazol-3(2*H*)-one

Crystal data

$\text{C}_{23}\text{H}_{23}\text{N}_5\text{O}$

$M_r = 385.46$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 15.2985(2) \text{ \AA}$

$b = 7.6827(1) \text{ \AA}$

$c = 19.6737(3) \text{ \AA}$

$\beta = 116.905(1)^\circ$

$V = 2062.03(5) \text{ \AA}^3$

$Z = 4$

$F(000) = 816$

$D_x = 1.242 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3425 reflections

$\theta = 2.8\text{--}22.2^\circ$

$\mu = 0.08 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Block, colourless

$0.45 \times 0.21 \times 0.10 \text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

5993 independent reflections

2881 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.048$

$\theta_{\text{max}} = 30.0^\circ$, $\theta_{\text{min}} = 1.5^\circ$

Absorption correction: multi-scan
(*SADABS*; Bruker, 2009) $h = -20 \rightarrow 21$
 $T_{\min} = 0.965$, $T_{\max} = 0.992$ $k = -10 \rightarrow 10$
 23014 measured reflections $l = -27 \rightarrow 27$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.059$ H atoms treated by a mixture of independent and constrained refinement
 $wR(F^2) = 0.161$ $w = 1/[\sigma^2(F_o^2) + (0.0683P)^2 + 0.0093P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $S = 1.03$ $(\Delta/\sigma)_{\max} < 0.001$
 5993 reflections $\Delta\rho_{\max} = 0.24 \text{ e } \text{\AA}^{-3}$
 311 parameters $\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$
 0 restraints Extinction correction: *SHELXTL* (Sheldrick, 2008),
 $F_c^* = kFc[1 + 0.001x Fc^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Primary atom site location: structure-invariant direct methods Extinction coefficient: 0.0055 (13)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.11416 (10)	0.47531 (17)	0.60512 (9)	0.0735 (5)
N1	0.04112 (10)	0.21383 (18)	0.60925 (9)	0.0508 (4)
N2	0.07320 (10)	0.04036 (17)	0.62683 (9)	0.0474 (4)
N3	0.28754 (10)	0.22512 (19)	0.62644 (8)	0.0470 (4)
N4	0.57171 (10)	0.41293 (19)	0.66310 (9)	0.0509 (4)
N5	0.53154 (10)	0.55923 (17)	0.62050 (8)	0.0426 (4)
C1	-0.09125 (16)	0.4099 (3)	0.58937 (15)	0.0687 (6)
C2	-0.16074 (18)	0.4715 (4)	0.6111 (2)	0.0865 (9)
C3	-0.17010 (19)	0.3990 (4)	0.6710 (2)	0.0855 (8)
C4	-0.10854 (18)	0.2665 (3)	0.71182 (17)	0.0727 (7)
C5	-0.03755 (15)	0.2068 (3)	0.69323 (13)	0.0568 (5)

supplementary materials

C6	-0.03009 (13)	0.2767 (2)	0.63100 (12)	0.0513 (5)
C7	0.11729 (13)	0.3159 (2)	0.60932 (10)	0.0491 (5)
C8	0.16367 (13)	0.0318 (2)	0.62798 (10)	0.0449 (4)
C9	0.19323 (12)	0.1939 (2)	0.61851 (9)	0.0428 (4)
C10	0.30947 (14)	0.3749 (2)	0.60912 (10)	0.0463 (4)
C11	0.40715 (12)	0.4167 (2)	0.62077 (9)	0.0410 (4)
C12	0.49667 (13)	0.3256 (2)	0.66248 (10)	0.0478 (5)
C13	0.43302 (12)	0.5662 (2)	0.59479 (9)	0.0409 (4)
C14	0.59465 (12)	0.6879 (2)	0.61384 (10)	0.0418 (4)
C15	0.66808 (15)	0.7576 (3)	0.67935 (12)	0.0586 (5)
C16	0.72986 (18)	0.8801 (3)	0.67306 (15)	0.0739 (7)
C17	0.71962 (16)	0.9330 (3)	0.60346 (14)	0.0670 (6)
C18	0.64705 (16)	0.8606 (3)	0.53863 (14)	0.0605 (6)
C19	0.58412 (15)	0.7374 (2)	0.54338 (11)	0.0508 (5)
C20	-0.00160 (14)	-0.0950 (2)	0.59196 (12)	0.0614 (5)
H20A	-0.0387	-0.0723	0.5384	0.092*
H20B	-0.0446	-0.0948	0.6154	0.092*
H20C	0.0296	-0.2066	0.5992	0.092*
C21	0.21737 (16)	-0.1350 (2)	0.64000 (15)	0.0748 (7)
H21A	0.2829	-0.1125	0.6477	0.112*
H21B	0.1844	-0.2079	0.5960	0.112*
H21C	0.2196	-0.1927	0.6840	0.112*
C22	0.51419 (15)	0.1564 (3)	0.70426 (13)	0.0727 (6)
H22A	0.5833	0.1332	0.7302	0.109*
H22B	0.4817	0.0645	0.6687	0.109*
H22C	0.4889	0.1629	0.7408	0.109*
C23	0.37292 (14)	0.7138 (3)	0.54809 (13)	0.0644 (6)
H23A	0.4032	0.8218	0.5715	0.097*
H23B	0.3084	0.7075	0.5448	0.097*
H23C	0.3684	0.7073	0.4979	0.097*
H1A	-0.0843 (13)	0.456 (2)	0.5486 (11)	0.055 (6)*
H2A	-0.1957 (19)	0.562 (4)	0.5826 (15)	0.106 (9)*
H3A	-0.2190 (17)	0.434 (3)	0.6873 (13)	0.088 (7)*
H4A	-0.1115 (17)	0.210 (3)	0.7549 (14)	0.091 (8)*
H5A	0.0095 (14)	0.115 (2)	0.7224 (11)	0.062 (6)*
H10A	0.2610 (12)	0.469 (2)	0.5891 (10)	0.052 (5)*
H15A	0.6715 (13)	0.724 (2)	0.7285 (11)	0.064 (6)*
H16A	0.7790 (17)	0.924 (3)	0.7169 (14)	0.092 (8)*
H17A	0.7596 (15)	1.023 (3)	0.5978 (12)	0.082 (7)*
H18A	0.6394 (13)	0.892 (2)	0.4898 (12)	0.067 (6)*
H19A	0.5326 (13)	0.682 (2)	0.4972 (11)	0.060 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0634 (9)	0.0410 (8)	0.1185 (13)	0.0000 (7)	0.0432 (9)	0.0135 (8)
N1	0.0454 (9)	0.0419 (8)	0.0686 (11)	-0.0020 (7)	0.0289 (8)	0.0045 (7)
N2	0.0494 (9)	0.0372 (8)	0.0616 (10)	-0.0071 (7)	0.0304 (8)	-0.0037 (7)

N3	0.0456 (9)	0.0494 (9)	0.0511 (9)	-0.0080 (7)	0.0263 (7)	-0.0023 (7)
N4	0.0458 (9)	0.0490 (9)	0.0560 (10)	-0.0008 (7)	0.0212 (7)	0.0111 (7)
N5	0.0415 (9)	0.0408 (8)	0.0467 (9)	-0.0013 (6)	0.0211 (7)	0.0051 (7)
C1	0.0535 (14)	0.0628 (14)	0.0775 (17)	0.0058 (11)	0.0188 (12)	0.0043 (12)
C2	0.0536 (15)	0.0703 (17)	0.113 (2)	0.0143 (13)	0.0180 (16)	-0.0166 (16)
C3	0.0572 (16)	0.093 (2)	0.109 (2)	-0.0042 (15)	0.0399 (16)	-0.0380 (18)
C4	0.0617 (15)	0.0759 (16)	0.0876 (18)	-0.0125 (13)	0.0400 (14)	-0.0276 (14)
C5	0.0496 (12)	0.0563 (12)	0.0661 (14)	-0.0056 (10)	0.0274 (11)	-0.0097 (11)
C6	0.0360 (10)	0.0458 (10)	0.0670 (13)	-0.0029 (8)	0.0189 (9)	-0.0075 (9)
C7	0.0427 (10)	0.0460 (10)	0.0557 (12)	-0.0055 (9)	0.0198 (9)	0.0049 (9)
C8	0.0464 (10)	0.0434 (10)	0.0523 (11)	-0.0062 (8)	0.0287 (9)	-0.0038 (8)
C9	0.0420 (10)	0.0455 (9)	0.0436 (10)	-0.0062 (8)	0.0216 (8)	-0.0023 (8)
C10	0.0465 (11)	0.0479 (10)	0.0465 (11)	-0.0042 (9)	0.0229 (9)	0.0013 (9)
C11	0.0428 (10)	0.0435 (9)	0.0420 (10)	-0.0061 (8)	0.0237 (8)	-0.0024 (8)
C12	0.0503 (11)	0.0479 (10)	0.0478 (11)	-0.0041 (9)	0.0243 (9)	0.0044 (8)
C13	0.0417 (10)	0.0425 (9)	0.0419 (10)	-0.0022 (7)	0.0218 (8)	-0.0005 (8)
C14	0.0415 (10)	0.0381 (9)	0.0508 (11)	-0.0025 (8)	0.0254 (9)	-0.0003 (8)
C15	0.0572 (12)	0.0649 (13)	0.0492 (13)	-0.0139 (10)	0.0200 (10)	0.0020 (10)
C16	0.0663 (15)	0.0719 (15)	0.0697 (17)	-0.0271 (12)	0.0187 (13)	-0.0036 (13)
C17	0.0657 (14)	0.0549 (12)	0.0818 (17)	-0.0181 (11)	0.0345 (13)	0.0019 (12)
C18	0.0769 (15)	0.0496 (11)	0.0662 (15)	-0.0079 (10)	0.0423 (13)	0.0047 (11)
C19	0.0615 (13)	0.0459 (10)	0.0501 (12)	-0.0115 (9)	0.0298 (10)	-0.0034 (9)
C20	0.0618 (13)	0.0536 (11)	0.0711 (14)	-0.0219 (10)	0.0322 (11)	-0.0095 (10)
C21	0.0783 (15)	0.0461 (12)	0.125 (2)	-0.0014 (11)	0.0680 (15)	-0.0048 (12)
C22	0.0644 (14)	0.0625 (13)	0.0870 (17)	0.0023 (11)	0.0306 (12)	0.0302 (12)
C23	0.0523 (12)	0.0575 (12)	0.0835 (15)	0.0062 (10)	0.0309 (11)	0.0151 (11)

Geometric parameters (Å, °)

O1—C7	1.227 (2)	C11—C13	1.386 (2)
N1—C7	1.404 (2)	C11—C12	1.422 (2)
N1—N2	1.4082 (19)	C12—C22	1.496 (2)
N1—C6	1.423 (2)	C13—C23	1.487 (2)
N2—C8	1.375 (2)	C14—C19	1.375 (2)
N2—C20	1.467 (2)	C14—C15	1.378 (3)
N3—C10	1.287 (2)	C15—C16	1.379 (3)
N3—C9	1.400 (2)	C15—H15A	0.978 (19)
N4—C12	1.325 (2)	C16—C17	1.368 (3)
N4—N5	1.3702 (19)	C16—H16A	0.92 (2)
N5—C13	1.357 (2)	C17—C18	1.374 (3)
N5—C14	1.429 (2)	C17—H17A	0.96 (2)
C1—C6	1.378 (3)	C18—C19	1.383 (3)
C1—C2	1.396 (4)	C18—H18A	0.95 (2)
C1—H1A	0.927 (18)	C19—H19A	0.990 (19)
C2—C3	1.368 (4)	C20—H20A	0.9600
C2—H2A	0.90 (3)	C20—H20B	0.9600
C3—C4	1.372 (4)	C20—H20C	0.9600
C3—H3A	0.98 (2)	C21—H21A	0.9600
C4—C5	1.372 (3)	C21—H21B	0.9600

supplementary materials

C4—H4A	0.97 (2)	C21—H21C	0.9600
C5—C6	1.388 (3)	C22—H22A	0.9600
C5—H5A	0.984 (19)	C22—H22B	0.9600
C7—C9	1.439 (2)	C22—H22C	0.9600
C8—C9	1.366 (2)	C23—H23A	0.9600
C8—C21	1.483 (2)	C23—H23B	0.9600
C10—C11	1.442 (2)	C23—H23C	0.9600
C10—H10A	0.982 (17)		
C7—N1—N2	109.36 (13)	C11—C12—C22	129.15 (16)
C7—N1—C6	123.92 (15)	N5—C13—C11	106.51 (14)
N2—N1—C6	118.45 (14)	N5—C13—C23	122.19 (15)
C8—N2—N1	106.59 (13)	C11—C13—C23	131.30 (16)
C8—N2—C20	122.69 (15)	C19—C14—C15	120.63 (17)
N1—N2—C20	116.40 (14)	C19—C14—N5	120.55 (16)
C10—N3—C9	120.18 (15)	C15—C14—N5	118.79 (16)
C12—N4—N5	105.24 (14)	C14—C15—C16	118.9 (2)
C13—N5—N4	112.08 (13)	C14—C15—H15A	118.6 (11)
C13—N5—C14	128.42 (14)	C16—C15—H15A	122.4 (11)
N4—N5—C14	119.27 (13)	C17—C16—C15	121.4 (2)
C6—C1—C2	118.4 (3)	C17—C16—H16A	120.3 (15)
C6—C1—H1A	119.1 (12)	C15—C16—H16A	118.3 (15)
C2—C1—H1A	122.5 (12)	C16—C17—C18	119.2 (2)
C3—C2—C1	121.1 (3)	C16—C17—H17A	122.6 (13)
C3—C2—H2A	125.9 (17)	C18—C17—H17A	118.2 (13)
C1—C2—H2A	113.0 (18)	C17—C18—C19	120.6 (2)
C2—C3—C4	119.6 (3)	C17—C18—H18A	121.0 (12)
C2—C3—H3A	124.3 (14)	C19—C18—H18A	118.4 (12)
C4—C3—H3A	116.1 (14)	C14—C19—C18	119.34 (19)
C5—C4—C3	120.6 (3)	C14—C19—H19A	119.2 (10)
C5—C4—H4A	115.9 (14)	C18—C19—H19A	121.5 (10)
C3—C4—H4A	123.5 (14)	N2—C20—H20A	109.5
C4—C5—C6	119.8 (2)	N2—C20—H20B	109.5
C4—C5—H5A	122.7 (11)	H20A—C20—H20B	109.5
C6—C5—H5A	117.5 (11)	N2—C20—H20C	109.5
C1—C6—C5	120.4 (2)	H20A—C20—H20C	109.5
C1—C6—N1	118.7 (2)	H20B—C20—H20C	109.5
C5—C6—N1	120.90 (17)	C8—C21—H21A	109.5
O1—C7—N1	123.40 (17)	C8—C21—H21B	109.5
O1—C7—C9	131.56 (17)	H21A—C21—H21B	109.5
N1—C7—C9	104.99 (14)	C8—C21—H21C	109.5
C9—C8—N2	110.30 (15)	H21A—C21—H21C	109.5
C9—C8—C21	128.03 (16)	H21B—C21—H21C	109.5
N2—C8—C21	121.65 (15)	C12—C22—H22A	109.5
C8—C9—N3	121.95 (15)	C12—C22—H22B	109.5
C8—C9—C7	108.22 (15)	H22A—C22—H22B	109.5
N3—C9—C7	129.41 (15)	C12—C22—H22C	109.5
N3—C10—C11	122.19 (18)	H22A—C22—H22C	109.5
N3—C10—H10A	121.7 (10)	H22B—C22—H22C	109.5
C11—C10—H10A	116.1 (10)	C13—C23—H23A	109.5

C13—C11—C12	105.06 (14)	C13—C23—H23B	109.5
C13—C11—C10	125.06 (16)	H23A—C23—H23B	109.5
C12—C11—C10	129.81 (16)	C13—C23—H23C	109.5
N4—C12—C11	111.10 (15)	H23A—C23—H23C	109.5
N4—C12—C22	119.73 (16)	H23B—C23—H23C	109.5
C7—N1—N2—C8	-7.52 (18)	N1—C7—C9—C8	-3.59 (19)
C6—N1—N2—C8	-157.20 (15)	O1—C7—C9—N3	1.0 (3)
C7—N1—N2—C20	-148.51 (16)	N1—C7—C9—N3	-176.21 (16)
C6—N1—N2—C20	61.8 (2)	C9—N3—C10—C11	176.07 (15)
C12—N4—N5—C13	1.35 (18)	N3—C10—C11—C13	171.42 (16)
C12—N4—N5—C14	176.16 (14)	N3—C10—C11—C12	-12.0 (3)
C6—C1—C2—C3	-1.7 (4)	N5—N4—C12—C11	-1.32 (19)
C1—C2—C3—C4	1.7 (4)	N5—N4—C12—C22	-179.90 (16)
C2—C3—C4—C5	0.3 (4)	C13—C11—C12—N4	0.84 (19)
C3—C4—C5—C6	-2.3 (3)	C10—C11—C12—N4	-176.23 (17)
C2—C1—C6—C5	-0.3 (3)	C13—C11—C12—C22	179.26 (19)
C2—C1—C6—N1	-179.76 (19)	C10—C11—C12—C22	2.2 (3)
C4—C5—C6—C1	2.3 (3)	N4—N5—C13—C11	-0.84 (18)
C4—C5—C6—N1	-178.27 (17)	C14—N5—C13—C11	-175.06 (15)
C7—N1—C6—C1	62.0 (2)	N4—N5—C13—C23	179.18 (16)
N2—N1—C6—C1	-153.01 (17)	C14—N5—C13—C23	5.0 (3)
C7—N1—C6—C5	-117.4 (2)	C12—C11—C13—N5	0.01 (17)
N2—N1—C6—C5	27.5 (2)	C10—C11—C13—N5	177.27 (15)
N2—N1—C7—O1	-170.72 (18)	C12—C11—C13—C23	179.99 (19)
C6—N1—C7—O1	-23.1 (3)	C10—C11—C13—C23	-2.8 (3)
N2—N1—C7—C9	6.80 (18)	C13—N5—C14—C19	-63.0 (2)
C6—N1—C7—C9	154.47 (17)	N4—N5—C14—C19	123.15 (18)
N1—N2—C8—C9	5.20 (19)	C13—N5—C14—C15	118.9 (2)
C20—N2—C8—C9	143.14 (17)	N4—N5—C14—C15	-55.0 (2)
N1—N2—C8—C21	-176.14 (18)	C19—C14—C15—C16	1.1 (3)
C20—N2—C8—C21	-38.2 (3)	N5—C14—C15—C16	179.18 (18)
N2—C8—C9—N3	172.28 (14)	C14—C15—C16—C17	-0.1 (3)
C21—C8—C9—N3	-6.3 (3)	C15—C16—C17—C18	-0.9 (4)
N2—C8—C9—C7	-1.0 (2)	C16—C17—C18—C19	0.9 (3)
C21—C8—C9—C7	-179.6 (2)	C15—C14—C19—C18	-1.1 (3)
C10—N3—C9—C8	172.48 (17)	N5—C14—C19—C18	-179.16 (16)
C10—N3—C9—C7	-15.8 (3)	C17—C18—C19—C14	0.1 (3)
O1—C7—C9—C8	173.6 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 and Cg2 are the centroids of the N4/N5/C11–C13 and C1–C6 rings, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C10—H10A \cdots O1	0.986 (18)	2.40 (2)	3.052 (3)	123.3 (14)
C19—H19A \cdots Cg2 ⁱ	0.990 (19)	2.656 (19)	3.452 (2)	137.4 (17)
C20—H20C \cdots Cg1 ⁱⁱ	0.96	2.85 (3)	3.720 (3)	149.(1)
C22—H22B \cdots Cg2 ⁱⁱⁱ	0.96	2.82 (3)	3.585 (3)	135.(1)

Symmetry codes: (i) $-x+1, -y-1, -z-1$; (ii) $-x, -y-2, -z-1$; (iii) $-x+1, y-1/2, -z-1/2$.

Fig. 1




Fig. 2

