

3-Mesityl-2-oxo-1-oxaspiro[4.4]non-3-en-4-yl benzoate

Chuan-ming Yu,^a Zong-cheng Wang,^a Huan Zhou,^b Ming-hua Ji^c and Jin-hao Zhao^{c*}

^aCollege of Pharmaceutical Sciences, Zhejiang University of Technology, Hangzhou 310032, People's Republic of China, ^bResearch Center of Analysis and Measurement, Zhejiang University of Technology, Hangzhou 310032, People's Republic of China, and ^cInstitute of Pesticide and Environmental Toxicology, Zhejiang University, Hangzhou 310029, People's Republic of China
Correspondence e-mail: jinhaozhao@zju.edu.cn


Received 18 May 2010; accepted 4 June 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; disorder in main residue; R factor = 0.051; wR factor = 0.164; data-to-parameter ratio = 17.4.

In the title compound, $C_{24}H_{24}O_4$, a derivative of the potent insecticide and miticide spiromesifen, one cyclopentane C atom is disordered over two positions with occupancies of 0.574 (12) and 0.426 (12), resulting in respective envelope and twisted conformations for the cyclopentane ring. The atom at the flap position is 0.620 (5) Å out of the mean plane formed by the other four atoms of the envelope form. The furan ring makes dihedral angles of 68.26 (3) and 69.38 (2)°, respectively, with the 2,4,6-trimethylphenyl and benzene rings. The dihedral angle between the two benzene rings is 62.27 (3)°.

Related literature

For the pesticide spiromesifen, the central unit of the title compound, see: Bayer Aktiengesellschaft (1995). For the synthesis and biological activity of spiromesifen derivatives, see: Ji *et al.* (2009); Zhao *et al.* (2009). For distance restraints, see: Watkin (1994).



Experimental

Crystal data

$C_{24}H_{24}O_4$
 $M_r = 376.43$
Monoclinic, $P2_1/c$
 $a = 8.4799$ (5) Å
 $b = 15.9912$ (9) Å
 $c = 15.9520$ (8) Å
 $\beta = 106.240$ (1)°

$V = 2076.8$ (2) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.08\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.57 \times 0.45 \times 0.32\text{ mm}$

Data collection

Rigaku R-Axis RAPID diffractometer
Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.955$, $T_{\max} = 0.975$

19157 measured reflections
4647 independent reflections
2622 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.164$
 $S = 1.00$
4647 reflections
267 parameters

22 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.25\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$

Data collection: *PROCESS-AUTO* (Rigaku, 2006); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

The authors thank Professor Jian-Ming Gu for help with the analysis of the crystal data. This work was supported by the National Natural Science Foundation of China (No. 30700532) and the Science and Technology Project of Zhejiang Province (No. 2009 C21014).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SI2265).

References

- Bayer Aktiengesellschaft (1995). WO Patent No. 9504719A1.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Ji, M.-H., Xia, M., Zhu, G.-N. & Zhao, J.-H. (2009). *Acta Cryst. E* **65**, o705.
Rigaku (2006). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
Rigaku (2007). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
Watkin, D. (1994). *Acta Cryst. A* **50**, 411–437.
Zhao, J. H., Ji, M. H., Xu, X. H., Cheng, J. L. & Zhu, G. N. (2009). *Chin. Chem. Lett.* **20**, 1307–1310.

supplementary materials

Acta Cryst. (2010). E66, o1624 [doi:10.1107/S1600536810021343]

3-Mesyl-2-oxo-1-oxaspiro[4.4]non-3-en-4-yl benzoate

C. Yu, Z. Wang, H. Zhou, M. Ji and J. Zhao

Comment

4-hydroxyl-3-(2,4,6-trimethylphenyl)-1-oxaspiro[4,4]non-3-en-2-one (HTPO) is a key intermediate of Spiromesifen, which is an efficient insecticide and miticide, developed by Bayer Aktiengesellschaft (1995), see (Ji *et al.* 2009). As part of our continuing interest in the design and synthesis of the new insecticide and miticide, we have isolated the title compound (I), by the condensation reaction of benzoyl chloride and HTPO as colorless crystals. The molecule of the title compound (Fig. 1), exhibits a similar conformation and the same double bond characteristics as reported for the chlorobenzoate structure (Ji *et al.* 2009).

There was an indication of positional disorder in the crystal structure, detected with the checkCIF structure validation Program *PLATON* (Spek, 2009), which showed in the alert level B section significant Hirshfeld rigid bond test differences, 10.75 s.u. for C23—C24. Atom C23 was split into atoms C23A and C23B, which were refined using SIMU and PART instructions of *SHELXL97*, and four distance restraints were applied (Watkin, 1994). The C22—C23A, C22—C23B, C24—C23A and C24—C23B bond distance were restrained to 1.510 (2) Å, 1.514 (2) Å, 1.517 (2) Å and 1.5151 (19) Å, respectively.

The cyclopentane ring with C23B displays an envelope conformation with the C24 atom at the flap position 0.620 (5) Å out of the mean plane formed by the other four atoms, whereas the cyclopentane ring with C23A displays a twisted conformation, twisted on C21—C22.

Experimental

4-hydroxyl-3-(2,4,6-trimethylphenyl)-1-oxaspiro[4,4]non-3-en-2-one (0.272 g, 1 mmol), 4-dimethylaminopyridine (0.012 g, 0.1 mmol), triethylamine (0.131 g, 1.3 mmol) and dry chloroform (10 ml) were added to a 25 ml round flask. Then the mixture was stirred and cooled to 273 K. Within 30 min benzoyl chloride (0.168 g, 1.2 mmol) was added dropwise to the solution at 273 K. After the reaction mixture was reacted at room temperature for 3 h, 1% HCl was added. The organic layer was washed to neutral with water and dried over Na₂SO₄. After filtered and concentrated, the organic residue was purified by silica gel column chromatography, eluted with ethyl acetate-petrum (1:3, v/v) to give a white solid (yield 83%, 0.312 g), which was then recrystallized from 95% ethanol to give colourless blocks.

Refinement

H atoms were included in calculated positions and refined using a riding model, with C—H distances constrained to 0.96 Å for methyl H atoms, 0.93 Å for aryl H atoms and 0.98 Å for the remainder, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms.

supplementary materials

Figures




Fig. 1. The molecular structure of (I), showing the atom-labelling scheme and the disordered CH₂ group indicated as dashed lines. Displacement ellipsoids are drawn at the 50% probability level.

3-Mesityl-2-oxo-1-oxaspiro[4.4]non-3-en-4-yl benzoate

Crystal data

C ₂₄ H ₂₄ O ₄	$F(000) = 800$
$M_r = 376.43$	$D_x = 1.204 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 10851 reflections
$a = 8.4799 (5) \text{ \AA}$	$\theta = 3.1\text{--}27.4^\circ$
$b = 15.9912 (9) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$c = 15.9520 (8) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 106.240 (1)^\circ$	Chunk, colorless
$V = 2076.8 (2) \text{ \AA}^3$	$0.57 \times 0.45 \times 0.32 \text{ mm}$
$Z = 4$	

Data collection

Rigaku R-AXIS RAPID diffractometer	4647 independent reflections
Radiation source: rolling anode graphite	2622 reflections with $I > 2\sigma(I)$
Detector resolution: 10.00 pixels mm ⁻¹	$R_{\text{int}} = 0.031$
ω scans	$\theta_{\text{max}} = 27.4^\circ$, $\theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.955$, $T_{\text{max}} = 0.975$	$k = -20 \rightarrow 17$
19157 measured reflections	$l = -17 \rightarrow 20$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.051$	H-atom parameters constrained
$wR(F^2) = 0.164$	$w = 1/[\sigma^2(F_o^2) + (0.0597P)^2 + 0.8207P]$
	where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.00$	$(\Delta/\sigma)_{\max} < 0.001$
4647 reflections	$\Delta\rho_{\max} = 0.25 \text{ e \AA}^{-3}$
267 parameters	$\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$
22 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.035 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O3	0.59902 (17)	0.68318 (9)	0.43446 (9)	0.0620 (4)	
O4	0.8496 (2)	0.73853 (11)	0.45217 (12)	0.0889 (6)	
O1	0.3476 (2)	0.84301 (10)	0.30647 (12)	0.0840 (5)	
O2	0.3938 (3)	0.83890 (14)	0.17466 (12)	0.1114 (8)	
C14	0.8250 (2)	0.61060 (12)	0.52516 (12)	0.0512 (5)	
C3	0.6519 (3)	0.70257 (13)	0.24733 (13)	0.0598 (5)	
C19	0.7173 (3)	0.54964 (13)	0.53702 (14)	0.0639 (6)	
H19	0.6052	0.5551	0.5106	0.077*	
C15	0.9908 (3)	0.60287 (13)	0.56565 (13)	0.0579 (5)	
H15	1.0632	0.6440	0.5583	0.070*	
C13	0.7675 (3)	0.68381 (13)	0.46840 (13)	0.0566 (5)	
C12	0.5273 (3)	0.73987 (13)	0.37080 (13)	0.0577 (5)	
C8	0.6291 (3)	0.61735 (14)	0.22738 (13)	0.0605 (5)	
C7	0.7338 (3)	0.57817 (15)	0.18622 (14)	0.0668 (6)	
H7	0.7190	0.5216	0.1728	0.080*	
C2	0.5435 (3)	0.74710 (13)	0.29100 (14)	0.0624 (6)	
C17	0.9423 (3)	0.47286 (15)	0.62716 (15)	0.0714 (6)	
H17	0.9821	0.4257	0.6606	0.086*	
C5	0.8798 (3)	0.70432 (17)	0.18587 (15)	0.0743 (7)	
H5	0.9642	0.7334	0.1720	0.089*	
C18	0.7768 (3)	0.48087 (15)	0.58814 (16)	0.0734 (7)	
H18	0.7048	0.4398	0.5962	0.088*	
C16	1.0493 (3)	0.53393 (15)	0.61723 (14)	0.0668 (6)	
H16	1.1609	0.5289	0.6451	0.080*	
C20	0.4067 (3)	0.79956 (14)	0.38972 (15)	0.0645 (6)	

supplementary materials

C4	0.7791 (3)	0.74694 (15)	0.22727 (14)	0.0678 (6)	
C6	0.8588 (3)	0.62051 (16)	0.16464 (15)	0.0710 (6)	
C1	0.4252 (3)	0.81278 (16)	0.24846 (17)	0.0796 (7)	
C21	0.4808 (3)	0.86271 (16)	0.46176 (17)	0.0774 (7)	
H21A	0.5716	0.8379	0.5058	0.093*	
H21B	0.5206	0.9115	0.4378	0.093*	
C9	0.8105 (4)	0.83721 (16)	0.25219 (18)	0.0909 (8)	
H9A	0.9100	0.8550	0.2402	0.136*	
H9B	0.8210	0.8438	0.3133	0.136*	
H9C	0.7204	0.8705	0.2190	0.136*	
C11	0.4958 (3)	0.56714 (16)	0.24929 (17)	0.0778 (7)	
H11A	0.5270	0.5551	0.3107	0.117*	
H11B	0.4801	0.5157	0.2170	0.117*	
H11C	0.3954	0.5986	0.2342	0.117*	
C24	0.2621 (3)	0.76212 (15)	0.4166 (2)	0.0859 (8)	
H24A	0.1634	0.7648	0.3679	0.103*	
H24B	0.2842	0.7039	0.4326	0.103*	
C10	0.9716 (4)	0.5763 (2)	0.1205 (2)	0.1052 (10)	
H10A	0.9949	0.5211	0.1443	0.158*	
H10B	1.0722	0.6072	0.1303	0.158*	
H10C	0.9193	0.5726	0.0589	0.158*	
C22	0.3449 (4)	0.8866 (2)	0.5007 (2)	0.1037 (10)	
H22A	0.2826	0.9332	0.4690	0.124*	
H22B	0.3895	0.9027	0.5613	0.124*	
C23A	0.2365 (12)	0.8105 (4)	0.4935 (5)	0.095 (3)	0.426 (12)
H23A	0.1223	0.8266	0.4829	0.115*	0.426 (12)
H23B	0.2690	0.7774	0.5465	0.115*	0.426 (12)
C23B	0.1941 (4)	0.8384 (3)	0.4509 (5)	0.091 (2)	0.574 (12)
H23C	0.1270	0.8716	0.4033	0.109*	0.574 (12)
H23D	0.1283	0.8222	0.4892	0.109*	0.574 (12)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O3	0.0500 (9)	0.0678 (9)	0.0656 (9)	0.0025 (7)	0.0116 (7)	0.0185 (7)
O4	0.0676 (11)	0.0851 (12)	0.1018 (13)	-0.0220 (9)	0.0034 (9)	0.0370 (10)
O1	0.0854 (13)	0.0734 (11)	0.0842 (11)	0.0284 (9)	0.0089 (9)	0.0066 (9)
O2	0.1274 (18)	0.1159 (16)	0.0777 (13)	0.0473 (14)	0.0069 (11)	0.0328 (11)
C14	0.0498 (12)	0.0531 (11)	0.0494 (10)	-0.0009 (9)	0.0120 (8)	0.0015 (8)
C3	0.0631 (14)	0.0602 (13)	0.0522 (11)	0.0049 (10)	0.0097 (9)	0.0130 (9)
C19	0.0525 (13)	0.0618 (13)	0.0733 (14)	-0.0042 (10)	0.0110 (10)	0.0110 (10)
C15	0.0499 (12)	0.0638 (13)	0.0577 (12)	-0.0023 (10)	0.0111 (9)	-0.0029 (9)
C13	0.0493 (12)	0.0626 (13)	0.0546 (11)	-0.0041 (10)	0.0090 (9)	0.0063 (9)
C12	0.0559 (13)	0.0528 (12)	0.0594 (12)	0.0028 (9)	0.0081 (9)	0.0076 (9)
C8	0.0583 (13)	0.0627 (13)	0.0573 (12)	0.0021 (10)	0.0110 (10)	0.0091 (9)
C7	0.0689 (15)	0.0656 (14)	0.0642 (13)	0.0049 (11)	0.0159 (11)	0.0052 (10)
C2	0.0634 (14)	0.0563 (12)	0.0613 (13)	0.0079 (10)	0.0074 (10)	0.0083 (9)
C17	0.0786 (17)	0.0606 (14)	0.0675 (14)	0.0115 (12)	0.0083 (12)	0.0103 (10)

C5	0.0707 (16)	0.0876 (18)	0.0652 (14)	-0.0082 (13)	0.0197 (12)	0.0191 (12)
C18	0.0730 (16)	0.0621 (14)	0.0813 (16)	-0.0060 (12)	0.0152 (12)	0.0175 (11)
C16	0.0598 (14)	0.0698 (14)	0.0634 (13)	0.0083 (11)	0.0050 (10)	0.0021 (10)
C20	0.0617 (14)	0.0572 (12)	0.0703 (14)	0.0025 (10)	0.0114 (11)	-0.0010 (10)
C4	0.0781 (16)	0.0649 (14)	0.0572 (13)	-0.0026 (12)	0.0135 (11)	0.0158 (10)
C6	0.0692 (16)	0.0805 (17)	0.0647 (14)	0.0046 (13)	0.0209 (11)	0.0103 (11)
C1	0.0821 (18)	0.0734 (16)	0.0718 (16)	0.0207 (13)	0.0028 (13)	0.0111 (12)
C21	0.0757 (17)	0.0659 (15)	0.0873 (17)	-0.0098 (12)	0.0176 (13)	-0.0127 (12)
C9	0.118 (2)	0.0694 (16)	0.0866 (18)	-0.0153 (15)	0.0303 (16)	0.0134 (13)
C11	0.0752 (17)	0.0737 (16)	0.0869 (17)	-0.0075 (13)	0.0268 (13)	-0.0015 (12)
C24	0.0620 (16)	0.0648 (15)	0.132 (2)	-0.0049 (12)	0.0293 (15)	-0.0144 (15)
C10	0.095 (2)	0.121 (3)	0.115 (2)	0.0060 (19)	0.0548 (19)	-0.0022 (19)
C22	0.091 (2)	0.122 (3)	0.095 (2)	0.0028 (19)	0.0220 (16)	-0.0362 (18)
C23A	0.096 (3)	0.096 (3)	0.095 (3)	-0.0014 (10)	0.0283 (13)	0.0004 (10)
C23B	0.089 (2)	0.090 (2)	0.093 (2)	0.0003 (10)	0.0265 (11)	-0.0017 (10)

Geometric parameters (Å, °)

O3—C12	1.370 (2)	C16—H16	0.9300
O3—C13	1.379 (2)	C20—C24	1.530 (3)
O4—C13	1.191 (2)	C20—C21	1.527 (3)
O1—C1	1.365 (3)	C4—C9	1.501 (3)
O1—C20	1.458 (3)	C6—C10	1.512 (4)
O2—C1	1.207 (3)	C21—C22	1.503 (4)
C14—C15	1.379 (3)	C21—H21A	0.9700
C14—C19	1.385 (3)	C21—H21B	0.9700
C14—C13	1.477 (3)	C9—H9A	0.9600
C3—C4	1.401 (3)	C9—H9B	0.9600
C3—C8	1.400 (3)	C9—H9C	0.9600
C3—C2	1.482 (3)	C11—H11A	0.9600
C19—C18	1.378 (3)	C11—H11B	0.9600
C19—H19	0.9300	C11—H11C	0.9600
C15—C16	1.382 (3)	C24—C23B	1.5151 (19)
C15—H15	0.9300	C24—C23A	1.517 (2)
C12—C2	1.323 (3)	C24—H24A	0.9700
C12—C20	1.490 (3)	C24—H24B	0.9700
C8—C7	1.392 (3)	C10—H10A	0.9600
C8—C11	1.505 (3)	C10—H10B	0.9600
C7—C6	1.380 (3)	C10—H10C	0.9600
C7—H7	0.9300	C22—C23A	1.510 (2)
C2—C1	1.480 (3)	C22—C23B	1.514 (2)
C17—C18	1.372 (3)	C22—H22A	0.9700
C17—C16	1.372 (3)	C22—H22B	0.9700
C17—H17	0.9300	C23A—H23A	0.9700
C5—C6	1.382 (4)	C23A—H23B	0.9700
C5—C4	1.395 (3)	C23B—H23C	0.9700
C5—H5	0.9300	C23B—H23D	0.9700
C18—H18	0.9300		
C12—O3—C13	118.83 (16)	O1—C1—C2	109.5 (2)

supplementary materials

C1—O1—C20	109.94 (17)	C22—C21—C20	106.1 (2)
C15—C14—C19	119.88 (19)	C22—C21—H21A	110.5
C15—C14—C13	118.49 (18)	C20—C21—H21A	110.5
C19—C14—C13	121.62 (18)	C22—C21—H21B	110.5
C4—C3—C8	120.4 (2)	C20—C21—H21B	110.5
C4—C3—C2	118.9 (2)	H21A—C21—H21B	108.7
C8—C3—C2	120.7 (2)	C4—C9—H9A	109.5
C18—C19—C14	119.7 (2)	C4—C9—H9B	109.5
C18—C19—H19	120.1	H9A—C9—H9B	109.5
C14—C19—H19	120.1	C4—C9—H9C	109.5
C14—C15—C16	120.0 (2)	H9A—C9—H9C	109.5
C14—C15—H15	120.0	H9B—C9—H9C	109.5
C16—C15—H15	120.0	C8—C11—H11A	109.5
O4—C13—O3	121.64 (19)	C8—C11—H11B	109.5
O4—C13—C14	127.1 (2)	H11A—C11—H11B	109.5
O3—C13—C14	111.30 (17)	C8—C11—H11C	109.5
C2—C12—O3	128.8 (2)	H11A—C11—H11C	109.5
C2—C12—C20	113.61 (18)	H11B—C11—H11C	109.5
O3—C12—C20	117.45 (18)	C23B—C24—C20	101.7 (3)
C7—C8—C3	118.7 (2)	C23A—C24—C20	109.5 (2)
C7—C8—C11	119.3 (2)	C23B—C24—H24A	85.6
C3—C8—C11	122.1 (2)	C23A—C24—H24A	109.8
C6—C7—C8	122.2 (2)	C20—C24—H24A	109.8
C6—C7—H7	118.9	C23B—C24—H24B	138.0
C8—C7—H7	118.9	C23A—C24—H24B	109.8
C12—C2—C1	105.3 (2)	C20—C24—H24B	109.8
C12—C2—C3	130.94 (19)	H24A—C24—H24B	108.2
C1—C2—C3	123.8 (2)	C6—C10—H10A	109.5
C18—C17—C16	120.4 (2)	C6—C10—H10B	109.5
C18—C17—H17	119.8	H10A—C10—H10B	109.5
C16—C17—H17	119.8	C6—C10—H10C	109.5
C6—C5—C4	122.4 (2)	H10A—C10—H10C	109.5
C6—C5—H5	118.8	H10B—C10—H10C	109.5
C4—C5—H5	118.8	C21—C22—C23A	106.5 (3)
C17—C18—C19	120.1 (2)	C21—C22—C23B	106.7 (3)
C17—C18—H18	119.9	C21—C22—H22A	110.4
C19—C18—H18	119.9	C23A—C22—H22A	110.4
C17—C16—C15	119.8 (2)	C23B—C22—H22A	82.0
C17—C16—H16	120.1	C21—C22—H22B	110.4
C15—C16—H16	120.1	C23A—C22—H22B	110.4
O1—C20—C12	101.61 (17)	C23B—C22—H22B	134.0
O1—C20—C24	110.1 (2)	H22A—C22—H22B	108.6
C12—C20—C24	117.10 (19)	C22—C23A—C24	104.5 (2)
O1—C20—C21	109.55 (19)	C22—C23A—H23A	110.8
C12—C20—C21	114.51 (19)	C24—C23A—H23A	110.8
C24—C20—C21	103.96 (19)	C22—C23A—H23B	110.8
C5—C4—C3	118.2 (2)	C24—C23A—H23B	110.9
C5—C4—C9	120.6 (2)	H23A—C23A—H23B	108.9
C3—C4—C9	121.1 (2)	C22—C23B—C24	104.4 (2)

C7—C6—C5	118.0 (2)	C22—C23B—H23C	110.9
C7—C6—C10	121.1 (3)	C24—C23B—H23C	110.9
C5—C6—C10	120.8 (2)	C22—C23B—H23D	110.9
O2—C1—O1	121.5 (2)	C24—C23B—H23D	110.9
O2—C1—C2	129.0 (3)	H23C—C23B—H23D	108.9
C15—C14—C19—C18	-1.1 (3)	C2—C12—C20—C21	-116.3 (2)
C13—C14—C19—C18	177.7 (2)	O3—C12—C20—C21	67.0 (3)
C19—C14—C15—C16	0.7 (3)	C6—C5—C4—C3	-0.3 (3)
C13—C14—C15—C16	-178.18 (19)	C6—C5—C4—C9	177.6 (2)
C12—O3—C13—O4	9.7 (3)	C8—C3—C4—C5	1.0 (3)
C12—O3—C13—C14	-170.80 (17)	C2—C3—C4—C5	-179.77 (19)
C15—C14—C13—O4	0.7 (3)	C8—C3—C4—C9	-176.9 (2)
C19—C14—C13—O4	-178.1 (2)	C2—C3—C4—C9	2.3 (3)
C15—C14—C13—O3	-178.75 (18)	C8—C7—C6—C5	0.6 (3)
C19—C14—C13—O3	2.4 (3)	C8—C7—C6—C10	179.6 (2)
C13—O3—C12—C2	63.9 (3)	C4—C5—C6—C7	-0.5 (4)
C13—O3—C12—C20	-119.9 (2)	C4—C5—C6—C10	-179.4 (2)
C4—C3—C8—C7	-0.9 (3)	C20—O1—C1—O2	179.0 (3)
C2—C3—C8—C7	179.92 (19)	C20—O1—C1—C2	-0.7 (3)
C4—C3—C8—C11	178.9 (2)	C12—C2—C1—O2	-177.9 (3)
C2—C3—C8—C11	-0.3 (3)	C3—C2—C1—O2	2.1 (5)
C3—C8—C7—C6	0.1 (3)	C12—C2—C1—O1	1.7 (3)
C11—C8—C7—C6	-179.8 (2)	C3—C2—C1—O1	-178.3 (2)
O3—C12—C2—C1	174.2 (2)	O1—C20—C21—C22	93.4 (3)
C20—C12—C2—C1	-2.1 (3)	C12—C20—C21—C22	-153.2 (2)
O3—C12—C2—C3	-5.8 (4)	C24—C20—C21—C22	-24.2 (3)
C20—C12—C2—C3	177.9 (2)	O1—C20—C24—C23B	-77.9 (3)
C4—C3—C2—C12	-111.4 (3)	C12—C20—C24—C23B	166.7 (3)
C8—C3—C2—C12	67.8 (3)	C21—C20—C24—C23B	39.3 (3)
C4—C3—C2—C1	68.6 (3)	O1—C20—C24—C23A	-109.0 (5)
C8—C3—C2—C1	-112.2 (3)	C12—C20—C24—C23A	135.6 (5)
C16—C17—C18—C19	1.4 (4)	C21—C20—C24—C23A	8.2 (5)
C14—C19—C18—C17	0.0 (4)	C20—C21—C22—C23A	31.8 (5)
C18—C17—C16—C15	-1.8 (4)	C20—C21—C22—C23B	-0.5 (4)
C14—C15—C16—C17	0.8 (3)	C21—C22—C23A—C24	-25.9 (8)
C1—O1—C20—C12	-0.5 (2)	C23B—C22—C23A—C24	69.0 (3)
C1—O1—C20—C24	-125.3 (2)	C23B—C24—C23A—C22	-69.3 (3)
C1—O1—C20—C21	121.0 (2)	C20—C24—C23A—C22	10.7 (8)
C2—C12—C20—O1	1.7 (2)	C21—C22—C23B—C24	25.4 (6)
O3—C12—C20—O1	-175.02 (17)	C23A—C22—C23B—C24	-69.1 (3)
C2—C12—C20—C24	121.7 (2)	C23A—C24—C23B—C22	68.8 (3)
O3—C12—C20—C24	-55.1 (3)	C20—C24—C23B—C22	-39.8 (5)

supplementary materials

Fig. 1

