

4-[(*E*)-(2,4,5-Trimethoxybenzylidene)-amino]-1,5-dimethyl-2-phenyl-1*H*-pyrazol-3(2*H*)-one

Hoong-Kun Fun,^{a,*‡} Madhukar Hemamalini,^a Abdullah M. Asiri^{b§} and Salman A. Khan^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Chemistry, Faculty of Science, King Abdu Aziz University, Jeddah, Saudi Arabia
Correspondence e-mail: hkfun@usm.my


Received 1 June 2010; accepted 7 June 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.001\text{ \AA}$; R factor = 0.040; wR factor = 0.123; data-to-parameter ratio = 16.3.

The title compound, $C_{21}H_{23}N_3O_4$, adopts an *E* configuration about the central $\text{C}=\text{N}$ double bond and the pyrazolone ring is almost planar, with a maximum deviation of $0.042(1)\text{ \AA}$. The central pyrazolone ring makes dihedral angles of $51.96(5)$ and $3.82(5)^\circ$ with the attached phenyl and the trimethoxy-substituted benzene rings, respectively. The dihedral angle between the phenyl ring and the trimethoxy-substituted benzene ring is $50.19(5)^\circ$ and an intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond generates an *S*(6) ring motif. The crystal structure is stabilized by intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds.

Related literature

For background to the applications of Schiff bases, see: Vukovic *et al.* (2010); Ramesh & Maheswaran (2003); Dongfang *et al.* (2008); Sastry & Rao (1988); Kamel *et al.* (2010). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



‡ Thomson Reuters ResearcherID: A-3561-2009.

§ On secondment to: The Center of Excellence for Advanced Materials Research, King Abdu Aziz University, Jeddah 21589, Saudi Arabia.

Experimental

Crystal data

$C_{21}H_{23}N_3O_4$
 $M_r = 381.42$
Monoclinic, $P2_1/c$
 $a = 21.0128(10)\text{ \AA}$
 $b = 7.4242(4)\text{ \AA}$
 $c = 12.5194(6)\text{ \AA}$
 $\beta = 98.675(1)^\circ$

$V = 1930.72(17)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.67 \times 0.27 \times 0.15\text{ mm}$

Data collection

Bruker APEXII DUO CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2009)
 $T_{\min} = 0.941$, $T_{\max} = 0.987$

23600 measured reflections
5614 independent reflections
4779 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.123$
 $S = 1.04$
5614 reflections
345 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.48\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.23\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C10—H10A…O1	0.954 (13)	2.331 (13)	3.0112 (11)	127.8 (10)
C4—H4A…O1 ⁱ	0.969 (13)	2.541 (13)	3.2628 (12)	131.4 (10)
C20—H20A…N3 ⁱⁱ	0.996 (14)	2.577 (14)	3.5383 (13)	162.1 (12)
C20—H20C…O2 ⁱⁱⁱ	0.977 (14)	2.509 (14)	3.4470 (13)	160.8 (12)
C20—H20C…O3 ⁱⁱⁱ	0.977 (14)	2.495 (15)	3.2779 (13)	137.0 (11)

Symmetry codes: (i) $x, -y - \frac{1}{2}, z + \frac{1}{2}$; (ii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (iii) $-x, -y + 1, -z$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and MH thank the Malaysian Government and Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012. MH also thanks Universiti Sains Malaysia for a post-doctoral research fellowship. AMA and SAK thank the Chemistry Department, King Abdul Aziz University, Jeddah, Saudi Arabia, for providing research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5480).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
Dongfang, X. U., Shuzhi, M. A., Guangying, D. U., Qizhuang, H. E. & Dazhi, S. (2008). *J. Rare Earths*, **26**, 643–647.
Kamel, M. M., Ali, H. I., Anwar, M. M., Mohamed, N. A. & Soliman, A. M. (2010). *Eur. J. Med. Chem.* **45**, 572–580.

- Ramesh, R. & Maheswaran, S. (2003). *J. Inorg. Biochem.*, **96**, 457–462.
Sastry, C. S. P. & Rao, A. R. M. (1988). *J. Pharmacol. Methods*, **26**, 643–647.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
Vukovic, N., Sukdolak, S., Solujic, S. & Niciforovic, N. (2010). *Food Chem.* **120**, 1011–1018.

supplementary materials

Acta Cryst. (2010). E66, o1656-o1657 [doi:10.1107/S1600536810021586]

4-[*(E*)-(2,4,5-Trimethoxybenzylidene)amino]-1,5-dimethyl-2-phenyl-1*H*-pyrazol-3(2*H*)-one

H.-K. Fun, M. Hemamalini, A. M. Asiri and S. A. Khan

Comment

Compounds with the structure of AC=NB are known as Schiff base, which can be synthesized from the condensation of amino and active carbonyl groups. Schiff base compounds have shown different therapeutic properties such as antibacterial (Vukovic *et al.*, 2010), antifungal (Ramesh & Maheswaran, 2003), antitumor (Dongfang *et al.*, 2008), anti-inflammatory (Sastry & Rao, 1988) and anticancer activities (Kamel *et al.*, 2010). Due to their importance, the crystal structure determination of the title compound was carried out and the results are presented here.

In the title compound (Fig. 1), the pyrazolone ring (N1/N2/C7–C9) is almost planar, with maximum deviation of 0.042 (1) Å for atom N2. The central pyrazolone (N1/N2/C7–C9) ring makes dihedral angles of 51.96 (5)° and 3.82 (5)° with the attached phenyl ring (C1–C6) and the trimethoxy substituted phenyl ring (C11–C16), respectively. The dihedral angle between the phenyl ring(C1–C6) and the trimethoxy substituted phenyl ring (C11–C16) is 50.19 (5)°. The three methoxy groups are coplanar with the benzene ring [torsion angles C19–O2–C13–C12 = 5.04 (16)°, C20–O3–C14–C15 = -0.36 (14)° and C21–O4–C16–C15 = -1.66 (13)°].

In the crystal packing (Fig. 2), the intramolecular C10—H10A···O1 hydrogen bonding generates an *S*(6) ring motif (Bernstein *et al.*, 1995). The crystal sturcture is futher stabilized by weak intermolecular C4—H4A···O1, C20—H20C···O2, C20—H20C···O3 and C20—H20A···N3 (Table 1) hydrogen bonds.

Experimental

A mixture of 4-aminophenazone (0.50 g, 0.0033 mol) and 2,4,5-tri-methoxy- benzaldehyde (0.65 g, 0.0033 mol) in methanol (15 ml) was refluxed for 5 h with stirring to give a light yellow precipitate. It was then filtered and washed with methanol to give the pure Schiff base and yellow blocks of (I) were recrystallized from methanol. Yield: 48.18%; Mp. 381°C; IR (KBr) ν_{max} cm⁻¹: 2937 (C—H), 1644 (C=C), 1609(C=O), 1591 (C=N), 1122 (N—N). ¹H-NMR (CDCl_3) δ: 10.02 ((s, 1H, CH olefinic), 7.67 (s, H3, CHaromatic), 6.49 (s, H6, CHaromatic), 7.47–7.26 (m, 5H, CHaromatic), 3.93 (s, OCH₃), 3.93 (s, OCH₃), 3.84 (s, OCH₃), 3.11(s, N-CH₃), 2.48 (s,-CH₃).

Refinement

All the H atoms were located from a difference Fourier map and refined freely [C—H = 0.945 (14)–1.008 (14) Å].

supplementary materials

Figures




Fig. 1. The molecular structure of (I) showing 50% probability displacement ellipsoids. The intramolecular hydrogen bond is shown as a dashed line.




Fig. 2. The crystal packing of (I) showing hydrogen-bonded (dashed lines) networks. H atoms not involved in the hydrogen bond interactions are omitted for clarity.

4-[(*E*)-(2,4,5-Trimethoxybenzylidene)amino]-1,5-dimethyl-2-phenyl- 1*H*-pyrazol-3(2*H*)-one

Crystal data

C ₂₁ H ₂₃ N ₃ O ₄	<i>F</i> (000) = 808
<i>M_r</i> = 381.42	<i>D_x</i> = 1.312 Mg m ⁻³
Monoclinic, <i>P</i> 2 ₁ /c	Mo <i>K</i> α radiation, λ = 0.71073 Å
Hall symbol: -P 2ybc	Cell parameters from 8559 reflections
<i>a</i> = 21.0128 (10) Å	θ = 2.9–34.8°
<i>b</i> = 7.4242 (4) Å	μ = 0.09 mm ⁻¹
<i>c</i> = 12.5194 (6) Å	<i>T</i> = 100 K
β = 98.675 (1)°	Blcok, yellow
<i>V</i> = 1930.72 (17) Å ³	0.67 × 0.27 × 0.15 mm
<i>Z</i> = 4	

Data collection

Bruker APEXII DUO CCD diffractometer	5614 independent reflections
Radiation source: fine-focus sealed tube	4779 reflections with $I > 2\sigma(I)$
graphite	R_{int} = 0.031
φ and ω scans	$\theta_{\text{max}} = 30.0^\circ$, $\theta_{\text{min}} = 1.0^\circ$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009)	$h = -29 \rightarrow 29$
$T_{\text{min}} = 0.941$, $T_{\text{max}} = 0.987$	$k = -10 \rightarrow 10$
23600 measured reflections	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.040$	Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.123$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0771P)^2 + 0.3259P]$
5614 reflections	where $P = (F_o^2 + 2F_c^2)/3$
345 parameters	$(\Delta/\sigma)_{\max} < 0.001$
0 restraints	$\Delta\rho_{\max} = 0.48 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.36912 (3)	-0.05337 (10)	0.48796 (5)	0.01654 (15)
O2	0.02634 (3)	0.14809 (11)	0.14482 (6)	0.02325 (17)
O3	0.06645 (3)	0.44390 (11)	0.06868 (6)	0.02140 (17)
O4	0.27842 (3)	0.35928 (10)	0.27787 (6)	0.01722 (15)
N1	0.35715 (4)	-0.34299 (11)	0.55298 (6)	0.01452 (16)
N2	0.30506 (4)	-0.45837 (11)	0.56225 (6)	0.01459 (16)
N3	0.22306 (4)	-0.11616 (11)	0.39236 (6)	0.01426 (16)
C1	0.47358 (5)	-0.34082 (14)	0.59345 (8)	0.01773 (19)
C2	0.52970 (5)	-0.35703 (15)	0.66770 (9)	0.0216 (2)
C3	0.52678 (5)	-0.39395 (14)	0.77564 (8)	0.0205 (2)
C4	0.46740 (5)	-0.41126 (14)	0.81062 (8)	0.01797 (19)
C5	0.41087 (5)	-0.39230 (14)	0.73777 (7)	0.01641 (18)
C6	0.41448 (4)	-0.35847 (13)	0.62948 (7)	0.01427 (18)
C7	0.25240 (4)	-0.38702 (13)	0.49748 (7)	0.01383 (18)
C8	0.26770 (4)	-0.22283 (13)	0.45728 (7)	0.01270 (17)
C9	0.33566 (4)	-0.18791 (13)	0.49592 (7)	0.01293 (17)
C10	0.24207 (4)	0.03035 (13)	0.35116 (7)	0.01384 (17)
C11	0.19674 (4)	0.14168 (13)	0.27961 (7)	0.01385 (18)
C12	0.13242 (4)	0.08699 (14)	0.24831 (7)	0.01533 (18)
C13	0.08971 (4)	0.18970 (14)	0.17891 (8)	0.01645 (18)
C14	0.11121 (4)	0.35193 (14)	0.13731 (7)	0.01642 (19)
C15	0.17427 (4)	0.40938 (14)	0.16780 (7)	0.01589 (18)
C16	0.21674 (4)	0.30561 (13)	0.24024 (7)	0.01424 (17)

supplementary materials

C17	0.32012 (5)	-0.65126 (14)	0.55935 (8)	0.0194 (2)
C18	0.18985 (5)	-0.48412 (14)	0.47905 (8)	0.01728 (19)
C19	0.00170 (5)	-0.00727 (18)	0.19178 (11)	0.0293 (3)
C20	0.08640 (5)	0.61077 (16)	0.02644 (9)	0.0223 (2)
C21	0.29988 (5)	0.52608 (14)	0.23842 (8)	0.01750 (19)
H1A	0.4749 (7)	-0.319 (2)	0.5178 (11)	0.025 (3)*
H2A	0.5716 (7)	-0.342 (2)	0.6413 (11)	0.027 (4)*
H3A	0.5674 (7)	-0.413 (2)	0.8265 (12)	0.028 (4)*
H4A	0.4646 (6)	-0.4380 (19)	0.8855 (11)	0.020 (3)*
H5A	0.3683 (7)	-0.407 (2)	0.7601 (11)	0.024 (3)*
H10A	0.2859 (6)	0.0685 (18)	0.3651 (10)	0.016 (3)*
H12A	0.1206 (7)	-0.025 (2)	0.2767 (11)	0.022 (3)*
H15A	0.1878 (7)	0.520 (2)	0.1400 (11)	0.020 (3)*
H17A	0.2798 (7)	-0.723 (2)	0.5642 (11)	0.024 (3)*
H17B	0.3365 (7)	-0.684 (2)	0.4918 (12)	0.027 (4)*
H17C	0.3507 (7)	-0.673 (2)	0.6234 (12)	0.030 (4)*
H18A	0.1555 (7)	-0.411 (2)	0.4352 (12)	0.030 (4)*
H18B	0.1925 (8)	-0.601 (2)	0.4416 (12)	0.033 (4)*
H18C	0.1744 (7)	-0.511 (2)	0.5471 (12)	0.032 (4)*
H19A	-0.0437 (8)	-0.018 (2)	0.1586 (13)	0.037 (4)*
H19B	0.0243 (8)	-0.119 (2)	0.1720 (13)	0.038 (4)*
H19C	0.0082 (8)	0.003 (3)	0.2715 (14)	0.042 (4)*
H20A	0.1208 (7)	0.586 (2)	-0.0182 (11)	0.024 (3)*
H20B	0.1018 (7)	0.696 (2)	0.0873 (12)	0.032 (4)*
H20C	0.0475 (7)	0.656 (2)	-0.0181 (11)	0.026 (4)*
H21A	0.3011 (7)	0.515 (2)	0.1612 (12)	0.023 (3)*
H21B	0.3418 (7)	0.545 (2)	0.2761 (11)	0.025 (4)*
H21C	0.2732 (7)	0.626 (2)	0.2555 (11)	0.024 (3)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0170 (3)	0.0133 (3)	0.0192 (3)	-0.0013 (3)	0.0023 (2)	0.0031 (3)
O2	0.0125 (3)	0.0240 (4)	0.0322 (4)	-0.0015 (3)	0.0001 (3)	0.0114 (3)
O3	0.0138 (3)	0.0224 (4)	0.0272 (4)	0.0030 (3)	0.0007 (3)	0.0132 (3)
O4	0.0142 (3)	0.0164 (3)	0.0202 (3)	-0.0017 (3)	-0.0002 (2)	0.0061 (3)
N1	0.0139 (3)	0.0120 (4)	0.0170 (3)	-0.0004 (3)	0.0000 (3)	0.0033 (3)
N2	0.0149 (3)	0.0111 (4)	0.0172 (3)	-0.0007 (3)	0.0004 (3)	0.0030 (3)
N3	0.0153 (3)	0.0137 (4)	0.0137 (3)	0.0033 (3)	0.0020 (3)	0.0020 (3)
C1	0.0173 (4)	0.0173 (5)	0.0189 (4)	0.0029 (4)	0.0038 (3)	0.0031 (4)
C2	0.0151 (4)	0.0213 (5)	0.0281 (5)	0.0019 (4)	0.0027 (4)	0.0041 (4)
C3	0.0190 (4)	0.0159 (4)	0.0243 (5)	0.0007 (4)	-0.0041 (3)	0.0008 (4)
C4	0.0218 (4)	0.0150 (4)	0.0158 (4)	0.0018 (4)	-0.0013 (3)	-0.0005 (3)
C5	0.0173 (4)	0.0151 (4)	0.0166 (4)	0.0019 (3)	0.0020 (3)	-0.0005 (3)
C6	0.0148 (4)	0.0110 (4)	0.0163 (4)	0.0020 (3)	-0.0001 (3)	0.0004 (3)
C7	0.0153 (4)	0.0132 (4)	0.0130 (4)	0.0014 (3)	0.0023 (3)	0.0005 (3)
C8	0.0140 (4)	0.0119 (4)	0.0122 (3)	0.0020 (3)	0.0020 (3)	0.0007 (3)
C9	0.0149 (4)	0.0119 (4)	0.0122 (3)	0.0023 (3)	0.0026 (3)	0.0010 (3)

C10	0.0145 (4)	0.0131 (4)	0.0138 (4)	0.0025 (3)	0.0018 (3)	0.0012 (3)
C11	0.0146 (4)	0.0132 (4)	0.0140 (4)	0.0023 (3)	0.0027 (3)	0.0024 (3)
C12	0.0149 (4)	0.0143 (4)	0.0171 (4)	0.0020 (3)	0.0037 (3)	0.0038 (3)
C13	0.0122 (4)	0.0176 (4)	0.0197 (4)	0.0017 (3)	0.0031 (3)	0.0039 (4)
C14	0.0144 (4)	0.0175 (4)	0.0176 (4)	0.0043 (3)	0.0030 (3)	0.0056 (3)
C15	0.0155 (4)	0.0154 (4)	0.0171 (4)	0.0023 (3)	0.0035 (3)	0.0051 (3)
C16	0.0131 (4)	0.0150 (4)	0.0147 (4)	0.0013 (3)	0.0024 (3)	0.0019 (3)
C17	0.0216 (4)	0.0111 (4)	0.0244 (5)	0.0012 (4)	-0.0003 (4)	0.0034 (4)
C18	0.0166 (4)	0.0155 (4)	0.0196 (4)	-0.0020 (3)	0.0025 (3)	0.0013 (3)
C19	0.0165 (4)	0.0268 (6)	0.0443 (7)	-0.0026 (4)	0.0037 (4)	0.0137 (5)
C20	0.0179 (4)	0.0221 (5)	0.0268 (5)	0.0031 (4)	0.0028 (4)	0.0124 (4)
C21	0.0194 (4)	0.0139 (4)	0.0194 (4)	-0.0017 (4)	0.0036 (3)	0.0029 (3)

Geometric parameters (\AA , $^\circ$)

O1—C9	1.2341 (12)	C8—C9	1.4603 (12)
O2—C13	1.3709 (11)	C10—C11	1.4613 (12)
O2—C19	1.4268 (14)	C10—H10A	0.954 (13)
O3—C14	1.3583 (11)	C11—C16	1.4011 (13)
O3—C20	1.4342 (13)	C11—C12	1.4090 (13)
O4—C16	1.3698 (11)	C12—C13	1.3808 (13)
O4—C21	1.4311 (12)	C12—H12A	0.949 (15)
N1—C9	1.3937 (12)	C13—C14	1.4130 (14)
N1—N2	1.4084 (11)	C14—C15	1.3897 (13)
N1—C6	1.4261 (11)	C15—C16	1.4029 (12)
N2—C7	1.3754 (11)	C15—H15A	0.953 (15)
N2—C17	1.4683 (13)	C17—H17A	1.008 (14)
N3—C10	1.2927 (12)	C17—H17B	0.990 (14)
N3—C8	1.3918 (11)	C17—H17C	0.963 (15)
C1—C6	1.3898 (13)	C18—H18A	1.000 (16)
C1—C2	1.3923 (13)	C18—H18B	0.990 (17)
C1—H1A	0.965 (14)	C18—H18C	0.977 (15)
C2—C3	1.3893 (15)	C19—H19A	0.986 (17)
C2—H2A	0.992 (15)	C19—H19B	1.006 (17)
C3—C4	1.3892 (15)	C19—H19C	0.990 (18)
C3—H3A	0.995 (15)	C20—H20A	0.996 (14)
C4—C5	1.3913 (13)	C20—H20B	1.007 (16)
C4—H4A	0.969 (13)	C20—H20C	0.976 (15)
C5—C6	1.3921 (13)	C21—H21A	0.974 (14)
C5—H5A	0.984 (14)	C21—H21B	0.945 (14)
C7—C8	1.3753 (13)	C21—H21C	0.972 (15)
C7—C18	1.4863 (13)		
C13—O2—C19	116.72 (8)	C13—C12—H12A	122.3 (8)
C14—O3—C20	117.04 (8)	C11—C12—H12A	116.3 (8)
C16—O4—C21	117.65 (7)	O2—C13—C12	125.46 (9)
C9—N1—N2	110.43 (7)	O2—C13—C14	115.26 (8)
C9—N1—C6	125.90 (8)	C12—C13—C14	119.28 (8)
N2—N1—C6	118.94 (7)	O3—C14—C15	124.05 (9)
C7—N2—N1	106.45 (7)	O3—C14—C13	115.62 (8)

supplementary materials

C7—N2—C17	121.21 (8)	C15—C14—C13	120.33 (8)
N1—N2—C17	114.72 (8)	C14—C15—C16	119.75 (9)
C10—N3—C8	119.36 (8)	C14—C15—H15A	119.3 (8)
C6—C1—C2	118.90 (9)	C16—C15—H15A	120.9 (8)
C6—C1—H1A	119.6 (8)	O4—C16—C11	116.75 (8)
C2—C1—H1A	121.5 (8)	O4—C16—C15	122.63 (9)
C3—C2—C1	120.62 (9)	C11—C16—C15	120.61 (8)
C3—C2—H2A	121.2 (8)	N2—C17—H17A	109.0 (8)
C1—C2—H2A	118.2 (8)	N2—C17—H17B	111.4 (9)
C4—C3—C2	119.89 (9)	H17A—C17—H17B	108.9 (12)
C4—C3—H3A	120.6 (9)	N2—C17—H17C	105.3 (9)
C2—C3—H3A	119.4 (9)	H17A—C17—H17C	108.8 (12)
C3—C4—C5	120.18 (9)	H17B—C17—H17C	113.3 (12)
C3—C4—H4A	120.8 (8)	C7—C18—H18A	111.7 (9)
C5—C4—H4A	119.0 (8)	C7—C18—H18B	112.7 (9)
C4—C5—C6	119.34 (9)	H18A—C18—H18B	107.7 (13)
C4—C5—H5A	121.7 (8)	C7—C18—H18C	111.5 (9)
C6—C5—H5A	118.9 (8)	H18A—C18—H18C	106.4 (12)
C1—C6—C5	121.05 (8)	H18B—C18—H18C	106.5 (13)
C1—C6—N1	118.69 (8)	O2—C19—H19A	106.4 (10)
C5—C6—N1	120.26 (8)	O2—C19—H19B	110.6 (9)
C8—C7—N2	110.21 (8)	H19A—C19—H19B	106.9 (14)
C8—C7—C18	128.54 (8)	O2—C19—H19C	110.6 (11)
N2—C7—C18	121.25 (8)	H19A—C19—H19C	114.1 (14)
C7—C8—N3	122.95 (8)	H19B—C19—H19C	108.1 (14)
C7—C8—C9	107.87 (8)	O3—C20—H20A	109.0 (9)
N3—C8—C9	129.17 (8)	O3—C20—H20B	110.2 (9)
O1—C9—N1	124.44 (8)	H20A—C20—H20B	111.3 (12)
O1—C9—C8	131.11 (8)	O3—C20—H20C	104.0 (9)
N1—C9—C8	104.37 (8)	H20A—C20—H20C	111.0 (11)
N3—C10—C11	120.57 (8)	H20B—C20—H20C	111.1 (12)
N3—C10—H10A	121.7 (8)	O4—C21—H21A	109.1 (9)
C11—C10—H10A	117.8 (8)	O4—C21—H21B	105.8 (9)
C16—C11—C12	118.58 (8)	H21A—C21—H21B	110.2 (12)
C16—C11—C10	120.31 (8)	O4—C21—H21C	111.1 (8)
C12—C11—C10	121.11 (8)	H21A—C21—H21C	112.6 (12)
C13—C12—C11	121.40 (9)	H21B—C21—H21C	107.8 (12)
C9—N1—N2—C7	8.54 (10)	C7—C8—C9—O1	-173.56 (9)
C6—N1—N2—C7	165.59 (8)	N3—C8—C9—O1	5.87 (16)
C9—N1—N2—C17	145.49 (8)	C7—C8—C9—N1	3.25 (10)
C6—N1—N2—C17	-57.45 (11)	N3—C8—C9—N1	-177.33 (9)
C6—C1—C2—C3	1.32 (16)	C8—N3—C10—C11	177.82 (8)
C1—C2—C3—C4	-1.34 (17)	N3—C10—C11—C16	176.26 (8)
C2—C3—C4—C5	0.21 (16)	N3—C10—C11—C12	-4.03 (14)
C3—C4—C5—C6	0.91 (16)	C16—C11—C12—C13	1.10 (14)
C2—C1—C6—C5	-0.17 (15)	C10—C11—C12—C13	-178.61 (9)
C2—C1—C6—N1	-179.57 (9)	C19—O2—C13—C12	5.04 (16)
C4—C5—C6—C1	-0.93 (15)	C19—O2—C13—C14	-175.23 (10)
C4—C5—C6—N1	178.45 (9)	C11—C12—C13—O2	-179.44 (9)

C9—N1—C6—C1	−65.96 (13)	C11—C12—C13—C14	0.83 (15)
N2—N1—C6—C1	140.85 (9)	C20—O3—C14—C15	−0.36 (14)
C9—N1—C6—C5	114.64 (11)	C20—O3—C14—C13	178.96 (9)
N2—N1—C6—C5	−38.55 (13)	O2—C13—C14—O3	−0.49 (13)
N1—N2—C7—C8	−6.30 (10)	C12—C13—C14—O3	179.26 (9)
C17—N2—C7—C8	−139.83 (9)	O2—C13—C14—C15	178.86 (9)
N1—N2—C7—C18	173.92 (8)	C12—C13—C14—C15	−1.40 (15)
C17—N2—C7—C18	40.39 (13)	O3—C14—C15—C16	179.28 (9)
N2—C7—C8—N3	−177.54 (8)	C13—C14—C15—C16	0.00 (15)
C18—C7—C8—N3	2.22 (15)	C21—O4—C16—C11	179.81 (8)
N2—C7—C8—C9	1.93 (10)	C21—O4—C16—C15	−1.66 (13)
C18—C7—C8—C9	−178.31 (9)	C12—C11—C16—O4	176.04 (8)
C10—N3—C8—C7	−174.99 (8)	C10—C11—C16—O4	−4.24 (13)
C10—N3—C8—C9	5.66 (14)	C12—C11—C16—C15	−2.52 (14)
N2—N1—C9—O1	169.88 (8)	C10—C11—C16—C15	177.20 (8)
C6—N1—C9—O1	14.78 (14)	C14—C15—C16—O4	−176.49 (9)
N2—N1—C9—C8	−7.20 (9)	C14—C15—C16—C11	1.98 (14)
C6—N1—C9—C8	−162.30 (8)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C10—H10A···O1	0.954 (13)	2.331 (13)	3.0112 (11)	127.8 (10)
C4—H4A···O1 ⁱ	0.969 (13)	2.541 (13)	3.2628 (12)	131.4 (10)
C20—H20A···N3 ⁱⁱ	0.996 (14)	2.577 (14)	3.5383 (13)	162.1 (12)
C20—H20C···O2 ⁱⁱⁱ	0.977 (14)	2.509 (14)	3.4470 (13)	160.8 (12)
C20—H20C···O3 ⁱⁱⁱ	0.977 (14)	2.495 (15)	3.2779 (13)	137.0 (11)

Symmetry codes: (i) $x, -y-1/2, z+1/2$; (ii) $x, -y+1/2, z-1/2$; (iii) $-x, -y+1, -z$.

supplementary materials

Fig. 1




Fig. 2

