

## N'-(4-Cyanobenzylidene)furan-2-carbohydrazide monohydrate

Yu-Feng Li and Fang-Fang Jian\*

Microscale Science Institute, Department of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China

Correspondence e-mail: liyufeng8111@163.com


Received 7 June 2010; accepted 10 June 2010

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.048;  $wR$  factor = 0.135; data-to-parameter ratio = 15.7.

In the title compound,  $\text{C}_{13}\text{H}_9\text{N}_3\text{O}_2\cdot\text{H}_2\text{O}$ , the dihedral angle between the aromatic rings is  $10.7(4)^\circ$  and an intramolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bond occurs. In the crystal, the components are linked by  $\text{N}-\text{H}\cdots\text{O}$ ,  $\text{O}-\text{H}\cdots\text{N}$  and  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds.

### Related literature

For a related structure and background references, see: Li *et al.* (2010).



### Experimental

#### Crystal data

$\text{C}_{13}\text{H}_9\text{N}_3\text{O}_2\cdot\text{H}_2\text{O}$	$V = 1239.3(4)\text{ \AA}^3$
$M_r = 257.25$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 7.0501(14)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$b = 14.295(3)\text{ \AA}$	$T = 293\text{ K}$
$c = 12.640(3)\text{ \AA}$	$0.22 \times 0.20 \times 0.18\text{ mm}$
$\beta = 103.38(3)^\circ$	

#### Data collection

Bruker SMART CCD diffractometer	2834 independent reflections
11389 measured reflections	1568 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.043$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.135$	$\Delta\rho_{\text{max}} = 0.24\text{ e \AA}^{-3}$
$S = 0.98$	$\Delta\rho_{\text{min}} = -0.16\text{ e \AA}^{-3}$
2834 reflections	
180 parameters	

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A $\cdots$ O1	0.86	2.33	2.692 (2)	106
N1—H1A $\cdots$ O3	0.86	2.07	2.920 (2)	169
O3—H3B $\cdots$ N3 <sup>i</sup>	1.00 (4)	1.99 (4)	2.980 (2)	172 (3)
O3—H3C $\cdots$ O2 <sup>ii</sup>	0.88 (3)	1.98 (3)	2.848 (2)	171 (3)

Symmetry codes: (i)  $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$ ; (ii)  $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1997); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5490).

### References

- Bruker (1997). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Li, Y.-F., Zhang, F.-G. & Jian, F.-F. (2010). *Acta Cryst. E* **66**, o1471.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

## **supplementary materials**

*Acta Cryst.* (2010). E66, o1670 [doi:10.1107/S1600536810022221]

## N<sup>1</sup>-(4-Cyanobenzylidene)furan-2-carbohydrazide monohydrate

Y.-F. Li and F.-F. Jian

### Experimental

A mixture of 4-formylbenzonitrile (0.1 mol), and furan-2-carbohydrazide (0.1 mol) was stirred in refluxing ethanol (20 mL) for 2 h to afford the title compound (0.090 mol, yield 90%). Colourless blocks of (I) were obtained by recrystallization from ethanol at room temperature.

### Refinement

H atoms were fixed geometrically and allowed to ride on their attached atoms, with C—H distances=0.97 Å, and with  $U_{\text{iso}}=1.2\text{--}1.5U_{\text{eq}}$ .

### Figures




Fig. 1. The structure of the title compound showing 30% probability displacement ellipsoids.

## N<sup>1</sup>-(4-Cyanobenzylidene)furan-2-carbohydrazide monohydrate

### Crystal data

C <sub>13</sub> H <sub>9</sub> N <sub>3</sub> O <sub>2</sub> ·H <sub>2</sub> O	$F(000) = 536$
$M_r = 257.25$	$D_x = 1.379 \text{ Mg m}^{-3}$
Monoclinic, P2 <sub>1</sub> /n	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 1568 reflections
$a = 7.0501 (14) \text{ \AA}$	$\theta = 2.7\text{--}25.5^\circ$
$b = 14.295 (3) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$c = 12.640 (3) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 103.38 (3)^\circ$	Block, colorless
$V = 1239.3 (4) \text{ \AA}^3$	$0.22 \times 0.20 \times 0.18 \text{ mm}$
$Z = 4$	

### Data collection

Bruker SMART CCD diffractometer	1568 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube graphite	$R_{\text{int}} = 0.043$ $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.3^\circ$

# supplementary materials

---

$\varphi$ and $\omega$ scans	$h = -8 \rightarrow 9$
11389 measured reflections	$k = -18 \rightarrow 18$
2834 independent reflections	$l = -16 \rightarrow 16$

## Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.048$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.135$	H atoms treated by a mixture of independent and constrained refinement
$S = 0.98$	$w = 1/[\sigma^2(F_o^2) + (0.0694P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
2834 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
180 parameters	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.16 \text{ e \AA}^{-3}$

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
N2	0.2572 (2)	0.42473 (10)	0.51829 (12)	0.0431 (4)
N1	0.2142 (2)	0.34390 (9)	0.45997 (12)	0.0440 (4)
H1A	0.1704	0.3458	0.3906	0.053*
O1	0.10154 (18)	0.19128 (9)	0.33670 (10)	0.0526 (4)
C6	0.2239 (3)	0.50009 (12)	0.46228 (15)	0.0461 (5)
H6A	0.1747	0.4964	0.3875	0.055*
C4	0.1844 (3)	0.17816 (13)	0.44486 (15)	0.0442 (5)
O3	0.0883 (2)	0.37589 (11)	0.22607 (12)	0.0654 (5)
C13	0.3865 (3)	0.85907 (14)	0.65549 (17)	0.0548 (5)
O2	0.3075 (2)	0.25365 (9)	0.61179 (11)	0.0595 (4)
C12	0.3138 (3)	0.60216 (13)	0.62661 (16)	0.0525 (5)
H12A	0.3218	0.5497	0.6709	0.063*
C7	0.2620 (3)	0.59186 (12)	0.51417 (15)	0.0420 (4)
C5	0.2414 (3)	0.26093 (12)	0.51272 (15)	0.0436 (4)

C8	0.2453 (3)	0.67132 (13)	0.44979 (16)	0.0518 (5)
H8A	0.2069	0.6655	0.3746	0.062*
C11	0.3534 (3)	0.68908 (13)	0.67294 (16)	0.0551 (5)
H11A	0.3887	0.6952	0.7482	0.066*
C10	0.3405 (3)	0.76786 (12)	0.60708 (16)	0.0453 (5)
C2	0.0589 (3)	0.10477 (13)	0.29261 (18)	0.0566 (6)
H2B	0.0003	0.0931	0.2200	0.068*
N3	0.4234 (3)	0.93164 (12)	0.69148 (17)	0.0721 (6)
C1	0.1133 (3)	0.03943 (15)	0.36854 (18)	0.0654 (6)
H1B	0.1006	-0.0249	0.3588	0.078*
C9	0.2850 (3)	0.75903 (13)	0.49571 (16)	0.0521 (5)
H9A	0.2741	0.8118	0.4516	0.062*
C3	0.1943 (3)	0.08674 (13)	0.46686 (18)	0.0617 (6)
H3A	0.2449	0.0593	0.5342	0.074*
H3B	0.025 (5)	0.438 (3)	0.208 (3)	0.156 (14)*
H3C	-0.005 (4)	0.337 (2)	0.197 (3)	0.120 (12)*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N2	0.0504 (9)	0.0360 (8)	0.0404 (9)	0.0034 (7)	0.0056 (7)	-0.0056 (7)
N1	0.0572 (9)	0.0360 (8)	0.0346 (9)	0.0018 (7)	0.0023 (7)	-0.0016 (6)
O1	0.0708 (9)	0.0435 (7)	0.0389 (8)	-0.0040 (7)	0.0030 (6)	0.0018 (6)
C6	0.0560 (11)	0.0405 (10)	0.0384 (11)	0.0028 (9)	0.0037 (8)	-0.0023 (8)
C4	0.0466 (10)	0.0473 (11)	0.0366 (11)	0.0005 (9)	0.0052 (8)	0.0017 (8)
O3	0.0897 (11)	0.0468 (9)	0.0497 (9)	-0.0047 (9)	-0.0044 (8)	0.0021 (7)
C13	0.0600 (12)	0.0438 (11)	0.0576 (13)	0.0031 (10)	0.0073 (10)	-0.0013 (10)
O2	0.0829 (10)	0.0532 (8)	0.0369 (8)	0.0059 (7)	0.0026 (7)	0.0064 (6)
C12	0.0739 (14)	0.0408 (10)	0.0418 (11)	0.0062 (10)	0.0114 (10)	0.0024 (8)
C7	0.0435 (10)	0.0381 (10)	0.0428 (11)	0.0050 (8)	0.0072 (8)	-0.0002 (8)
C5	0.0451 (10)	0.0468 (11)	0.0377 (11)	0.0064 (9)	0.0074 (8)	0.0064 (8)
C8	0.0653 (12)	0.0467 (11)	0.0392 (12)	0.0028 (9)	0.0038 (9)	-0.0003 (8)
C11	0.0768 (14)	0.0483 (11)	0.0369 (11)	0.0042 (10)	0.0064 (10)	-0.0036 (9)
C10	0.0469 (10)	0.0383 (10)	0.0490 (12)	0.0020 (8)	0.0077 (8)	-0.0048 (8)
C2	0.0708 (14)	0.0460 (11)	0.0488 (12)	-0.0087 (10)	0.0052 (10)	-0.0062 (9)
N3	0.0868 (14)	0.0448 (11)	0.0786 (14)	-0.0037 (10)	0.0064 (11)	-0.0129 (9)
C1	0.0812 (15)	0.0401 (11)	0.0656 (15)	-0.0037 (11)	-0.0021 (12)	-0.0018 (10)
C9	0.0625 (12)	0.0396 (10)	0.0513 (13)	0.0008 (9)	0.0075 (10)	0.0067 (9)
C3	0.0757 (14)	0.0430 (11)	0.0575 (14)	0.0020 (10)	-0.0030 (11)	0.0125 (9)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

N2—C6	1.281 (2)	C12—C11	1.374 (3)
N2—N1	1.3664 (18)	C12—C7	1.391 (3)
N1—C5	1.352 (2)	C12—H12A	0.9300
N1—H1A	0.8600	C7—C8	1.386 (2)
O1—C2	1.361 (2)	C8—C9	1.383 (2)
O1—C4	1.370 (2)	C8—H8A	0.9300
C6—C7	1.464 (2)	C11—C10	1.391 (3)

## supplementary materials

---

C6—H6A	0.9300	C11—H11A	0.9300
C4—C3	1.335 (2)	C10—C9	1.377 (3)
C4—C5	1.462 (2)	C2—C1	1.331 (3)
O3—H3B	1.00 (4)	C2—H2B	0.9300
O3—H3C	0.87 (3)	C1—C3	1.414 (3)
C13—N3	1.139 (2)	C1—H1B	0.9300
C13—C10	1.445 (3)	C9—H9A	0.9300
O2—C5	1.235 (2)	C3—H3A	0.9300
C6—N2—N1	115.07 (15)	C9—C8—C7	121.00 (18)
C5—N1—N2	119.16 (15)	C9—C8—H8A	119.5
C5—N1—H1A	120.4	C7—C8—H8A	119.5
N2—N1—H1A	120.4	C12—C11—C10	119.89 (18)
C2—O1—C4	106.68 (15)	C12—C11—H11A	120.1
N2—C6—C7	121.01 (17)	C10—C11—H11A	120.1
N2—C6—H6A	119.5	C9—C10—C11	120.09 (17)
C7—C6—H6A	119.5	C9—C10—C13	119.90 (17)
C3—C4—O1	109.38 (16)	C11—C10—C13	120.01 (18)
C3—C4—C5	132.55 (18)	C1—C2—O1	110.06 (19)
O1—C4—C5	118.07 (15)	C1—C2—H2B	125.0
H3B—O3—H3C	102 (3)	O1—C2—H2B	125.0
N3—C13—C10	178.5 (2)	C2—C1—C3	106.77 (19)
C11—C12—C7	120.69 (17)	C2—C1—H1B	126.6
C11—C12—H12A	119.7	C3—C1—H1B	126.6
C7—C12—H12A	119.7	C10—C9—C8	119.63 (17)
C8—C7—C12	118.66 (16)	C10—C9—H9A	120.2
C8—C7—C6	119.31 (17)	C8—C9—H9A	120.2
C12—C7—C6	122.03 (16)	C4—C3—C1	107.11 (19)
O2—C5—N1	123.42 (16)	C4—C3—H3A	126.4
O2—C5—C4	120.96 (16)	C1—C3—H3A	126.4
N1—C5—C4	115.60 (16)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
N1—H1A…O1	0.86	2.33	2.692 (2)	106
N1—H1A…O3	0.86	2.07	2.920 (2)	169
O3—H3B…N3 <sup>i</sup>	1.00 (4)	1.99 (4)	2.980 (2)	172 (3)
O3—H3C…O2 <sup>ii</sup>	0.88 (3)	1.98 (3)	2.848 (2)	171 (3)

Symmetry codes: (i)  $x-1/2, -y+3/2, z-1/2$ ; (ii)  $x-1/2, -y+1/2, z-1/2$ .

Fig. 1

