

(2E)-1-(2-Bromophenyl)-3-(3,4,5-trimethoxyphenyl)prop-2-en-1-one

Jerry P. Jasinski,^{a*} Ray J. Butcher,^b K. Veena,^c B. Narayana^d and H. S. Yathirajan^e

^aDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, ^bDepartment of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA, ^cDepartment of Studies in Chemistry, Mangalore University, Mangalagangotri, 574 199, India, ^dDepartment of Studies in Chemistry, Mangalore University, Mangalagangotri 574 199, India, and ^eDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India

Correspondence e-mail: jjasinski@keene.edu


Received 9 June 2010; accepted 10 June 2010

Key indicators: single-crystal X-ray study; $T = 110$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.039; wR factor = 0.112; data-to-parameter ratio = 15.6.

In the chalcone title compound, $\text{C}_{18}\text{H}_{17}\text{BrO}_4$, the dihedral angle between the mean planes of the 2-bromo- and 3,4,5-trimethoxy-substituted benzene rings is $89.3(1)^\circ$. The angles between the mean plane of the prop-2-en-1-one group and the 2-bromophenyl and 3,4,5-trimethoxyphenyl ring planes are $59.7(1)$ and $40.5(8)^\circ$, respectively. While no classical hydrogen bonds are present, three weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ interactions and weak $\text{C}-\text{H}\cdots\text{Br}$ and $\text{C}-\text{H}\cdots\text{Cg}$ π -ring stacking interactions [$\text{C}-\text{H}\cdots\text{Cg}$ distance = $3.377(2)$ Å] are observed, which contribute to the stability of crystal packing.

Related literature

For the radical quenching properties of included phenol groups, see: Dhar (1981). For the anticancer activity of chalcones, see: Dimmock *et al.* (1999). For related structures, see: Chantrapromma *et al.* (2009); Patil *et al.* (2006); Suwunwong *et al.* (2009). For bond distances and angles, see: Allen (2002).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{17}\text{BrO}_4$
 $M_r = 377.23$

Orthorhombic, $Pbca$
 $a = 9.9616(4)$ Å

$b = 13.6020(13)$ Å
 $c = 24.4162(17)$ Å
 $V = 3308.4(4)$ Å³
 $Z = 8$

Mo $K\alpha$ radiation
 $\mu = 2.50$ mm⁻¹
 $T = 110$ K
 $0.47 \times 0.42 \times 0.31$ mm

Data collection

Oxford Diffraction Xcalibur diffractometer with a Ruby (Gemini Cu) detector
Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2007)
 $T_{\min} = 0.499$, $T_{\max} = 1.000$
8122 measured reflections
3296 independent reflections
2940 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.112$
 $S = 1.04$
3296 reflections

211 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.46$ e Å⁻³
 $\Delta\rho_{\min} = -0.67$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg2 is the centroid of the C10–C15 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C6}-\text{H6A}\cdots\text{O1}^i$	0.95	2.44	3.233 (3)	140
$\text{C9}-\text{H9A}\cdots\text{O2}^{ii}$	0.95	2.51	3.308 (3)	141
$\text{C15}-\text{H15A}\cdots\text{O2}^{ii}$	0.95	2.53	3.202 (2)	128
$\text{C17}-\text{H17C}\cdots\text{Br1}^{iii}$	0.98	2.99	3.746 (2)	135
$\text{C17}-\text{H17A}\cdots\text{Cg2}^{iv}$	0.98	2.83	3.379 (2)	125

Symmetry codes: (i) $x - \frac{1}{2}, y, -z + \frac{3}{2}$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, z$; (iii) $-x + \frac{1}{2}, y - \frac{1}{2}, z$; (iv) $x - \frac{1}{2}, -y + \frac{1}{2}, -z + 1$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2007); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

KV thanks UGC for a Junior Research Fellowship and for an SAP chemical grant. HSY thanks UOM for sabbatical leave. RJB acknowledges the NSF MRI program (grant No. CHE-0619278) for funds to purchase the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2317).

References

Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
Chantrapromma, S., Suwunwong, T., Karalai, C. & Fun, H.-K. (2009). *Acta Cryst.* **E65**, o893–o894.
Dhar, D. N. (1981). *The Chemistry of Chalcones and Related Compounds*. New York: John Wiley.
Dimmock, J. R., Elias, D. W., Beazely, M. A. & Kandepu, N. M. (1999). *Curr. Med. Chem.* **6**, 1125–1149.
Oxford Diffraction (2007). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
Patil, P. S., Rosli, M. M., Fun, H.-K., Razak, I. A. & Dharmaprakash, S. M. (2006). *Acta Cryst.* **E62**, o4644–o4645.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Suwunwong, T., Chantrapromma, S. & Fun, H.-K. (2009). *Acta Cryst.* **E65**, o120.

supplementary materials

Acta Cryst. (2010). E66, o1676 [doi:10.1107/S160053681002235X]

(2E)-1-(2-Bromophenyl)-3-(3,4,5-trimethoxyphenyl)prop-2-en-1-one

J. P. Jasinski, R. J. Butcher, K. Veena, B. Narayana and H. S. Yathirajan

Comment

Chalcones, or 1,3-diaryl-2-propen-1-ones, belong to the flavonoid family. Chemically they consist of open-chain flavonoids in which the two aromatic rings are joined by a three-carbon α,β -unsaturated carbonyl system. A vast number of naturally occurring chalcones are polyhydroxylated in the aryl rings. The radical quenching properties of the phenol groups present in many chalcones have raised interest in using the compounds or chalcone rich plant extracts as drugs or food preservatives (Dhar, 1981). Chalcones have been reported to possess many useful biological properties, including anti-inflammatory, antimicrobial, antifungal, antioxidant, cytotoxic, anticancer activities (Dimmock *et al.*, 1999). The crystal structures of some closely related chalcones, *viz.*, (*E*)-1-(4-bromophenyl)-3-(3,4,5-trimethoxyphenyl)prop-2-en-1-one (Suwunwong *et al.*, 2009), (*E*)-1-(4-bromophenyl)-3-(2,4,6-trimethoxyphenyl)prop-2-en-1-one (Chantrapromma *et al.*, 2009) and 1-(4-bromophenyl)-3-(2,4,5-trimethoxyphenyl)prop-2-en-1-one (Patil *et al.*, 2006) have been reported. Hence in continuation with the synthesis and crystal structure determination and also owing to the importance of these flavanoid analogs, this new bromo-trimethoxy substituted chalcone, (I), $C_{18}H_{17}BrO_4$, is synthesized and its crystal structure is reported.

The title compound, (I), $C_{18}H_{17}BrO_4$, is a chalcone with 2-bromophenyl and 3,4,5-trimethoxyphenyl rings bonded at opposite sides of a propene group (Fig. 2). The dihedral angle between mean planes of the benzene rings in the *ortho*-bromo and *meta*-*para*-trimethoxy substituted rings is $89.3(1)^\circ$. The angles between the mean plane of the prop-2-ene-1-one group (C1/C7/O1/C8) and the mean planes of the benzene rings in the 2-bromophenyl (C1–C6) and 3,4,5-trimethoxyphenyl rings (C10–C15) are $59.7(1)^\circ$ and $40.5(8)^\circ$, respectively. Bond distances and angles are in normal ranges (Allen, 2002). While no classical hydrogen bonds are present, three weak intermolecular C—H \cdots O interactions (Fig. 3) and weak C—H \cdots Br (Table 1) and C17—H17A \cdots Cg2 π -ring stacking interactions (H17A \cdots Cg2 = 2.83 Å; H17A–Perp = 2.82 Å; C17—H17A \cdots Cg2 = 125° ; C17 \cdots Cg2—H17A = 3.379(2) Å; Cg2 = C10–C15) are observed which contribute to the stability of crystal packing.

Experimental

A 50% KOH solution was added to a mixture of 2-bromo acetophenone (0.01 mol, 1.99 g) and 3,4,5-trimethoxy benzaldehyde (0.01 mol, 1.96 g) in 25 ml of ethanol (Fig. 1). The mixture was stirred for an hour at room temperature and the precipitate was collected by filtration and purified by recrystallization from ethanol. The single-crystal was grown from ethyl acetate by slow evaporation method and yield of the compound was 45% (m.p. 325–327 K). Analytical data: Found (Calculated) for $C_{18}H_{17}BrO_4$: C %: 57.26 (57.31%); H%: 4.49 (4.54%).

Refinement

The H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C–H distances = 0.95–0.96 Å and with $U_{iso}(H) = 1.18–1.50 U_{eq}(C)$.

Figures




Fig. 1. Reaction Scheme for the title compound.




Fig. 2. Molecular structure of (I), $C_{18}H_{17}BrO_4$, showing the atom labeling scheme and 50% probability displacement ellipsoids.




Fig. 3. Packing diagram of the title compound, $C_{18}H_{17}BrO_4$, viewed down the a axis. Dashed lines indicate weak C—H...O intermolecular hydrogen bond interactions linking the molecules into chains along the (011).

(2E)-1-(2-Bromophenyl)-3-(3,4,5-trimethoxyphenyl)prop-2-en-1-one

Crystal data

$C_{18}H_{17}BrO_4$

$M_r = 377.23$

Orthorhombic, $Pbca$

Hall symbol: -P 2ac 2ab

$a = 9.9616$ (4) Å

$b = 13.6020$ (13) Å

$c = 24.4162$ (17) Å

$V = 3308.4$ (4) Å³

$Z = 8$

$F(000) = 1536$

$D_x = 1.515$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4251 reflections

$\theta = 4.4$ – 74.1°

$\mu = 2.50$ mm⁻¹

$T = 110$ K

Chunk, colorless

$0.47 \times 0.42 \times 0.31$ mm

Data collection

Oxford Diffraction Xcalibur
diffractometer with a Ruby (Gemini Cu) detector

Radiation source: Enhance (Cu) X-ray Source

3296 independent reflections

2940 reflections with $I > 2\sigma(I)$

graphite $R_{\text{int}} = 0.022$
 Detector resolution: 10.5081 pixels mm⁻¹ $\theta_{\text{max}} = 26.3^\circ$, $\theta_{\text{min}} = 2.6^\circ$
 ω scans $h = -12 \rightarrow 7$
 Absorption correction: multi-scan $k = -16 \rightarrow 15$
 (*CrysAlis RED*; Oxford Diffraction, 2007)
 $T_{\text{min}} = 0.499$, $T_{\text{max}} = 1.000$ $l = -30 \rightarrow 28$
 8122 measured reflections

Refinement

Refinement on F^2 Primary atom site location: structure-invariant direct methods
 Least-squares matrix: full Secondary atom site location: difference Fourier map
 $R[F^2 > 2\sigma(F^2)] = 0.039$ Hydrogen site location: inferred from neighbouring sites
 $wR(F^2) = 0.112$ H-atom parameters constrained
 $S = 1.04$ $w = 1/[\sigma^2(F_o^2) + (0.0769P)^2 + 1.9413P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 3296 reflections $(\Delta/\sigma)_{\text{max}} = 0.003$
 211 parameters $\Delta\rho_{\text{max}} = 0.46 \text{ e } \text{\AA}^{-3}$
 0 restraints $\Delta\rho_{\text{min}} = -0.67 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. IR data (KBr) $\nu \text{ cm}^{-1}$: 2998 cm⁻¹, 2937 cm⁻¹, 2839 cm⁻¹ (C—H al. str), 3058 cm⁻¹ (C—H ar.str) 1646 cm⁻¹ (C=O), 1580 cm⁻¹ (C=C); 1245 cm⁻¹ (C—O—C).

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.74430 (2)	0.76760 (2)	0.612337 (10)	0.02659 (13)
O1	0.74217 (17)	0.57654 (14)	0.69518 (10)	0.0340 (5)
O2	0.31902 (16)	0.16171 (11)	0.60595 (6)	0.0191 (3)
O3	0.17918 (15)	0.20383 (11)	0.51830 (6)	0.0180 (3)
O4	0.14169 (16)	0.39311 (11)	0.48455 (6)	0.0203 (3)
C1	0.5557 (2)	0.68349 (15)	0.68770 (8)	0.0159 (4)
C2	0.6035 (2)	0.77077 (15)	0.66443 (8)	0.0169 (4)
C3	0.5453 (2)	0.86108 (16)	0.67668 (9)	0.0219 (4)

supplementary materials

H3A	0.5798	0.9198	0.6610	0.026*
C4	0.4364 (2)	0.86455 (17)	0.71200 (9)	0.0255 (5)
H4A	0.3966	0.9260	0.7208	0.031*
C5	0.3854 (2)	0.77852 (17)	0.73449 (9)	0.0254 (5)
H5A	0.3101	0.7811	0.7584	0.031*
C6	0.4441 (2)	0.68889 (16)	0.72215 (9)	0.0203 (4)
H6A	0.4079	0.6303	0.7373	0.024*
C7	0.6256 (2)	0.58607 (16)	0.68075 (9)	0.0196 (4)
C8	0.5507 (2)	0.50263 (15)	0.65798 (9)	0.0183 (4)
H8A	0.5862	0.4384	0.6630	0.022*
C9	0.4355 (2)	0.51190 (14)	0.63061 (9)	0.0157 (4)
H9A	0.3975	0.5758	0.6282	0.019*
C10	0.3627 (2)	0.43154 (15)	0.60391 (8)	0.0149 (4)
C11	0.3777 (2)	0.33362 (15)	0.62149 (8)	0.0157 (4)
H11A	0.4324	0.3185	0.6522	0.019*
C12	0.3114 (2)	0.25927 (15)	0.59324 (9)	0.0151 (4)
C13	0.2319 (2)	0.28097 (16)	0.54734 (9)	0.0144 (4)
C14	0.2176 (2)	0.37893 (15)	0.53031 (9)	0.0158 (4)
C15	0.2808 (2)	0.45419 (15)	0.55925 (9)	0.0157 (4)
H15A	0.2681	0.5207	0.5486	0.019*
C16	0.3980 (2)	0.13512 (16)	0.65246 (10)	0.0242 (5)
H16A	0.3929	0.0639	0.6581	0.036*
H16B	0.3637	0.1690	0.6850	0.036*
H16C	0.4916	0.1542	0.6462	0.036*
C17	0.0357 (2)	0.19852 (18)	0.51743 (10)	0.0249 (5)
H17A	0.0077	0.1400	0.4970	0.037*
H17B	-0.0005	0.2574	0.4997	0.037*
H17C	0.0018	0.1945	0.5550	0.037*
C18	0.1511 (3)	0.48850 (18)	0.45889 (11)	0.0319 (6)
H18A	0.0980	0.4887	0.4251	0.048*
H18B	0.2452	0.5027	0.4502	0.048*
H18C	0.1165	0.5388	0.4839	0.048*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.02410 (18)	0.0327 (2)	0.02294 (19)	-0.00535 (9)	0.00684 (8)	-0.00220 (9)
O1	0.0244 (9)	0.0223 (9)	0.0552 (13)	0.0010 (6)	-0.0201 (8)	-0.0062 (9)
O2	0.0224 (8)	0.0124 (7)	0.0226 (8)	-0.0033 (6)	-0.0055 (6)	0.0007 (6)
O3	0.0159 (7)	0.0188 (7)	0.0194 (7)	-0.0023 (6)	-0.0009 (6)	-0.0073 (6)
O4	0.0254 (8)	0.0187 (7)	0.0169 (7)	-0.0003 (6)	-0.0078 (6)	0.0006 (6)
C1	0.0177 (9)	0.0161 (10)	0.0138 (9)	-0.0046 (8)	-0.0051 (8)	-0.0020 (7)
C2	0.0177 (10)	0.0213 (11)	0.0117 (9)	-0.0025 (8)	0.0008 (8)	-0.0023 (7)
C3	0.0317 (12)	0.0159 (10)	0.0180 (10)	-0.0011 (9)	-0.0002 (9)	0.0020 (8)
C4	0.0342 (13)	0.0220 (11)	0.0204 (10)	0.0057 (10)	0.0026 (10)	-0.0045 (8)
C5	0.0251 (11)	0.0334 (13)	0.0178 (10)	-0.0009 (10)	0.0052 (9)	-0.0040 (9)
C6	0.0247 (10)	0.0207 (10)	0.0157 (10)	-0.0063 (9)	-0.0027 (8)	0.0007 (8)
C7	0.0208 (10)	0.0183 (10)	0.0197 (10)	-0.0017 (8)	-0.0051 (8)	0.0001 (8)

C8	0.0204 (10)	0.0117 (8)	0.0229 (11)	-0.0009 (8)	-0.0033 (9)	-0.0015 (8)
C9	0.0184 (10)	0.0126 (9)	0.0160 (10)	0.0005 (8)	0.0010 (8)	-0.0011 (7)
C10	0.0142 (9)	0.0141 (9)	0.0164 (9)	-0.0010 (8)	0.0018 (8)	-0.0034 (7)
C11	0.0152 (9)	0.0154 (9)	0.0163 (9)	0.0005 (8)	-0.0024 (8)	-0.0015 (8)
C12	0.0122 (9)	0.0150 (9)	0.0180 (10)	0.0000 (7)	0.0021 (8)	0.0008 (8)
C13	0.0116 (9)	0.0171 (10)	0.0144 (10)	-0.0023 (7)	0.0020 (7)	-0.0038 (8)
C14	0.0134 (8)	0.0194 (10)	0.0145 (9)	0.0019 (8)	0.0010 (8)	-0.0029 (8)
C15	0.0156 (8)	0.0133 (9)	0.0182 (10)	0.0028 (8)	0.0021 (8)	-0.0019 (8)
C16	0.0263 (11)	0.0166 (9)	0.0296 (12)	-0.0004 (9)	-0.0076 (10)	0.0053 (8)
C17	0.0171 (10)	0.0285 (12)	0.0290 (12)	-0.0073 (9)	-0.0057 (9)	0.0013 (9)
C18	0.0448 (15)	0.0260 (12)	0.0251 (12)	-0.0022 (11)	-0.0135 (11)	0.0079 (9)

Geometric parameters (Å, °)

Br1—C2	1.894 (2)	C8—H8A	0.9500
O1—C7	1.221 (3)	C9—C10	1.465 (3)
O2—C12	1.365 (2)	C9—H9A	0.9500
O2—C16	1.428 (3)	C10—C15	1.396 (3)
O3—C13	1.371 (2)	C10—C11	1.407 (3)
O3—C17	1.431 (3)	C11—C12	1.391 (3)
O4—C14	1.363 (3)	C11—H11A	0.9500
O4—C18	1.444 (3)	C12—C13	1.404 (3)
C1—C6	1.396 (3)	C13—C14	1.403 (3)
C1—C2	1.400 (3)	C14—C15	1.394 (3)
C1—C7	1.506 (3)	C15—H15A	0.9500
C2—C3	1.391 (3)	C16—H16A	0.9800
C3—C4	1.386 (3)	C16—H16B	0.9800
C3—H3A	0.9500	C16—H16C	0.9800
C4—C5	1.389 (3)	C17—H17A	0.9800
C4—H4A	0.9500	C17—H17B	0.9800
C5—C6	1.385 (3)	C17—H17C	0.9800
C5—H5A	0.9500	C18—H18A	0.9800
C6—H6A	0.9500	C18—H18B	0.9800
C7—C8	1.468 (3)	C18—H18C	0.9800
C8—C9	1.334 (3)		
C12—O2—C16	117.25 (16)	C12—C11—C10	119.09 (19)
C13—O3—C17	115.38 (17)	C12—C11—H11A	120.5
C14—O4—C18	116.55 (17)	C10—C11—H11A	120.5
C6—C1—C2	118.09 (19)	O2—C12—C11	124.57 (19)
C6—C1—C7	118.82 (18)	O2—C12—C13	114.68 (18)
C2—C1—C7	122.91 (19)	C11—C12—C13	120.74 (19)
C3—C2—C1	121.3 (2)	O3—C13—C14	122.3 (2)
C3—C2—Br1	118.25 (16)	O3—C13—C12	117.91 (19)
C1—C2—Br1	120.35 (15)	C14—C13—C12	119.56 (19)
C4—C3—C2	119.3 (2)	O4—C14—C15	124.20 (19)
C4—C3—H3A	120.3	O4—C14—C13	115.69 (18)
C2—C3—H3A	120.3	C15—C14—C13	120.1 (2)
C3—C4—C5	120.3 (2)	C14—C15—C10	119.83 (19)
C3—C4—H4A	119.9	C14—C15—H15A	120.1

supplementary materials

C5—C4—H4A	119.9	C10—C15—H15A	120.1
C6—C5—C4	120.0 (2)	O2—C16—H16A	109.5
C6—C5—H5A	120.0	O2—C16—H16B	109.5
C4—C5—H5A	120.0	H16A—C16—H16B	109.5
C5—C6—C1	120.9 (2)	O2—C16—H16C	109.5
C5—C6—H6A	119.5	H16A—C16—H16C	109.5
C1—C6—H6A	119.5	H16B—C16—H16C	109.5
O1—C7—C8	120.7 (2)	O3—C17—H17A	109.5
O1—C7—C1	120.0 (2)	O3—C17—H17B	109.5
C8—C7—C1	119.23 (18)	H17A—C17—H17B	109.5
C9—C8—C7	123.64 (19)	O3—C17—H17C	109.5
C9—C8—H8A	118.2	H17A—C17—H17C	109.5
C7—C8—H8A	118.2	H17B—C17—H17C	109.5
C8—C9—C10	125.33 (18)	O4—C18—H18A	109.5
C8—C9—H9A	117.3	O4—C18—H18B	109.5
C10—C9—H9A	117.3	H18A—C18—H18B	109.5
C15—C10—C11	120.61 (19)	O4—C18—H18C	109.5
C15—C10—C9	118.17 (18)	H18A—C18—H18C	109.5
C11—C10—C9	121.18 (19)	H18B—C18—H18C	109.5
C6—C1—C2—C3	2.4 (3)	C9—C10—C11—C12	-177.03 (19)
C7—C1—C2—C3	-172.66 (19)	C16—O2—C12—C11	1.8 (3)
C6—C1—C2—Br1	-174.42 (15)	C16—O2—C12—C13	-179.68 (19)
C7—C1—C2—Br1	10.5 (3)	C10—C11—C12—O2	179.33 (19)
C1—C2—C3—C4	-1.0 (3)	C10—C11—C12—C13	0.9 (3)
Br1—C2—C3—C4	175.89 (17)	C17—O3—C13—C14	-68.0 (3)
C2—C3—C4—C5	-0.6 (3)	C17—O3—C13—C12	117.1 (2)
C3—C4—C5—C6	0.7 (4)	O2—C12—C13—O3	-4.5 (3)
C4—C5—C6—C1	0.8 (3)	C11—C12—C13—O3	174.12 (18)
C2—C1—C6—C5	-2.3 (3)	O2—C12—C13—C14	-179.56 (18)
C7—C1—C6—C5	173.0 (2)	C11—C12—C13—C14	-1.0 (3)
C6—C1—C7—O1	-117.3 (3)	C18—O4—C14—C15	13.2 (3)
C2—C1—C7—O1	57.7 (3)	C18—O4—C14—C13	-165.9 (2)
C6—C1—C7—C8	60.9 (3)	O3—C13—C14—O4	3.4 (3)
C2—C1—C7—C8	-124.0 (2)	C12—C13—C14—O4	178.29 (18)
O1—C7—C8—C9	-164.7 (2)	O3—C13—C14—C15	-175.64 (18)
C1—C7—C8—C9	17.1 (3)	C12—C13—C14—C15	-0.8 (3)
C7—C8—C9—C10	175.4 (2)	O4—C14—C15—C10	-176.39 (19)
C8—C9—C10—C15	-153.2 (2)	C13—C14—C15—C10	2.6 (3)
C8—C9—C10—C11	24.9 (3)	C11—C10—C15—C14	-2.7 (3)
C15—C10—C11—C12	1.0 (3)	C9—C10—C15—C14	175.34 (19)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg2 is the centroid of the C10—C15 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C6—H6A \cdots O1 ⁱ	0.95	2.44	3.233 (3)	140
C9—H9A \cdots O2 ⁱⁱ	0.95	2.51	3.308 (3)	141
C15—H15A \cdots O2 ⁱⁱ	0.95	2.53	3.202 (2)	128

C17—H17C···Br1 ⁱⁱⁱ	0.98	2.99	3.746 (2)	135
C17—H17A···Cg2 ^{iv}	0.98	2.83	3.379 (2)	125

Symmetry codes: (i) $x-1/2, y, -z+3/2$; (ii) $-x+1/2, y+1/2, z$; (iii) $-x+1/2, y-1/2, z$; (iv) $x-1/2, -y+1/2, -z+1$.

Fig. 1




Fig. 2




Fig. 3

