

1-[1-(4-Bromophenyl)ethylidene]-4-(2,4-dimethoxyphenyl)thiosemicarbazide

Muhammad Yaqub,^a Humayun Pervez,^a Nadia Arif,^a M. Nawaz Tahir^{b*} and Mazhar Hussain^a

^aDepartment of Chemistry, Bahauddin Zakariya University, Multan 60800, Pakistan, and

^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 12 June 2010; accepted 12 June 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.041; wR factor = 0.100; data-to-parameter ratio = 19.8.

In the title compound, $\text{C}_{17}\text{H}_{18}\text{BrN}_3\text{O}_2\text{S}$, the dihedral angle between the aromatic rings is $9.15(17)^\circ$. A bifurcated intramolecular $\text{N}-\text{H}\cdots(\text{N},\text{O})$ hydrogen bond generates two $S(5)$ rings and a weak intramolecular $\text{C}-\text{H}\cdots\text{S}$ interaction completes an $S(6)$ ring motif. In the crystal, inversion dimers linked by pairs of $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds generate $R_2^2(8)$ loops and weak $\text{C}-\text{H}\cdots\text{S}$ and $\text{C}-\text{H}\cdots\pi$ interactions are also present.

Related literature

For the pharmacological applications of thiosemicarbazones see: Beraldo & Gambino (2004); Pervez *et al.* (2008, 2010*a,b*). For related structures, see: Jian *et al.* (2005); Martínez *et al.* (2006). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{18}\text{BrN}_3\text{O}_2\text{S}$

$M_r = 408.31$

Monoclinic, $P2_1/n$

$a = 5.8390(2)$ Å

$b = 30.3335(11)$ Å

$c = 9.9423(4)$ Å

$\beta = 94.910(2)^\circ$

$V = 1754.49(11)$ Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 2.48$ mm⁻¹

$T = 296$ K

$0.25 \times 0.22 \times 0.20$ mm

Data collection

Bruker Kappa APEXII CCD

diffractometer

Absorption correction: multi-scan

(*SADABS*; Bruker, 2005)

$T_{\min} = 0.642$, $T_{\max} = 0.652$

17115 measured reflections

4346 independent reflections

2654 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.100$

$S = 1.01$

4346 reflections

220 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.42$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.36$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of C1–C6 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}$	0.86	2.12	2.573 (3)	113
$\text{N1}-\text{H1}\cdots\text{N3}$	0.86	2.05	2.538 (3)	115
$\text{N2}-\text{H2A}\cdots\text{S1}^{\text{i}}$	0.86	2.84	3.662 (2)	161
$\text{C2}-\text{H2}\cdots\text{S1}$	0.93	2.58	3.248 (3)	129
$\text{C17}-\text{H17A}\cdots\text{S1}^{\text{ii}}$	0.96	2.86	3.774 (3)	161
$\text{C8}-\text{H8A}\cdots\text{Cg1}^{\text{iii}}$	0.96	2.98	3.860 (3)	153

Symmetry codes: (i) $-x + 2, -y, -z + 2$; (ii) $x + 1, y, z$; (iii) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

NA gratefully acknowledges the Higher Education Commission (HEC), Islamabad, Pakistan, for providing her with a Scholarship under the Indigenous PhD Program and also for partial funding of this research work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5499).

References

- Beraldo, H. & Gambino, D. (2004). *Mini Rev. Med. Chem.* **4**, 159–165.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Jian, F., Li, Y. & Xiao, H. (2005). *Acta Cryst. E* **61**, o2219–o2220.
- Martínez, J., Adrio, L. A., Antelo, J. M., Ortigueira, J. M., Pereira, M. T., López-Torres, M. & Vila, M. J. (2006). *J. Organomet. Chem.* **691**, 2891–2901.
- Pervez, H., Iqbal, M. S., Tahir, M. Y., Nasim, F. H., Choudhary, M. I. Khan, K. M. & Yaqub, M. (2008). *J. Enz. Inhib. Med. Chem.* **23**, 848–854.
- Pervez, H., Manzoor, N., Yaqub, M., Khan, A., Khan, K. M., Nasim, F. H. & Choudhary, M. I. (2010*a*). *Lett. Drug Des. Discov.* **7**, 102–108.
- Pervez, H., Iqbal, M. S., Saira, N., Yaqub, M. & Tahir, M. N. (2010*b*). *Acta Cryst. E* **66**, o1169–o1170.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o1696 [doi:10.1107/S1600536810022622]

1-[1-(4-Bromophenyl)ethylidene]-4-(2,4-dimethoxyphenyl)thiosemicarbazide

M. Yaqub, H. Pervez, N. Arif, M. N. Tahir and M. Hussain

Comment

Thiosemicarbazones have wide pharmacological properties (Beraldo & Gambino, 2004). Prompted by this, we recently reported the synthesis and medicinal importance of some isatins-thiosemicarbazones (Pervez *et al.*, 2008, 2010*a,b*). Now, we report the synthesis and crystal structure of the title compound (I), (Fig. 1).

The crystal structure of (II) *i.e.* 4-fluoroacetophenone-*N*-propylthiosemicarbazone (Martinez *et al.*, 2006) and (III) *i.e.* 4-phenyl-1-(1-phenylethylidene)thiosemicarbazide (Jian, *et al.*, 2005) have been published. The title compound (I) is different from (II) and (III) due to attachment of substituents at the phenyl rings.

In (I), the phenyl ring A (C1–C6) of 2,4-dimethoxyanilino group, B (C11–C16) of 4-bromophenyl are planar with r. m. s. deviations of 0.0034 and 0.0036 Å, respectively. The thiosemicarbazone moiety C (N1–N3/C9/S1) is also planar with r. m. s. deviation of 0.0062 Å from its mean square plane. The dihedral angle between A/B, A/C and B/C is 9.15 (17), 2.07 (17) and 9.12 (16)°, respectively. Two S(5) ring motifs (Bernstein *et al.*, 1995) (Table 1, Fig. 1) are formed due to strong intramolecular H-bonding of N—H···N and N—H···O types. The weak interaction of C—H···S type completes an S(6) ring motif. The molecules are dimerized due to intermolecular interactions of N—H···S type and complete $R_2^2(8)$ ring motif (Fig. 2). The dimers are interlinked through C—H···S interactions (Table 1). The C—H··· π interaction (Table 1) also play role in stabilizing the molecules.

Experimental

A solution of 4-(2,4-dimethoxyphenyl)thiosemicarbazide (0.15 g, 0.66 mole) in warm ethanol (20 ml) was added drop wise to the stirred solution of 4-bromoacetophenone (0.13 g, 0.66 mol) in warm ethanol (10 ml) containing 2–3 drops of acetic acid. The resultant mixture was then heated under reflux for 30 min. After cooling the reaction mixture to room temperature, the yellow solid was collected by suction filtration, washing with ethanol furnished the title compound in pure form (0.21 g, 95%), m.p. 502 K. Colourless prisms of (I) were grown in chloroform-petroleum ether (1:5) system at room temperature by diffusion method.

Refinement

The H-atoms were positioned geometrically (N—H = 0.86 Å, C—H = 0.93–0.96 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N})$, where $x = 1.5$ for methyl and $x = 1.2$ for all other H-atoms.

Figures




Fig. 1. View of (I) with displacement ellipsoids drawn at the 50% probability level. The dotted lines indicate the intra-molecular H-bonds.




Fig. 2. The partial packing of (I), which shows that molecules form dimers.

1-[1-(4-Bromophenyl)ethylidene]-4-(2,4-dimethoxyphenyl)thiosemicarbazide

Crystal data

$C_{17}H_{18}BrN_3O_2S$

$M_r = 408.31$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 5.8390$ (2) Å

$b = 30.3335$ (11) Å

$c = 9.9423$ (4) Å

$\beta = 94.910$ (2)°

$V = 1754.49$ (11) Å³

$Z = 4$

$F(000) = 832$

$D_x = 1.546$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2654 reflections

$\theta = 2.2$ – 28.3 °

$\mu = 2.48$ mm⁻¹

$T = 296$ K

Prism, colorless

$0.25 \times 0.22 \times 0.20$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer

Radiation source: fine-focus sealed tube graphite

Detector resolution: 7.5 pixels mm⁻¹

ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$T_{\min} = 0.642$, $T_{\max} = 0.652$

17115 measured reflections

4346 independent reflections

2654 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.044$

$\theta_{\max} = 28.3$ °, $\theta_{\min} = 2.2$ °

$h = -4 \rightarrow 7$

$k = -40 \rightarrow 40$

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.041$$

$$wR(F^2) = 0.100$$

$$S = 1.01$$

4346 reflections

220 parameters

0 restraints

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0413P)^2 + 0.4088P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	1.68144 (7)	0.28876 (1)	0.66065 (3)	0.0846 (1)
S1	0.69373 (12)	0.00989 (2)	0.87217 (7)	0.0577 (2)
O1	0.7242 (3)	0.14610 (6)	0.55058 (18)	0.0598 (7)
O2	0.0390 (3)	0.09126 (7)	0.3147 (2)	0.0710 (8)
N1	0.7530 (3)	0.08160 (7)	0.7192 (2)	0.0504 (7)
N2	1.0174 (3)	0.06989 (7)	0.8946 (2)	0.0490 (7)
N3	1.1192 (3)	0.10729 (7)	0.8520 (2)	0.0465 (7)
C1	0.5629 (4)	0.08091 (8)	0.6222 (2)	0.0453 (8)
C2	0.3956 (5)	0.04927 (10)	0.6085 (3)	0.0684 (11)
C3	0.2178 (5)	0.05173 (10)	0.5075 (3)	0.0707 (11)
C4	0.2059 (4)	0.08609 (9)	0.4186 (3)	0.0536 (9)
C5	0.3721 (4)	0.11870 (9)	0.4316 (3)	0.0519 (9)
C6	0.5493 (4)	0.11606 (8)	0.5315 (2)	0.0449 (8)
C7	0.7459 (5)	0.17845 (10)	0.4497 (3)	0.0718 (11)
C8	-0.1278 (5)	0.05715 (11)	0.2945 (3)	0.0714 (11)
C9	0.8214 (4)	0.05526 (8)	0.8225 (2)	0.0426 (8)
C10	1.2991 (4)	0.12299 (8)	0.9180 (2)	0.0413 (8)
C11	1.3922 (4)	0.16334 (8)	0.8596 (2)	0.0397 (7)
C12	1.2659 (4)	0.18592 (9)	0.7571 (3)	0.0522 (9)
C13	1.3493 (5)	0.22333 (9)	0.7001 (3)	0.0585 (10)
C14	1.5632 (4)	0.23844 (9)	0.7440 (3)	0.0511 (9)
C15	1.6935 (5)	0.21732 (9)	0.8453 (3)	0.0560 (10)
C16	1.6074 (4)	0.17997 (9)	0.9025 (3)	0.0508 (9)
C17	1.4082 (5)	0.10312 (9)	1.0456 (3)	0.0565 (9)

supplementary materials

H1	0.84389	0.10350	0.71009	0.0605*
H2	0.40157	0.02564	0.66831	0.0820*
H2A	1.07456	0.05588	0.96481	0.0587*
H3	0.10583	0.02985	0.50022	0.0849*
H5	0.36375	0.14251	0.37241	0.0623*
H7A	0.61321	0.19727	0.44397	0.1075*
H7B	0.75783	0.16426	0.36429	0.1075*
H7C	0.88133	0.19576	0.47252	0.1075*
H8A	-0.21919	0.05609	0.37027	0.1075*
H8B	-0.05148	0.02938	0.28602	0.1075*
H8C	-0.22514	0.06296	0.21373	0.1075*
H12	1.12106	0.17552	0.72613	0.0626*
H13	1.26085	0.23820	0.63226	0.0701*
H15	1.83820	0.22799	0.87536	0.0672*
H16	1.69579	0.16564	0.97145	0.0610*
H17A	1.48971	0.07685	1.02462	0.0848*
H17B	1.29134	0.09589	1.10418	0.0848*
H17C	1.51361	0.12388	1.08973	0.0848*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.1088 (3)	0.0700 (2)	0.0736 (2)	-0.0456 (2)	0.0001 (2)	0.0106 (2)
S1	0.0585 (4)	0.0489 (4)	0.0642 (4)	-0.0114 (3)	-0.0030 (3)	0.0194 (3)
O1	0.0677 (12)	0.0508 (11)	0.0587 (12)	-0.0148 (9)	-0.0080 (9)	0.0172 (9)
O2	0.0668 (12)	0.0697 (14)	0.0712 (13)	0.0015 (11)	-0.0242 (10)	0.0076 (11)
N1	0.0543 (12)	0.0453 (12)	0.0496 (12)	-0.0150 (10)	-0.0074 (10)	0.0128 (10)
N2	0.0522 (12)	0.0458 (12)	0.0470 (12)	-0.0095 (10)	-0.0066 (10)	0.0129 (10)
N3	0.0493 (12)	0.0431 (12)	0.0464 (12)	-0.0070 (9)	-0.0005 (10)	0.0087 (9)
C1	0.0483 (14)	0.0439 (14)	0.0429 (13)	-0.0044 (11)	-0.0008 (11)	0.0061 (11)
C2	0.077 (2)	0.0571 (18)	0.0670 (19)	-0.0215 (15)	-0.0181 (16)	0.0215 (15)
C3	0.0721 (19)	0.063 (2)	0.072 (2)	-0.0253 (16)	-0.0223 (16)	0.0155 (16)
C4	0.0529 (16)	0.0520 (16)	0.0540 (16)	0.0031 (12)	-0.0056 (13)	-0.0032 (13)
C5	0.0644 (17)	0.0443 (14)	0.0462 (15)	0.0062 (12)	-0.0006 (13)	0.0072 (11)
C6	0.0517 (15)	0.0390 (13)	0.0442 (14)	-0.0022 (11)	0.0051 (12)	0.0012 (10)
C7	0.090 (2)	0.0543 (19)	0.070 (2)	-0.0172 (16)	0.0010 (17)	0.0197 (15)
C8	0.0591 (18)	0.076 (2)	0.075 (2)	0.0018 (16)	-0.0180 (15)	-0.0104 (17)
C9	0.0461 (14)	0.0407 (13)	0.0412 (13)	-0.0013 (10)	0.0043 (11)	0.0027 (10)
C10	0.0435 (14)	0.0416 (13)	0.0384 (13)	0.0037 (10)	0.0018 (11)	-0.0006 (10)
C11	0.0417 (13)	0.0388 (13)	0.0384 (12)	-0.0005 (10)	0.0025 (10)	-0.0044 (10)
C12	0.0475 (15)	0.0506 (15)	0.0572 (16)	-0.0094 (12)	-0.0030 (12)	0.0095 (13)
C13	0.0617 (18)	0.0552 (17)	0.0567 (17)	-0.0082 (13)	-0.0053 (14)	0.0109 (13)
C14	0.0603 (16)	0.0469 (15)	0.0465 (15)	-0.0135 (12)	0.0075 (13)	-0.0047 (12)
C15	0.0539 (16)	0.0591 (18)	0.0544 (16)	-0.0181 (13)	0.0004 (13)	-0.0090 (13)
C16	0.0507 (15)	0.0525 (16)	0.0480 (14)	-0.0024 (12)	-0.0029 (12)	0.0005 (12)
C17	0.0650 (17)	0.0505 (16)	0.0515 (16)	-0.0036 (13)	-0.0099 (13)	0.0063 (12)

Geometric parameters (Å, °)

Br1—C14	1.895 (3)	C11—C12	1.387 (4)
S1—C9	1.660 (2)	C12—C13	1.376 (4)
O1—C6	1.370 (3)	C13—C14	1.366 (4)
O1—C7	1.417 (3)	C14—C15	1.368 (4)
O2—C4	1.367 (3)	C15—C16	1.382 (4)
O2—C8	1.423 (4)	C2—H2	0.9300
N1—C1	1.407 (3)	C3—H3	0.9300
N1—C9	1.335 (3)	C5—H5	0.9300
N2—N3	1.365 (3)	C7—H7A	0.9600
N2—C9	1.371 (3)	C7—H7B	0.9600
N3—C10	1.282 (3)	C7—H7C	0.9600
N1—H1	0.8600	C8—H8A	0.9600
N2—H2A	0.8600	C8—H8B	0.9600
C1—C2	1.368 (4)	C8—H8C	0.9600
C1—C6	1.394 (3)	C12—H12	0.9300
C2—C3	1.383 (4)	C13—H13	0.9300
C3—C4	1.365 (4)	C15—H15	0.9300
C4—C5	1.384 (4)	C16—H16	0.9300
C5—C6	1.374 (3)	C17—H17A	0.9600
C10—C11	1.478 (3)	C17—H17B	0.9600
C10—C17	1.497 (4)	C17—H17C	0.9600
C11—C16	1.387 (3)		
C6—O1—C7	118.27 (19)	C14—C15—C16	119.2 (3)
C4—O2—C8	117.3 (2)	C11—C16—C15	121.6 (3)
C1—N1—C9	133.2 (2)	C1—C2—H2	119.00
N3—N2—C9	118.23 (19)	C3—C2—H2	119.00
N2—N3—C10	120.6 (2)	C2—C3—H3	120.00
C9—N1—H1	113.00	C4—C3—H3	120.00
C1—N1—H1	113.00	C4—C5—H5	120.00
N3—N2—H2A	121.00	C6—C5—H5	120.00
C9—N2—H2A	121.00	O1—C7—H7A	109.00
C2—C1—C6	118.1 (2)	O1—C7—H7B	109.00
N1—C1—C6	115.5 (2)	O1—C7—H7C	109.00
N1—C1—C2	126.4 (2)	H7A—C7—H7B	109.00
C1—C2—C3	121.3 (3)	H7A—C7—H7C	109.00
C2—C3—C4	120.3 (3)	H7B—C7—H7C	109.00
O2—C4—C3	124.8 (2)	O2—C8—H8A	109.00
O2—C4—C5	115.9 (2)	O2—C8—H8B	109.00
C3—C4—C5	119.3 (3)	O2—C8—H8C	109.00
C4—C5—C6	120.3 (3)	H8A—C8—H8B	109.00
O1—C6—C5	124.6 (2)	H8A—C8—H8C	109.00
C1—C6—C5	120.7 (2)	H8B—C8—H8C	109.00
O1—C6—C1	114.75 (19)	C11—C12—H12	119.00
S1—C9—N1	127.77 (18)	C13—C12—H12	119.00
S1—C9—N2	119.25 (17)	C12—C13—H13	120.00
N1—C9—N2	112.9 (2)	C14—C13—H13	120.00

supplementary materials

C11—C10—C17	121.5 (2)	C14—C15—H15	120.00
N3—C10—C11	114.80 (19)	C16—C15—H15	120.00
N3—C10—C17	123.7 (2)	C11—C16—H16	119.00
C10—C11—C16	122.3 (2)	C15—C16—H16	119.00
C12—C11—C16	117.2 (2)	C10—C17—H17A	109.00
C10—C11—C12	120.5 (2)	C10—C17—H17B	109.00
C11—C12—C13	121.7 (2)	C10—C17—H17C	109.00
C12—C13—C14	119.4 (3)	H17A—C17—H17B	109.00
Br1—C14—C13	119.2 (2)	H17A—C17—H17C	109.00
C13—C14—C15	120.9 (3)	H17B—C17—H17C	110.00
Br1—C14—C15	119.90 (19)		
C7—O1—C6—C1	169.2 (2)	C2—C3—C4—O2	179.3 (3)
C7—O1—C6—C5	-10.1 (3)	C2—C3—C4—C5	-0.8 (4)
C8—O2—C4—C3	-3.1 (4)	O2—C4—C5—C6	-178.9 (2)
C8—O2—C4—C5	176.9 (2)	C3—C4—C5—C6	1.1 (4)
C9—N1—C1—C2	-2.8 (4)	C4—C5—C6—O1	178.5 (2)
C9—N1—C1—C6	178.0 (2)	C4—C5—C6—C1	-0.8 (4)
C1—N1—C9—S1	0.2 (4)	N3—C10—C11—C12	11.5 (3)
C1—N1—C9—N2	-177.7 (2)	N3—C10—C11—C16	-167.7 (2)
C9—N2—N3—C10	178.1 (2)	C17—C10—C11—C12	-167.6 (2)
N3—N2—C9—S1	-179.91 (16)	C17—C10—C11—C16	13.1 (4)
N3—N2—C9—N1	-1.8 (3)	C10—C11—C12—C13	-179.4 (2)
N2—N3—C10—C11	179.44 (19)	C16—C11—C12—C13	-0.1 (4)
N2—N3—C10—C17	-1.4 (3)	C10—C11—C16—C15	178.9 (2)
N1—C1—C2—C3	-179.0 (3)	C12—C11—C16—C15	-0.3 (4)
C6—C1—C2—C3	0.2 (4)	C11—C12—C13—C14	0.9 (4)
N1—C1—C6—O1	0.1 (3)	C12—C13—C14—Br1	177.8 (2)
N1—C1—C6—C5	179.5 (2)	C12—C13—C14—C15	-1.2 (4)
C2—C1—C6—O1	-179.3 (2)	Br1—C14—C15—C16	-178.2 (2)
C2—C1—C6—C5	0.2 (4)	C13—C14—C15—C16	0.7 (4)
C1—C2—C3—C4	0.1 (4)	C14—C15—C16—C11	0.1 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of C1–C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots O1	0.86	2.12	2.573 (3)	113
N1—H1 \cdots N3	0.86	2.05	2.538 (3)	115
N2—H2A \cdots S1 ⁱ	0.86	2.84	3.662 (2)	161
C2—H2 \cdots S1	0.93	2.58	3.248 (3)	129
C17—H17A \cdots S1 ⁱⁱ	0.96	2.86	3.774 (3)	161
C8—H8A \cdots Cg1 ⁱⁱⁱ	0.96	2.98	3.860 (3)	153

Symmetry codes: (i) $-x+2, -y, -z+2$; (ii) $x+1, y, z$; (iii) $x-1, y, z$.

Fig. 1




Fig. 2

