

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***trans*-3-(3,4-Dimethoxyphenyl)-2-(4-nitrophenyl)prop-2-enitrile**Abdullah M. Asiri,^a Salman A. Khan,^a Kong Wai Tan^b and Seik Weng Ng^{b*}^aChemistry Department, Faculty of Science, King Abdul Aziz University, PO Box 80203, Jeddah 21589, Saudi Arabia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 15 June 2010; accepted 16 June 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.047; wR factor = 0.133; data-to-parameter ratio = 16.0.

The asymmetric unit of the title compound, $\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_4$, contains two independent molecules in which the benzene rings are in a *trans* arrangement with respect to the $\text{C}=\text{C}$ double bond and the rings are inclined by 4.3 (1) and 22.1 (1)° with respect to each other.

Related literature

For the crystal structure of α -((4-methoxyphenyl)methylene)-4-nitrobenzeneacetonitrile, see: Vrcelj *et al.* (2002). For background literature on this class of pigments, see: Asiri (1999).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_4$	$\gamma = 100.156$ (1)°
$M_r = 310.30$	$V = 1453.3$ (2) Å ³
Triclinic, $P\bar{1}$	$Z = 4$
$a = 10.2211$ (8) Å	Mo $K\alpha$ radiation
$b = 11.9460$ (9) Å	$\mu = 0.10$ mm ⁻¹
$c = 12.2764$ (10) Å	$T = 100$ K
$\alpha = 91.094$ (1)°	$0.40 \times 0.20 \times 0.10$ mm
$\beta = 99.542$ (1)°	

Data collection

Bruker SMART APEX diffractometer	6628 independent reflections
13853 measured reflections	4851 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$	415 parameters
$wR(F^2) = 0.133$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.40$ e Å ⁻³
6628 reflections	$\Delta\rho_{\text{min}} = -0.24$ e Å ⁻³

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdul Aziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5073).

References

- Asiri, A. M. (1999). *Dyes Pigments*, **42**, 209–213.
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Vrcelj, R. M., Shepherd, E. E. A., Yoon, C.-S., Sherwood, J. N. & Kennedy, A. R. (2002). *Cryst. Growth Des.* **2**, 606–617.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.

supplementary materials

Acta Cryst. (2010). E66, o1733 [doi:10.1107/S1600536810023196]

***trans*-3-(3,4-Dimethoxyphenyl)-2-(4-nitrophenyl)prop-2-enitrile**

A. M. Asiri, S. A. Khan, K. W. Tan and S. W. Ng

Comment

Organic photochromic compounds having donor and acceptor parts that are conjugated are potential optical materials. Such compounds, as exemplified by the title compound, are synthesized from carbonyl compounds having an active methylene group by using the Knoevenagel condensation (Asiri, 1999). The title compound (Fig. 1) features a double-bond with two aromatic substituents in *trans*-positions.

α -((4-Methoxyphenyl)methylene)-4-nitrobenzeneacetonitrile, a stilbene derivative, exists in a *cis* and a *trans* form; interestingly, the *trans* form crystallizes in three modifications (Vrcelj *et al.*, 2002).

Experimental

3,4-Dimethoxybenzaldehyde (0.41 g, 2.5 mmol) and 4-nitrobenzyl cyanide (0.40 g, 2.5 mmol) were heated in ethanol (15 ml) for 3 h; several drops of pyridine were added. The reaction was monitored by TLC. The solution was cooled and the residue recrystallized from a methanol-chloroform (1/1) mixture.

Refinement

H-atoms were placed in calculated positions [C–H 0.95 to 0.98 Å, $U(\text{H})$ 1.2 to 1.5 $U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_4$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

***trans*-3-(3,4-Dimethoxyphenyl)-2-(4-nitrophenyl)prop-2-enitrile**

Crystal data

$\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_4$

$M_r = 310.30$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 10.2211$ (8) Å

$b = 11.9460$ (9) Å

$Z = 4$

$F(000) = 648$

$D_x = 1.418$ Mg m $^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4130 reflections

$\theta = 2.4$ – 28.1°

supplementary materials

$c = 12.2764 (10) \text{ \AA}$
 $\alpha = 91.094 (1)^\circ$
 $\beta = 99.542 (1)^\circ$
 $\gamma = 100.156 (1)^\circ$
 $V = 1453.3 (2) \text{ \AA}^3$

$\mu = 0.10 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
Prism, orange
 $0.40 \times 0.20 \times 0.10 \text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
Radiation source: fine-focus sealed tube
graphite
 ω scans
13853 measured reflections
6628 independent reflections

4851 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$
 $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 1.7^\circ$
 $h = -13 \rightarrow 13$
 $k = -13 \rightarrow 15$
 $l = -15 \rightarrow 15$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.133$
 $S = 1.02$
6628 reflections
415 parameters
0 restraints

Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0681P)^2 + 0.1844P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.40 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.83856 (13)	0.82806 (10)	0.70740 (9)	0.0232 (3)
O2	0.77000 (13)	0.93616 (10)	0.53428 (9)	0.0240 (3)
O3	1.55559 (14)	0.19841 (12)	0.52637 (11)	0.0352 (3)
O4	1.53950 (14)	0.26192 (12)	0.36149 (10)	0.0320 (3)
O5	0.70198 (12)	0.97785 (10)	0.86081 (9)	0.0229 (3)
O6	0.86075 (12)	0.93425 (10)	1.03307 (9)	0.0217 (3)
O7	0.08469 (14)	1.67994 (12)	1.06765 (12)	0.0343 (3)
O8	0.21823 (14)	1.70388 (11)	1.22686 (10)	0.0331 (3)

N1	1.05331 (17)	0.49336 (14)	0.73956 (12)	0.0312 (4)
N2	1.51106 (15)	0.25875 (13)	0.45509 (12)	0.0241 (3)
N3	0.34748 (15)	1.21226 (13)	0.81431 (11)	0.0231 (3)
N4	0.17862 (16)	1.65717 (13)	1.13420 (12)	0.0250 (3)
C1	1.04870 (18)	0.72332 (15)	0.52312 (13)	0.0206 (4)
C2	0.99041 (17)	0.73654 (14)	0.61828 (13)	0.0199 (4)
H2	1.0132	0.6944	0.6813	0.024*
C3	0.90092 (17)	0.80994 (14)	0.62037 (12)	0.0186 (4)
C4	0.86490 (17)	0.87172 (14)	0.52605 (13)	0.0193 (4)
C5	0.92618 (18)	0.86239 (15)	0.43441 (13)	0.0207 (4)
H5	0.9058	0.9065	0.3723	0.025*
C6	1.01697 (18)	0.78904 (15)	0.43308 (13)	0.0214 (4)
H6	1.0583	0.7834	0.3698	0.026*
C7	0.86525 (19)	0.76349 (16)	0.80335 (13)	0.0243 (4)
H7A	0.8146	0.7843	0.8594	0.036*
H7B	0.9620	0.7800	0.8334	0.036*
H7C	0.8374	0.6820	0.7828	0.036*
C8	0.72379 (19)	0.99399 (16)	0.43758 (13)	0.0259 (4)
H8A	0.6562	1.0379	0.4537	0.039*
H8B	0.6834	0.9379	0.3766	0.039*
H8C	0.8003	1.0455	0.4165	0.039*
C9	1.13862 (18)	0.64400 (15)	0.51114 (13)	0.0222 (4)
H9	1.1877	0.6586	0.4521	0.027*
C10	1.16493 (18)	0.55272 (15)	0.56936 (13)	0.0211 (4)
C11	1.10295 (18)	0.52212 (15)	0.66500 (13)	0.0215 (4)
C12	1.25355 (17)	0.47620 (14)	0.53921 (13)	0.0188 (4)
C13	1.29928 (17)	0.39916 (14)	0.61403 (13)	0.0198 (4)
H13	1.2727	0.3960	0.6846	0.024*
C14	1.38285 (17)	0.32729 (15)	0.58691 (13)	0.0204 (4)
H14	1.4135	0.2749	0.6380	0.024*
C15	1.42069 (17)	0.33351 (14)	0.48391 (13)	0.0199 (4)
C16	1.37606 (17)	0.40682 (15)	0.40728 (13)	0.0214 (4)
H16	1.4025	0.4085	0.3366	0.026*
C17	1.29240 (17)	0.47784 (15)	0.43440 (13)	0.0205 (4)
H17	1.2606	0.5284	0.3818	0.025*
C20	0.61948 (17)	1.17523 (14)	1.06498 (13)	0.0185 (3)
C21	0.61639 (17)	1.11728 (15)	0.96284 (13)	0.0198 (4)
H21	0.5576	1.1331	0.8989	0.024*
C22	0.69803 (17)	1.03785 (14)	0.95513 (12)	0.0187 (4)
C23	0.78556 (17)	1.01356 (14)	1.04955 (13)	0.0186 (4)
C24	0.78929 (17)	1.07024 (15)	1.15006 (13)	0.0205 (4)
H24	0.8479	1.0541	1.2140	0.025*
C25	0.70765 (17)	1.15032 (15)	1.15751 (13)	0.0204 (4)
H25	0.7116	1.1890	1.2267	0.024*
C26	0.62349 (19)	1.00501 (16)	0.76028 (13)	0.0241 (4)
H26A	0.6351	0.9559	0.6990	0.036*
H26B	0.6537	1.0849	0.7454	0.036*
H26C	0.5281	0.9929	0.7676	0.036*
C27	0.95890 (18)	0.91253 (16)	1.12473 (13)	0.0236 (4)

supplementary materials

H27A	1.0062	0.8538	1.1021	0.035*
H27B	0.9133	0.8863	1.1863	0.035*
H27C	1.0241	0.9827	1.1483	0.035*
C28	0.53948 (17)	1.26103 (15)	1.08144 (13)	0.0200 (4)
H28	0.5597	1.2965	1.1537	0.024*
C29	0.44121 (17)	1.29992 (14)	1.01262 (12)	0.0179 (3)
C30	0.39106 (17)	1.24995 (14)	0.90229 (13)	0.0179 (3)
C31	0.37513 (17)	1.39410 (14)	1.04263 (12)	0.0177 (3)
C32	0.25478 (18)	1.41207 (15)	0.97922 (13)	0.0209 (4)
H32	0.2156	1.3634	0.9158	0.025*
C33	0.19096 (18)	1.49957 (15)	1.00682 (14)	0.0229 (4)
H33	0.1105	1.5126	0.9619	0.028*
C34	0.24767 (17)	1.56705 (14)	1.10128 (13)	0.0204 (4)
C35	0.36790 (18)	1.55373 (14)	1.16528 (13)	0.0199 (4)
H35	0.4058	1.6023	1.2290	0.024*
C36	0.43196 (18)	1.46850 (14)	1.13491 (13)	0.0194 (4)
H36	0.5160	1.4600	1.1772	0.023*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0321 (7)	0.0269 (7)	0.0152 (5)	0.0114 (6)	0.0105 (5)	0.0017 (5)
O2	0.0294 (7)	0.0285 (7)	0.0184 (6)	0.0152 (6)	0.0064 (5)	0.0021 (5)
O3	0.0390 (9)	0.0396 (8)	0.0343 (7)	0.0228 (7)	0.0102 (6)	0.0074 (6)
O4	0.0344 (8)	0.0400 (8)	0.0257 (6)	0.0113 (6)	0.0132 (6)	-0.0049 (6)
O5	0.0274 (7)	0.0286 (7)	0.0150 (5)	0.0121 (6)	0.0034 (5)	-0.0039 (5)
O6	0.0244 (7)	0.0258 (7)	0.0184 (6)	0.0130 (5)	0.0052 (5)	0.0013 (5)
O7	0.0304 (8)	0.0300 (8)	0.0450 (8)	0.0147 (6)	0.0043 (6)	-0.0018 (6)
O8	0.0397 (8)	0.0300 (8)	0.0324 (7)	0.0083 (6)	0.0131 (6)	-0.0085 (6)
N1	0.0406 (10)	0.0357 (10)	0.0234 (7)	0.0156 (8)	0.0132 (7)	0.0058 (7)
N2	0.0222 (8)	0.0264 (8)	0.0246 (7)	0.0052 (7)	0.0062 (6)	-0.0032 (6)
N3	0.0266 (8)	0.0261 (8)	0.0179 (7)	0.0080 (7)	0.0043 (6)	0.0004 (6)
N4	0.0249 (8)	0.0212 (8)	0.0317 (8)	0.0057 (7)	0.0120 (7)	-0.0003 (6)
C1	0.0233 (9)	0.0210 (9)	0.0181 (8)	0.0041 (7)	0.0056 (7)	-0.0005 (6)
C2	0.0232 (9)	0.0209 (9)	0.0161 (7)	0.0041 (7)	0.0045 (7)	0.0015 (6)
C3	0.0220 (9)	0.0196 (9)	0.0145 (7)	0.0016 (7)	0.0064 (6)	-0.0025 (6)
C4	0.0210 (9)	0.0189 (9)	0.0181 (8)	0.0043 (7)	0.0038 (6)	-0.0025 (6)
C5	0.0255 (10)	0.0225 (9)	0.0148 (7)	0.0059 (7)	0.0036 (7)	0.0022 (6)
C6	0.0246 (9)	0.0252 (9)	0.0163 (7)	0.0052 (7)	0.0079 (7)	0.0002 (6)
C7	0.0322 (10)	0.0275 (10)	0.0154 (7)	0.0067 (8)	0.0092 (7)	0.0018 (7)
C8	0.0324 (11)	0.0286 (10)	0.0197 (8)	0.0152 (8)	0.0027 (7)	0.0025 (7)
C9	0.0206 (9)	0.0270 (10)	0.0191 (8)	0.0023 (7)	0.0057 (7)	-0.0010 (7)
C10	0.0214 (9)	0.0264 (9)	0.0167 (8)	0.0063 (7)	0.0047 (7)	-0.0001 (7)
C11	0.0247 (10)	0.0245 (9)	0.0173 (8)	0.0100 (8)	0.0034 (7)	-0.0001 (7)
C12	0.0186 (9)	0.0209 (9)	0.0163 (7)	0.0020 (7)	0.0037 (6)	-0.0009 (6)
C13	0.0220 (9)	0.0231 (9)	0.0144 (7)	0.0032 (7)	0.0047 (6)	-0.0002 (6)
C14	0.0211 (9)	0.0208 (9)	0.0188 (8)	0.0032 (7)	0.0030 (7)	0.0013 (6)
C15	0.0172 (9)	0.0214 (9)	0.0209 (8)	0.0032 (7)	0.0038 (6)	-0.0048 (7)

C16	0.0206 (9)	0.0285 (10)	0.0151 (7)	0.0027 (7)	0.0054 (6)	-0.0010 (7)
C17	0.0239 (9)	0.0236 (9)	0.0144 (7)	0.0037 (7)	0.0045 (7)	0.0017 (6)
C20	0.0190 (9)	0.0212 (9)	0.0173 (7)	0.0060 (7)	0.0061 (6)	0.0022 (6)
C21	0.0206 (9)	0.0242 (9)	0.0155 (7)	0.0060 (7)	0.0038 (6)	0.0010 (6)
C22	0.0210 (9)	0.0214 (9)	0.0147 (7)	0.0034 (7)	0.0063 (6)	-0.0006 (6)
C23	0.0196 (9)	0.0196 (9)	0.0184 (8)	0.0053 (7)	0.0068 (7)	0.0020 (6)
C24	0.0200 (9)	0.0259 (9)	0.0172 (8)	0.0071 (7)	0.0044 (6)	0.0030 (7)
C25	0.0248 (9)	0.0235 (9)	0.0143 (7)	0.0065 (7)	0.0055 (7)	-0.0012 (6)
C26	0.0296 (10)	0.0298 (10)	0.0144 (7)	0.0101 (8)	0.0030 (7)	-0.0016 (7)
C27	0.0229 (10)	0.0286 (10)	0.0219 (8)	0.0119 (8)	0.0036 (7)	0.0027 (7)
C28	0.0241 (9)	0.0231 (9)	0.0148 (7)	0.0063 (7)	0.0069 (7)	0.0003 (6)
C29	0.0208 (9)	0.0210 (9)	0.0140 (7)	0.0054 (7)	0.0075 (6)	0.0018 (6)
C30	0.0203 (9)	0.0185 (9)	0.0179 (8)	0.0075 (7)	0.0071 (7)	0.0041 (6)
C31	0.0200 (9)	0.0210 (9)	0.0141 (7)	0.0048 (7)	0.0073 (6)	0.0047 (6)
C32	0.0239 (9)	0.0243 (9)	0.0157 (7)	0.0075 (7)	0.0036 (7)	-0.0007 (6)
C33	0.0221 (9)	0.0278 (10)	0.0211 (8)	0.0089 (8)	0.0051 (7)	0.0037 (7)
C34	0.0226 (9)	0.0185 (9)	0.0237 (8)	0.0064 (7)	0.0115 (7)	0.0025 (7)
C35	0.0244 (9)	0.0202 (9)	0.0163 (7)	0.0031 (7)	0.0079 (7)	0.0013 (6)
C36	0.0203 (9)	0.0234 (9)	0.0158 (7)	0.0053 (7)	0.0048 (6)	0.0031 (6)

Geometric parameters (Å, °)

O1—C3	1.3639 (18)	C13—C14	1.385 (2)
O1—C7	1.435 (2)	C13—H13	0.9500
O2—C4	1.355 (2)	C14—C15	1.382 (2)
O2—C8	1.439 (2)	C14—H14	0.9500
O3—N2	1.2248 (19)	C15—C16	1.375 (2)
O4—N2	1.2308 (18)	C16—C17	1.379 (2)
O5—C22	1.3602 (18)	C16—H16	0.9500
O5—C26	1.432 (2)	C17—H17	0.9500
O6—C23	1.3528 (19)	C20—C25	1.400 (2)
O6—C27	1.438 (2)	C20—C21	1.414 (2)
O7—N4	1.226 (2)	C20—C28	1.450 (2)
O8—N4	1.2299 (19)	C21—C22	1.381 (2)
N1—C11	1.148 (2)	C21—H21	0.9500
N2—C15	1.471 (2)	C22—C23	1.410 (2)
N3—C30	1.148 (2)	C23—C24	1.388 (2)
N4—C34	1.472 (2)	C24—C25	1.387 (2)
C1—C6	1.396 (2)	C24—H24	0.9500
C1—C2	1.415 (2)	C25—H25	0.9500
C1—C9	1.453 (2)	C26—H26A	0.9800
C2—C3	1.377 (2)	C26—H26B	0.9800
C2—H2	0.9500	C26—H26C	0.9800
C3—C4	1.417 (2)	C27—H27A	0.9800
C4—C5	1.387 (2)	C27—H27B	0.9800
C5—C6	1.386 (2)	C27—H27C	0.9800
C5—H5	0.9500	C28—C29	1.355 (2)
C6—H6	0.9500	C28—H28	0.9500
C7—H7A	0.9800	C29—C30	1.444 (2)

supplementary materials

C7—H7B	0.9800	C29—C31	1.481 (2)
C7—H7C	0.9800	C31—C32	1.395 (2)
C8—H8A	0.9800	C31—C36	1.404 (2)
C8—H8B	0.9800	C32—C33	1.390 (2)
C8—H8C	0.9800	C32—H32	0.9500
C9—C10	1.357 (2)	C33—C34	1.381 (2)
C9—H9	0.9500	C33—H33	0.9500
C10—C11	1.447 (2)	C34—C35	1.381 (2)
C10—C12	1.478 (2)	C35—C36	1.381 (2)
C12—C13	1.397 (2)	C35—H35	0.9500
C12—C17	1.408 (2)	C36—H36	0.9500
C3—O1—C7	117.62 (13)	C15—C16—H16	120.5
C4—O2—C8	117.01 (12)	C17—C16—H16	120.5
C22—O5—C26	117.68 (13)	C16—C17—C12	120.72 (16)
C23—O6—C27	117.60 (12)	C16—C17—H17	119.6
O3—N2—O4	123.80 (15)	C12—C17—H17	119.6
O3—N2—C15	118.02 (13)	C25—C20—C21	118.28 (15)
O4—N2—C15	118.16 (15)	C25—C20—C28	116.82 (14)
O7—N4—O8	124.42 (15)	C21—C20—C28	124.90 (15)
O7—N4—C34	117.73 (14)	C22—C21—C20	120.62 (15)
O8—N4—C34	117.85 (15)	C22—C21—H21	119.7
C6—C1—C2	118.54 (15)	C20—C21—H21	119.7
C6—C1—C9	116.93 (14)	O5—C22—C21	124.87 (15)
C2—C1—C9	124.53 (15)	O5—C22—C23	114.96 (14)
C3—C2—C1	120.58 (15)	C21—C22—C23	120.17 (14)
C3—C2—H2	119.7	O6—C23—C24	125.05 (15)
C1—C2—H2	119.7	O6—C23—C22	115.42 (13)
O1—C3—C2	125.12 (15)	C24—C23—C22	119.53 (15)
O1—C3—C4	114.88 (14)	C25—C24—C23	120.21 (15)
C2—C3—C4	119.99 (14)	C25—C24—H24	119.9
O2—C4—C5	125.28 (15)	C23—C24—H24	119.9
O2—C4—C3	115.28 (13)	C24—C25—C20	121.20 (14)
C5—C4—C3	119.44 (15)	C24—C25—H25	119.4
C6—C5—C4	120.24 (15)	C20—C25—H25	119.4
C6—C5—H5	119.9	O5—C26—H26A	109.5
C4—C5—H5	119.9	O5—C26—H26B	109.5
C5—C6—C1	121.08 (14)	H26A—C26—H26B	109.5
C5—C6—H6	119.5	O5—C26—H26C	109.5
C1—C6—H6	119.5	H26A—C26—H26C	109.5
O1—C7—H7A	109.5	H26B—C26—H26C	109.5
O1—C7—H7B	109.5	O6—C27—H27A	109.5
H7A—C7—H7B	109.5	O6—C27—H27B	109.5
O1—C7—H7C	109.5	H27A—C27—H27B	109.5
H7A—C7—H7C	109.5	O6—C27—H27C	109.5
H7B—C7—H7C	109.5	H27A—C27—H27C	109.5
O2—C8—H8A	109.5	H27B—C27—H27C	109.5
O2—C8—H8B	109.5	C29—C28—C20	131.78 (15)
H8A—C8—H8B	109.5	C29—C28—H28	114.1
O2—C8—H8C	109.5	C20—C28—H28	114.1

H8A—C8—H8C	109.5	C28—C29—C30	121.52 (15)
H8B—C8—H8C	109.5	C28—C29—C31	124.14 (14)
C10—C9—C1	131.43 (15)	C30—C29—C31	114.33 (14)
C10—C9—H9	114.3	N3—C30—C29	177.59 (18)
C1—C9—H9	114.3	C32—C31—C36	118.01 (15)
C9—C10—C11	121.11 (15)	C32—C31—C29	120.82 (15)
C9—C10—C12	123.52 (15)	C36—C31—C29	121.17 (15)
C11—C10—C12	115.34 (15)	C33—C32—C31	121.49 (16)
N1—C11—C10	177.26 (19)	C33—C32—H32	119.3
C13—C12—C17	118.40 (15)	C31—C32—H32	119.3
C13—C12—C10	120.50 (14)	C34—C33—C32	118.21 (16)
C17—C12—C10	121.09 (15)	C34—C33—H33	120.9
C14—C13—C12	121.04 (14)	C32—C33—H33	120.9
C14—C13—H13	119.5	C35—C34—C33	122.21 (15)
C12—C13—H13	119.5	C35—C34—N4	118.82 (15)
C15—C14—C13	118.55 (16)	C33—C34—N4	118.97 (16)
C15—C14—H14	120.7	C34—C35—C36	118.78 (15)
C13—C14—H14	120.7	C34—C35—H35	120.6
C16—C15—C14	122.18 (16)	C36—C35—H35	120.6
C16—C15—N2	118.92 (14)	C35—C36—C31	121.19 (16)
C14—C15—N2	118.90 (15)	C35—C36—H36	119.4
C15—C16—C17	119.09 (14)	C31—C36—H36	119.4
C6—C1—C2—C3	2.1 (3)	C25—C20—C21—C22	0.3 (3)
C9—C1—C2—C3	-176.79 (16)	C28—C20—C21—C22	179.07 (16)
C7—O1—C3—C2	-1.8 (2)	C26—O5—C22—C21	5.0 (2)
C7—O1—C3—C4	176.78 (15)	C26—O5—C22—C23	-175.15 (15)
C1—C2—C3—O1	179.45 (16)	C20—C21—C22—O5	179.90 (16)
C1—C2—C3—C4	1.0 (3)	C20—C21—C22—C23	0.1 (3)
C8—O2—C4—C5	3.9 (2)	C27—O6—C23—C24	-4.6 (2)
C8—O2—C4—C3	-175.80 (15)	C27—O6—C23—C22	175.45 (15)
O1—C3—C4—O2	-2.5 (2)	O5—C22—C23—O6	0.0 (2)
C2—C3—C4—O2	176.11 (15)	C21—C22—C23—O6	179.80 (15)
O1—C3—C4—C5	177.76 (15)	O5—C22—C23—C24	179.98 (15)
C2—C3—C4—C5	-3.6 (3)	C21—C22—C23—C24	-0.2 (3)
O2—C4—C5—C6	-176.56 (16)	O6—C23—C24—C25	179.92 (16)
C3—C4—C5—C6	3.1 (3)	C22—C23—C24—C25	-0.1 (3)
C4—C5—C6—C1	0.0 (3)	C23—C24—C25—C20	0.5 (3)
C2—C1—C6—C5	-2.6 (3)	C21—C20—C25—C24	-0.6 (3)
C9—C1—C6—C5	176.37 (16)	C28—C20—C25—C24	-179.45 (16)
C6—C1—C9—C10	-163.11 (19)	C25—C20—C28—C29	-175.78 (18)
C2—C1—C9—C10	15.8 (3)	C21—C20—C28—C29	5.4 (3)
C1—C9—C10—C11	-2.8 (3)	C20—C28—C29—C30	4.5 (3)
C1—C9—C10—C12	175.28 (17)	C20—C28—C29—C31	-176.36 (17)
C9—C10—C12—C13	166.39 (17)	C28—C29—C31—C32	-164.68 (16)
C11—C10—C12—C13	-15.4 (2)	C30—C29—C31—C32	14.5 (2)
C9—C10—C12—C17	-14.5 (3)	C28—C29—C31—C36	15.9 (3)
C11—C10—C12—C17	163.69 (16)	C30—C29—C31—C36	-164.91 (15)
C17—C12—C13—C14	1.1 (3)	C36—C31—C32—C33	-1.0 (2)
C10—C12—C13—C14	-179.74 (16)	C29—C31—C32—C33	179.49 (15)

supplementary materials

C12—C13—C14—C15	0.2 (3)	C31—C32—C33—C34	-1.9 (3)
C13—C14—C15—C16	-1.3 (3)	C32—C33—C34—C35	3.2 (3)
C13—C14—C15—N2	179.07 (15)	C32—C33—C34—N4	-177.59 (15)
O3—N2—C15—C16	176.16 (16)	O7—N4—C34—C35	168.39 (16)
O4—N2—C15—C16	-2.5 (2)	O8—N4—C34—C35	-12.2 (2)
O3—N2—C15—C14	-4.2 (2)	O7—N4—C34—C33	-10.9 (2)
O4—N2—C15—C14	177.10 (16)	O8—N4—C34—C33	168.55 (16)
C14—C15—C16—C17	1.0 (3)	C33—C34—C35—C36	-1.4 (3)
N2—C15—C16—C17	-179.36 (15)	N4—C34—C35—C36	179.41 (15)
C15—C16—C17—C12	0.4 (3)	C34—C35—C36—C31	-1.8 (2)
C13—C12—C17—C16	-1.4 (3)	C32—C31—C36—C35	2.9 (2)
C10—C12—C17—C16	179.45 (16)	C29—C31—C36—C35	-177.60 (15)

Fig. 1

