

tert-Butyl 2-[4-(2-{4-[(tert-butoxycarbonyl)methoxy]-3-methylphenyl}-2-propyl)-2-methylphenoxy]acetate

Qamar Ali,^a Sammer Yousuf,^a Muhammad Raza Shah^a and Seik Weng Ng^{b*}

^aHEJ Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan, and

^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 14 June 2010; accepted 17 June 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$;
 R factor = 0.060; wR factor = 0.184; data-to-parameter ratio = 20.2.

In the molecule of the title compound, $C_{29}H_{40}O_6$, the carbon atom belonging to the propyl chain is connected to two aromatic rings that open up the $C_{\text{aryl}}-\text{C}-C_{\text{aryl}}$ angle to $111.5(1)^\circ$. The four-atom $-\text{O}-\text{CH}_2-\text{C}(=\text{O})-\text{O}-$ linkage between the aromatic ring and the *tert*-butyl group assumes a (–)anti-periplanar conformation for one substituent and a (–)syn-periplanar conformation for the other substituent; the $\text{O}-\text{C}-\text{C}-\text{O}$ torsion angles are $-173.7(2)$ and $-10.2(3)^\circ$.

Related literature

For the crystal structure of a related V-shaped molecule, see: Shah *et al.* (2010).



Experimental

Crystal data

$C_{29}H_{40}O_6$
 $M_r = 484.61$
Triclinic, $P\bar{1}$
 $a = 8.3154(6)\text{ \AA}$
 $b = 12.5589(8)\text{ \AA}$
 $c = 13.9410(9)\text{ \AA}$
 $\alpha = 101.782(1)^\circ$
 $\beta = 97.529(1)^\circ$
 $\gamma = 94.156(1)^\circ$
 $V = 1405.52(16)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.08\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.45 \times 0.35 \times 0.30\text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
13784 measured reflections

6428 independent reflections
4482 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$
 $wR(F^2) = 0.184$
 $S = 1.03$
6428 reflections

318 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.35\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.19\text{ e \AA}^{-3}$

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

The authors thank the Higher Education Commission of Pakistan and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5103).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2002). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Shah, K., Yousuf, S., Raza Shah, M. & Ng, S. W. (2010). *Acta Cryst. E* **66**, o1705.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Westrip, S. P. (2010). publCIF. In preparation.

supplementary materials

Acta Cryst. (2010). E66, o1750 [doi:10.1107/S1600536810023433]

tert-Butyl 2-[4-(2-{4-[(*tert*-butoxycarbonyl)methoxy]-3-methylphenyl}-2-propyl)-2-methylphenoxy]acetate

Q. Ali, S. Yousuf, M. Raza Shah and S. W. Ng

Comment

We are interested in the solid-state structures of *V*-shaped molecules. A recent study reported the crystal structure of 9,9-bis[4-(*tert*-butoxycarbonylmethoxy)phenyl]fluorene, a compound used as dissolution inhibitor for protecting photosensitive poly(benzoxazole)s (Shah *et al.*, 2010). The *V* shape induced by the fluorenyl portion is now replaced by an *i*-propyl unit to furnish a similarly shaped molecule (Scheme I, Fig. 1). In the title molecule, the carbon atom belonging to the propyl chain is connected to two aromatic rings that open up the Caryl–C–Caryl to 111.5 (1) $^{\circ}$. The four-atom –O–CH₂–C(=O)–O– chain between the aromatic ring and the *tert*-butyl group assumes a *W* shape for one substituent and a *U* shape for the other substituent [O–C–C–O torsion angle -173.7 (2) $^{\circ}$ and -10.2 (3) $^{\circ}$].

Experimental

2,2-Bis(4-hydroxy-3-methylphenyl)propane (0.50 g) and potassium carbonate (0.85 g) were placed in acetone (25 ml) and to this was added *tert*-butyl bromoacetate (0.75 ml, 5 mmol). The mixture was stirred at room temperature for 3 h. The solvent was evaporated under reduced pressure and the residue was dissolved in a mixture of water (50 ml) and dichloromethane (50 ml). The aqueous layer was extracted three times with dichloromethane. The combined organic phases were evaporated under reduced pressure and the solid material was recrystallized from *n*-hexane. Colourless crystals were obtained in 80% yield.

Refinement

H atoms were placed in calculated positions [C–H = 0.93–0.97 Å] and were included in the refinement in the riding model approximation, with $U_{iso}(\text{H}) = 1.2\text{--}1.5U_{eq}(\text{C})$.

Figures




Fig. 1. Displacement ellipsoid plot (Barbour, 2001) of C₂₉H₄₀O₆ at the 50% probability level. H atoms are drawn as spheres of arbitrary radius.

tert-Butyl 2-[4-(2-{4-[(*tert*-butoxycarbonyl)methoxy]-3-methylphenyl}-2-propyl)-2-methylphenoxy]acetate

Crystal data

C₂₉H₄₀O₆

Z = 2

supplementary materials

$M_r = 484.61$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 8.3154 (6)$ Å

$b = 12.5589 (8)$ Å

$c = 13.9410 (9)$ Å

$\alpha = 101.782 (1)^\circ$

$\beta = 97.529 (1)^\circ$

$\gamma = 94.156 (1)^\circ$

$V = 1405.52 (16)$ Å³

$F(000) = 524$

$D_x = 1.145$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4014 reflections

$\theta = 2.5\text{--}27.3^\circ$

$\mu = 0.08$ mm⁻¹

$T = 293$ K

Block, colourless

$0.45 \times 0.35 \times 0.30$ mm

Data collection

Bruker SMART APEX
diffractometer

4482 reflections with $I > 2\sigma(I)$

Radiation source: fine-focus sealed tube

$R_{\text{int}} = 0.021$

graphite

$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 1.7^\circ$

ω scans

$h = -10 \rightarrow 10$

13784 measured reflections

$k = -16 \rightarrow 14$

6428 independent reflections

$l = -18 \rightarrow 17$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct
methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.060$

Hydrogen site location: inferred from neighbouring
sites

$wR(F^2) = 0.184$

H-atom parameters constrained

$S = 1.03$

$w = 1/[\sigma^2(F_o^2) + (0.0989P)^2 + 0.2549P]$

where $P = (F_o^2 + 2F_c^2)/3$

6428 reflections

$(\Delta/\sigma)_{\text{max}} = 0.001$

318 parameters

$\Delta\rho_{\text{max}} = 0.35$ e Å⁻³

0 restraints

$\Delta\rho_{\text{min}} = -0.19$ e Å⁻³

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.54867 (16)	0.16145 (12)	0.44990 (9)	0.0584 (4)
O2	0.28231 (18)	0.17998 (13)	0.40617 (11)	0.0644 (4)
O3	0.22035 (16)	0.13010 (10)	0.58548 (10)	0.0537 (3)
O4	0.13940 (17)	0.83903 (11)	1.01874 (10)	0.0608 (4)
O5	0.4354 (2)	0.79162 (18)	1.21064 (11)	0.0891 (6)
O6	0.43525 (16)	0.77133 (14)	1.04833 (10)	0.0665 (4)
C1	0.6069 (3)	0.17993 (18)	0.35761 (14)	0.0571 (5)
C2	0.7872 (3)	0.1715 (3)	0.3802 (2)	0.1093 (12)
H2A	0.8337	0.2275	0.4369	0.164*

H2B	0.8384	0.1808	0.3242	0.164*
H2C	0.8044	0.1010	0.3939	0.164*
C3	0.5730 (4)	0.29242 (19)	0.34262 (19)	0.0811 (7)
H3A	0.6249	0.3465	0.3994	0.122*
H3B	0.4575	0.2972	0.3344	0.122*
H3C	0.6150	0.3051	0.2846	0.122*
C4	0.5265 (4)	0.0919 (2)	0.27129 (18)	0.0882 (8)
H4A	0.4108	0.0961	0.2625	0.132*
H4B	0.5486	0.0216	0.2839	0.132*
H4C	0.5685	0.1016	0.2124	0.132*
C5	0.3936 (2)	0.16414 (14)	0.46314 (13)	0.0459 (4)
C6	0.3813 (2)	0.14131 (16)	0.56480 (14)	0.0516 (4)
H6A	0.4307	0.0747	0.5700	0.062*
H6B	0.4434	0.2005	0.6146	0.062*
C7	0.1432 (2)	0.22212 (13)	0.61535 (12)	0.0424 (4)
C8	0.2100 (2)	0.32846 (15)	0.62278 (14)	0.0501 (4)
H8	0.3146	0.3418	0.6082	0.060*
C9	0.1196 (2)	0.41480 (14)	0.65211 (14)	0.0506 (4)
H9	0.1661	0.4860	0.6585	0.061*
C10	-0.0379 (2)	0.39793 (13)	0.67218 (12)	0.0429 (4)
C11	-0.0997 (2)	0.29071 (14)	0.66524 (12)	0.0432 (4)
H11	-0.2045	0.2775	0.6795	0.052*
C12	-0.0121 (2)	0.20183 (13)	0.63788 (12)	0.0423 (4)
C13	-0.0854 (3)	0.08640 (15)	0.63031 (16)	0.0593 (5)
H13A	-0.1955	0.0875	0.6441	0.089*
H13B	-0.0223	0.0540	0.6774	0.089*
H13C	-0.0851	0.0443	0.5646	0.089*
C14	-0.1407 (2)	0.49513 (14)	0.69162 (14)	0.0480 (4)
C15	-0.3066 (2)	0.46110 (18)	0.7189 (2)	0.0755 (7)
H15A	-0.3641	0.4036	0.6669	0.113*
H15B	-0.3691	0.5228	0.7274	0.113*
H15C	-0.2904	0.4356	0.7796	0.113*
C16	-0.1702 (3)	0.53592 (18)	0.59406 (15)	0.0711 (7)
H16A	-0.0673	0.5558	0.5749	0.107*
H16B	-0.2309	0.5985	0.6037	0.107*
H16C	-0.2308	0.4788	0.5429	0.107*
C17	-0.0548 (2)	0.58614 (13)	0.77766 (12)	0.0403 (4)
C18	-0.0610 (2)	0.69584 (13)	0.77638 (12)	0.0402 (4)
H18	-0.1110	0.7143	0.7192	0.048*
C19	0.0042 (2)	0.77945 (13)	0.85692 (12)	0.0420 (4)
C20	0.0800 (2)	0.75110 (14)	0.94170 (13)	0.0445 (4)
C21	0.0898 (2)	0.64272 (16)	0.94550 (14)	0.0522 (4)
H21	0.1415	0.6243	1.0023	0.063*
C22	0.0220 (2)	0.56143 (15)	0.86392 (14)	0.0494 (4)
H22	0.0281	0.4886	0.8671	0.059*
C23	-0.0095 (3)	0.89746 (16)	0.85277 (17)	0.0642 (6)
H23A	0.0977	0.9358	0.8633	0.096*
H23B	-0.0696	0.9303	0.9034	0.096*
H23C	-0.0651	0.9015	0.7890	0.096*

supplementary materials

C24	0.1975 (2)	0.8205 (2)	1.11273 (14)	0.0632 (6)
H24A	0.1263	0.7618	1.1251	0.076*
H24B	0.1881	0.8857	1.1619	0.076*
C25	0.3708 (2)	0.79182 (16)	1.12921 (13)	0.0527 (5)
C26	0.6032 (2)	0.7409 (2)	1.04549 (17)	0.0702 (6)
C27	0.6246 (5)	0.6415 (3)	1.0903 (3)	0.1357 (14)
H27A	0.6150	0.6602	1.1595	0.203*
H27B	0.7303	0.6178	1.0827	0.203*
H27C	0.5419	0.5836	1.0572	0.203*
C28	0.7205 (4)	0.8347 (3)	1.1020 (3)	0.1313 (15)
H28A	0.7017	0.8982	1.0754	0.197*
H28B	0.8299	0.8172	1.0967	0.197*
H28C	0.7055	0.8493	1.1704	0.197*
C29	0.6096 (4)	0.7084 (4)	0.9371 (2)	0.1321 (15)
H29A	0.5931	0.7701	0.9073	0.198*
H29B	0.5257	0.6502	0.9072	0.198*
H29C	0.7142	0.6842	0.9270	0.198*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0552 (8)	0.0840 (10)	0.0451 (7)	0.0226 (7)	0.0167 (6)	0.0235 (7)
O2	0.0593 (8)	0.0816 (10)	0.0560 (8)	0.0185 (7)	0.0039 (7)	0.0224 (7)
O3	0.0603 (8)	0.0434 (7)	0.0634 (8)	0.0169 (6)	0.0273 (6)	0.0100 (6)
O4	0.0636 (8)	0.0574 (8)	0.0506 (8)	0.0147 (6)	-0.0056 (6)	-0.0076 (6)
O5	0.0735 (11)	0.1484 (18)	0.0461 (9)	0.0218 (11)	0.0038 (7)	0.0219 (9)
O6	0.0487 (8)	0.1064 (12)	0.0437 (7)	0.0157 (7)	0.0103 (6)	0.0098 (7)
C1	0.0673 (12)	0.0676 (13)	0.0446 (10)	0.0186 (10)	0.0208 (9)	0.0193 (9)
C2	0.0730 (17)	0.195 (4)	0.093 (2)	0.048 (2)	0.0443 (15)	0.075 (2)
C3	0.113 (2)	0.0603 (14)	0.0764 (16)	0.0089 (13)	0.0251 (14)	0.0213 (12)
C4	0.136 (2)	0.0705 (15)	0.0585 (14)	0.0123 (15)	0.0343 (15)	0.0028 (11)
C5	0.0526 (10)	0.0408 (9)	0.0452 (9)	0.0152 (7)	0.0102 (8)	0.0055 (7)
C6	0.0563 (11)	0.0560 (11)	0.0490 (10)	0.0241 (9)	0.0171 (8)	0.0141 (8)
C7	0.0503 (9)	0.0407 (9)	0.0375 (8)	0.0122 (7)	0.0120 (7)	0.0055 (7)
C8	0.0454 (9)	0.0462 (10)	0.0595 (11)	0.0042 (8)	0.0177 (8)	0.0074 (8)
C9	0.0538 (10)	0.0344 (9)	0.0622 (11)	0.0019 (7)	0.0144 (8)	0.0049 (8)
C10	0.0462 (9)	0.0370 (8)	0.0426 (9)	0.0070 (7)	0.0067 (7)	0.0004 (7)
C11	0.0430 (9)	0.0422 (9)	0.0429 (9)	0.0042 (7)	0.0113 (7)	0.0029 (7)
C12	0.0540 (10)	0.0355 (8)	0.0372 (8)	0.0059 (7)	0.0112 (7)	0.0041 (6)
C13	0.0721 (13)	0.0401 (10)	0.0686 (13)	0.0044 (9)	0.0302 (10)	0.0066 (9)
C14	0.0489 (10)	0.0385 (9)	0.0524 (10)	0.0110 (7)	0.0031 (8)	0.0010 (7)
C15	0.0450 (11)	0.0517 (12)	0.119 (2)	0.0112 (9)	0.0138 (11)	-0.0087 (12)
C16	0.0976 (17)	0.0548 (12)	0.0504 (11)	0.0289 (11)	-0.0129 (11)	-0.0050 (9)
C17	0.0419 (8)	0.0387 (8)	0.0405 (8)	0.0093 (7)	0.0107 (7)	0.0048 (7)
C18	0.0458 (9)	0.0396 (9)	0.0369 (8)	0.0100 (7)	0.0101 (7)	0.0079 (6)
C19	0.0441 (9)	0.0388 (9)	0.0443 (9)	0.0086 (7)	0.0130 (7)	0.0061 (7)
C20	0.0413 (9)	0.0461 (9)	0.0428 (9)	0.0090 (7)	0.0075 (7)	-0.0005 (7)
C21	0.0564 (11)	0.0567 (11)	0.0432 (9)	0.0166 (8)	0.0007 (8)	0.0105 (8)

C22	0.0586 (11)	0.0391 (9)	0.0525 (10)	0.0141 (8)	0.0060 (8)	0.0129 (8)
C23	0.0833 (15)	0.0417 (10)	0.0646 (13)	0.0103 (10)	0.0068 (11)	0.0058 (9)
C24	0.0547 (11)	0.0802 (15)	0.0449 (10)	0.0143 (10)	0.0049 (8)	-0.0102 (9)
C25	0.0516 (10)	0.0627 (12)	0.0382 (9)	0.0016 (8)	0.0045 (8)	0.0010 (8)
C26	0.0450 (11)	0.0988 (18)	0.0643 (13)	0.0143 (11)	0.0131 (9)	0.0065 (12)
C27	0.140 (3)	0.116 (3)	0.163 (4)	0.069 (2)	0.040 (3)	0.029 (3)
C28	0.0605 (16)	0.132 (3)	0.174 (4)	-0.0194 (17)	0.0303 (19)	-0.027 (3)
C29	0.087 (2)	0.228 (5)	0.081 (2)	0.045 (2)	0.0380 (17)	0.007 (2)

Geometric parameters (\AA , $^\circ$)

O1—C5	1.328 (2)	C13—H13C	0.96
O1—C1	1.488 (2)	C14—C15	1.533 (3)
O2—C5	1.194 (2)	C14—C17	1.534 (2)
O3—C7	1.379 (2)	C14—C16	1.545 (3)
O3—C6	1.409 (2)	C15—H15A	0.96
O4—C20	1.384 (2)	C15—H15B	0.96
O4—C24	1.407 (3)	C15—H15C	0.96
O5—C25	1.191 (2)	C16—H16A	0.96
O6—C25	1.298 (2)	C16—H16B	0.96
O6—C26	1.478 (2)	C16—H16C	0.96
C1—C4	1.502 (3)	C17—C18	1.386 (2)
C1—C2	1.507 (3)	C17—C22	1.389 (2)
C1—C3	1.511 (3)	C18—C19	1.391 (2)
C2—H2A	0.96	C18—H18	0.93
C2—H2B	0.96	C19—C20	1.389 (2)
C2—H2C	0.96	C19—C23	1.507 (3)
C3—H3A	0.96	C20—C21	1.381 (3)
C3—H3B	0.96	C21—C22	1.389 (3)
C3—H3C	0.96	C21—H21	0.93
C4—H4A	0.96	C22—H22	0.93
C4—H4B	0.96	C23—H23A	0.96
C4—H4C	0.96	C23—H23B	0.96
C5—C6	1.517 (3)	C23—H23C	0.96
C6—H6A	0.97	C24—C25	1.510 (3)
C6—H6B	0.97	C24—H24A	0.97
C7—C12	1.387 (2)	C24—H24B	0.97
C7—C8	1.387 (3)	C26—C28	1.488 (4)
C8—C9	1.388 (3)	C26—C29	1.492 (4)
C8—H8	0.93	C26—C27	1.517 (5)
C9—C10	1.385 (3)	C27—H27A	0.96
C9—H9	0.93	C27—H27B	0.96
C10—C11	1.386 (2)	C27—H27C	0.96
C10—C14	1.540 (2)	C28—H28A	0.96
C11—C12	1.392 (2)	C28—H28B	0.96
C11—H11	0.93	C28—H28C	0.96
C12—C13	1.509 (2)	C29—H29A	0.96
C13—H13A	0.96	C29—H29B	0.96
C13—H13B	0.96	C29—H29C	0.96

supplementary materials

C5—O1—C1	122.52 (14)	C14—C15—H15B	109.5
C7—O3—C6	119.71 (14)	H15A—C15—H15B	109.5
C20—O4—C24	119.61 (16)	C14—C15—H15C	109.5
C25—O6—C26	123.23 (16)	H15A—C15—H15C	109.5
O1—C1—C4	109.56 (18)	H15B—C15—H15C	109.5
O1—C1—C2	101.98 (16)	C14—C16—H16A	109.5
C4—C1—C2	111.8 (2)	C14—C16—H16B	109.5
O1—C1—C3	109.83 (17)	H16A—C16—H16B	109.5
C4—C1—C3	111.7 (2)	C14—C16—H16C	109.5
C2—C1—C3	111.5 (2)	H16A—C16—H16C	109.5
C1—C2—H2A	109.5	H16B—C16—H16C	109.5
C1—C2—H2B	109.5	C18—C17—C22	117.09 (15)
H2A—C2—H2B	109.5	C18—C17—C14	121.98 (15)
C1—C2—H2C	109.5	C22—C17—C14	120.75 (15)
H2A—C2—H2C	109.5	C17—C18—C19	122.83 (15)
H2B—C2—H2C	109.5	C17—C18—H18	118.6
C1—C3—H3A	109.5	C19—C18—H18	118.6
C1—C3—H3B	109.5	C20—C19—C18	118.19 (15)
H3A—C3—H3B	109.5	C20—C19—C23	120.99 (16)
C1—C3—H3C	109.5	C18—C19—C23	120.81 (16)
H3A—C3—H3C	109.5	C21—C20—O4	124.86 (16)
H3B—C3—H3C	109.5	C21—C20—C19	120.66 (16)
C1—C4—H4A	109.5	O4—C20—C19	114.47 (15)
C1—C4—H4B	109.5	C20—C21—C22	119.52 (16)
H4A—C4—H4B	109.5	C20—C21—H21	120.2
C1—C4—H4C	109.5	C22—C21—H21	120.2
H4A—C4—H4C	109.5	C21—C22—C17	121.70 (16)
H4B—C4—H4C	109.5	C21—C22—H22	119.2
O2—C5—O1	127.28 (17)	C17—C22—H22	119.2
O2—C5—C6	125.41 (17)	C19—C23—H23A	109.5
O1—C5—C6	107.31 (15)	C19—C23—H23B	109.5
O3—C6—C5	113.92 (15)	H23A—C23—H23B	109.5
O3—C6—H6A	108.8	C19—C23—H23C	109.5
C5—C6—H6A	108.8	H23A—C23—H23C	109.5
O3—C6—H6B	108.8	H23B—C23—H23C	109.5
C5—C6—H6B	108.8	O4—C24—C25	116.98 (17)
H6A—C6—H6B	107.7	O4—C24—H24A	108.1
O3—C7—C12	114.91 (15)	C25—C24—H24A	108.1
O3—C7—C8	124.60 (16)	O4—C24—H24B	108.1
C12—C7—C8	120.48 (15)	C25—C24—H24B	108.1
C7—C8—C9	119.50 (17)	H24A—C24—H24B	107.3
C7—C8—H8	120.2	O5—C25—O6	126.65 (19)
C9—C8—H8	120.2	O5—C25—C24	120.14 (18)
C10—C9—C8	121.84 (16)	O6—C25—C24	113.20 (16)
C10—C9—H9	119.1	O6—C26—C28	109.1 (2)
C8—C9—H9	119.1	O6—C26—C29	102.79 (19)
C9—C10—C11	116.96 (15)	C28—C26—C29	116.1 (3)
C9—C10—C14	119.77 (15)	O6—C26—C27	109.7 (2)
C11—C10—C14	123.03 (15)	C28—C26—C27	110.2 (3)

C10—C11—C12	123.05 (16)	C29—C26—C27	108.5 (3)
C10—C11—H11	118.5	C26—C27—H27A	109.5
C12—C11—H11	118.5	C26—C27—H27B	109.5
C7—C12—C11	118.11 (15)	H27A—C27—H27B	109.5
C7—C12—C13	120.80 (15)	C26—C27—H27C	109.5
C11—C12—C13	121.08 (16)	H27A—C27—H27C	109.5
C12—C13—H13A	109.5	H27B—C27—H27C	109.5
C12—C13—H13B	109.5	C26—C28—H28A	109.5
H13A—C13—H13B	109.5	C26—C28—H28B	109.5
C12—C13—H13C	109.5	H28A—C28—H28B	109.5
H13A—C13—H13C	109.5	C26—C28—H28C	109.5
H13B—C13—H13C	109.5	H28A—C28—H28C	109.5
C15—C14—C17	106.90 (16)	H28B—C28—H28C	109.5
C15—C14—C10	111.80 (15)	C26—C29—H29A	109.5
C17—C14—C10	111.47 (14)	C26—C29—H29B	109.5
C15—C14—C16	108.36 (18)	H29A—C29—H29B	109.5
C17—C14—C16	111.53 (15)	C26—C29—H29C	109.5
C10—C14—C16	106.79 (15)	H29A—C29—H29C	109.5
C14—C15—H15A	109.5	H29B—C29—H29C	109.5
C5—O1—C1—C4	-62.9 (2)	C15—C14—C17—C18	-95.7 (2)
C5—O1—C1—C2	178.6 (2)	C10—C14—C17—C18	141.89 (16)
C5—O1—C1—C3	60.2 (3)	C16—C14—C17—C18	22.6 (2)
C1—O1—C5—O2	0.6 (3)	C15—C14—C17—C22	79.1 (2)
C1—O1—C5—C6	179.73 (16)	C10—C14—C17—C22	-43.3 (2)
C7—O3—C6—C5	-80.1 (2)	C16—C14—C17—C22	-162.58 (17)
O2—C5—C6—O3	5.4 (3)	C22—C17—C18—C19	-0.6 (3)
O1—C5—C6—O3	-173.69 (15)	C14—C17—C18—C19	174.44 (15)
C6—O3—C7—C12	-177.31 (15)	C17—C18—C19—C20	0.6 (3)
C6—O3—C7—C8	3.2 (3)	C17—C18—C19—C23	-178.41 (17)
O3—C7—C8—C9	178.89 (17)	C24—O4—C20—C21	-8.0 (3)
C12—C7—C8—C9	-0.6 (3)	C24—O4—C20—C19	170.98 (16)
C7—C8—C9—C10	-1.5 (3)	C18—C19—C20—C21	0.0 (3)
C8—C9—C10—C11	2.3 (3)	C23—C19—C20—C21	178.97 (18)
C8—C9—C10—C14	-172.32 (17)	C18—C19—C20—O4	-179.11 (15)
C9—C10—C11—C12	-1.1 (3)	C23—C19—C20—O4	-0.1 (2)
C14—C10—C11—C12	173.34 (16)	O4—C20—C21—C22	178.45 (17)
O3—C7—C12—C11	-177.78 (14)	C19—C20—C21—C22	-0.5 (3)
C8—C7—C12—C11	1.8 (3)	C20—C21—C22—C17	0.6 (3)
O3—C7—C12—C13	0.7 (2)	C18—C17—C22—C21	0.0 (3)
C8—C7—C12—C13	-179.79 (18)	C14—C17—C22—C21	-175.09 (17)
C10—C11—C12—C7	-0.9 (3)	C20—O4—C24—C25	82.9 (2)
C10—C11—C12—C13	-179.34 (17)	C26—O6—C25—O5	1.7 (4)
C9—C10—C14—C15	-176.28 (18)	C26—O6—C25—C24	-179.7 (2)
C11—C10—C14—C15	9.4 (3)	O4—C24—C25—O5	168.5 (2)
C9—C10—C14—C17	-56.7 (2)	O4—C24—C25—O6	-10.2 (3)
C11—C10—C14—C17	129.00 (18)	C25—O6—C26—C28	-65.1 (3)
C9—C10—C14—C16	65.4 (2)	C25—O6—C26—C29	171.0 (3)
C11—C10—C14—C16	-109.0 (2)	C25—O6—C26—C27	55.7 (3)

supplementary materials

Fig. 1

