

## 4-[(9-Ethyl-9*H*-carbazol-3-yl)methylideneamino]-1,5-dimethyl-2-phenyl-1*H*-pyrazol-3(2*H*)-one

Abdullah M. Asiri,<sup>a</sup> Salman A. Khan,<sup>a</sup> Kong Wai Tan<sup>b</sup> and Seik Weng Ng<sup>b\*</sup>

<sup>a</sup>Chemistry Department, Faculty of Science, King Abdul Aziz University, PO Box 80203, Jeddah 21589, Saudi Arabia, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 15 June 2010; accepted 19 June 2010

Key indicators: single-crystal X-ray study;  $T = 100\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.038;  $wR$  factor = 0.104; data-to-parameter ratio = 16.8.

The imino–carbon double bond in the title Schiff base,  $C_{26}H_{24}N_4O$ , has an *E* configuration. The 13-membered carbazolyl fused-ring (r.m.s. deviation =  $0.056\text{ \AA}$ ) is nearly coplanar with five-membered pyrazole ring (r.m.s. deviation =  $0.036\text{ \AA}$ ) [dihedral angle between the two systems =  $10.4(1)^\circ$ ]; the phenyl substituent is twisted by  $51.1(1)^\circ$  with respect to the five-membered ring.

### Related literature

For background to this class of Schiff bases, see: Montalvo-González & Ariza-Castolo (2003).



### Experimental

#### Crystal data

$C_{26}H_{24}N_4O$   
 $M_r = 408.49$   
Monoclinic,  $P2_1/n$   
 $a = 10.4458(6)\text{ \AA}$   
 $b = 18.2674(11)\text{ \AA}$   
 $c = 10.8989(6)\text{ \AA}$   
 $\beta = 96.127(1)^\circ$

$V = 2067.8(2)\text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.08\text{ mm}^{-1}$   
 $T = 100\text{ K}$   
 $0.25 \times 0.20 \times 0.20\text{ mm}$

#### Data collection

Bruker SMART APEX  
diffractometer  
19772 measured reflections

4756 independent reflections  
4000 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.031$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$   
 $wR(F^2) = 0.104$   
 $S = 1.02$   
4756 reflections

283 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.24\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.25\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdul Aziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2169).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Montalvo-González, R. & Ariza-Castolo, A. (2003). *J. Mol. Struct.* **655**, 375–389.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.

## **supplementary materials**

*Acta Cryst.* (2010). E66, o1782 [doi:10.1107/S1600536810023779]

## 4-[(9-Ethyl-9*H*-carbazol-3-yl)methylideneamino]-1,5-dimethyl-2-phenyl-1*H*-pyrazol-3(2*H*)-one

A. M. Asiri, S. A. Khan, K. W. Tan and S. W. Ng

### Comment

4-Aminoantipyrine (4-amino-1,2-dihydro-1,5-dimethyl-2-phenyl-3*H*-pyrazol-3-one) possesses a pyrazolone unit, a feature that is important to the design of non-steroidal anti-inflammatory chemicals. Its amino group group allows the chemical to condense with aromatic aldehydes to yield Schiff bases. The crystal structures of a large number of such Schiff bases have been reported. For the Schiff base derived from the benzaldehyde homolog, the phenyl and pyrazoly rings are nearly coplanar (Montalvo-González & Ariza-Castolo, 2003). In the title carbazole-aldehyde analog (Scheme I, Fig. 1), the 13-membered carbazolyl fused-ring is nearly coplanar with 5-membered pyrazolyl ring [dihedral angle between the two systems 10.4 (1) °]. The phenyl substituent is twisted by 51.1 (1) ° with respect to the 5-membered ring.

### Experimental

9-Ethylcarbazole-3-carboxaldehyde (0.50 g, 2.2 mmol) and 4-aminoantipyrine (0.45 g, 2.2 mmol) here heated in methanol (15 ml) for 5 h to afford a light yellow precipitate. The solid material was collected and recrystallized from methanol.

### Refinement

Carbon-bound H-atoms were placed in calculated positions [C–H 0.95 to 0.98 Å,  $U(H)$  1.2 to 1.5  $U_{\text{eq}}(\text{C})$ ] and were included in the refinement in the riding model approximation.

### Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of  $\text{C}_{26}\text{H}_{24}\text{N}_4\text{O}$  at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

## 4-[(9-Ethyl-9*H*-carbazol-3-yl)methylideneamino]-1,5-dimethyl- 2-phenyl-1*H*-pyrazol-3(2*H*)-one

### Crystal data

$\text{C}_{26}\text{H}_{24}\text{N}_4\text{O}$	$F(000) = 864$
$M_r = 408.49$	$D_x = 1.312 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 7258 reflections
$a = 10.4458 (6) \text{ \AA}$	$\theta = 2.2\text{--}28.3^\circ$
$b = 18.2674 (11) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$c = 10.8989 (6) \text{ \AA}$	$T = 100 \text{ K}$

# supplementary materials

---

$\beta = 96.127(1)^\circ$  Prism, yellow  
 $V = 2067.8(2)\text{ \AA}^3$   $0.25 \times 0.20 \times 0.20\text{ mm}$   
 $Z = 4$

## Data collection

Bruker SMART APEX diffractometer  
Radiation source: fine-focus sealed tube  
graphite  
 $\omega$  scans  
19772 measured reflections  
4756 independent reflections

4000 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.031$   
 $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.2^\circ$   
 $h = -13 \rightarrow 13$   
 $k = -23 \rightarrow 23$   
 $l = -14 \rightarrow 14$

## Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.038$   
 $wR(F^2) = 0.104$   
 $S = 1.02$   
4756 reflections  
283 parameters  
0 restraints

Primary atom site location: structure-invariant direct methods  
Secondary atom site location: difference Fourier map  
Hydrogen site location: inferred from neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0525P)^2 + 0.720P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.24\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.25\text{ e \AA}^{-3}$

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.60645 (8)	0.54578 (4)	0.40022 (8)	0.02038 (19)
N1	1.35714 (9)	0.68186 (6)	0.65745 (9)	0.0206 (2)
N2	0.75630 (9)	0.69353 (6)	0.47222 (9)	0.0175 (2)
N3	0.43648 (9)	0.70479 (5)	0.32066 (9)	0.0178 (2)
N4	0.44409 (9)	0.62867 (5)	0.33728 (9)	0.0179 (2)
C1	1.36337 (11)	0.75748 (7)	0.66819 (11)	0.0196 (2)
C2	1.46850 (12)	0.80129 (8)	0.70917 (12)	0.0250 (3)
H2	1.5508	0.7805	0.7330	0.030*
C3	1.44906 (13)	0.87620 (8)	0.71402 (12)	0.0269 (3)
H3	1.5192	0.9072	0.7419	0.032*

## supplementary materials

---

C4	1.32833 (13)	0.90704 (7)	0.67875 (11)	0.0255 (3)
H4	1.3177	0.9586	0.6833	0.031*
C5	1.22368 (12)	0.86349 (7)	0.63713 (11)	0.0213 (3)
H5	1.1420	0.8848	0.6126	0.026*
C6	1.24060 (11)	0.78774 (7)	0.63197 (10)	0.0178 (2)
C7	1.15504 (11)	0.72649 (6)	0.60145 (10)	0.0165 (2)
C8	1.02461 (11)	0.72027 (6)	0.56055 (10)	0.0167 (2)
H8	0.9737	0.7629	0.5434	0.020*
C9	0.96945 (11)	0.65098 (6)	0.54505 (10)	0.0172 (2)
C10	1.04661 (12)	0.58849 (6)	0.57261 (11)	0.0193 (2)
H10	1.0070	0.5416	0.5660	0.023*
C11	1.17744 (11)	0.59244 (6)	0.60877 (11)	0.0196 (2)
H11	1.2283	0.5496	0.6241	0.023*
C12	1.23104 (11)	0.66235 (7)	0.62169 (10)	0.0179 (2)
C13	1.46091 (11)	0.63045 (7)	0.69520 (11)	0.0223 (3)
H13A	1.5445	0.6535	0.6832	0.027*
H13B	1.4509	0.5864	0.6420	0.027*
C14	1.46196 (15)	0.60749 (9)	0.82923 (13)	0.0357 (3)
H14A	1.5318	0.5723	0.8501	0.054*
H14B	1.3794	0.5847	0.8416	0.054*
H14C	1.4754	0.6506	0.8825	0.054*
C15	0.83444 (11)	0.63985 (6)	0.49658 (10)	0.0175 (2)
H15	0.8037	0.5913	0.4827	0.021*
C16	0.56810 (11)	0.60962 (6)	0.38971 (10)	0.0165 (2)
C17	0.63052 (11)	0.67924 (6)	0.42004 (10)	0.0163 (2)
C18	0.54672 (11)	0.73382 (6)	0.37959 (10)	0.0169 (2)
C19	0.56536 (12)	0.81422 (6)	0.39240 (11)	0.0208 (3)
H19A	0.5360	0.8382	0.3139	0.031*
H19B	0.6569	0.8248	0.4147	0.031*
H19C	0.5155	0.8327	0.4571	0.031*
C20	0.31008 (11)	0.73921 (7)	0.31473 (12)	0.0221 (3)
H20A	0.3203	0.7914	0.3339	0.033*
H20B	0.2596	0.7160	0.3748	0.033*
H20C	0.2654	0.7334	0.2316	0.033*
C21	0.35842 (11)	0.58088 (6)	0.26541 (10)	0.0170 (2)
C22	0.31925 (12)	0.59525 (7)	0.14157 (11)	0.0207 (3)
H22	0.3482	0.6380	0.1034	0.025*
C23	0.23735 (12)	0.54634 (7)	0.07451 (11)	0.0217 (3)
H23	0.2100	0.5560	-0.0099	0.026*
C24	0.19515 (11)	0.48364 (7)	0.12940 (11)	0.0209 (3)
H24	0.1394	0.4503	0.0830	0.025*
C25	0.23514 (12)	0.47000 (7)	0.25284 (11)	0.0208 (2)
H25	0.2067	0.4269	0.2906	0.025*
C26	0.31609 (11)	0.51841 (6)	0.32186 (11)	0.0186 (2)
H26	0.3422	0.5090	0.4066	0.022*

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0190 (4)	0.0166 (4)	0.0248 (4)	0.0015 (3)	-0.0012 (3)	0.0022 (3)
N1	0.0129 (5)	0.0230 (5)	0.0254 (5)	0.0016 (4)	-0.0005 (4)	-0.0015 (4)
N2	0.0141 (5)	0.0215 (5)	0.0164 (4)	-0.0007 (4)	-0.0006 (4)	-0.0016 (4)
N3	0.0157 (5)	0.0142 (5)	0.0227 (5)	0.0007 (4)	-0.0022 (4)	0.0005 (4)
N4	0.0160 (5)	0.0145 (5)	0.0222 (5)	-0.0003 (4)	-0.0030 (4)	0.0019 (4)
C1	0.0169 (6)	0.0236 (6)	0.0187 (5)	-0.0015 (5)	0.0040 (4)	-0.0004 (4)
C2	0.0173 (6)	0.0327 (7)	0.0250 (6)	-0.0050 (5)	0.0031 (5)	-0.0012 (5)
C3	0.0263 (6)	0.0318 (7)	0.0229 (6)	-0.0139 (5)	0.0040 (5)	-0.0009 (5)
C4	0.0346 (7)	0.0208 (6)	0.0214 (6)	-0.0079 (5)	0.0043 (5)	0.0010 (5)
C5	0.0249 (6)	0.0203 (6)	0.0186 (5)	-0.0017 (5)	0.0024 (5)	0.0018 (4)
C6	0.0166 (5)	0.0219 (6)	0.0153 (5)	-0.0020 (4)	0.0029 (4)	0.0001 (4)
C7	0.0174 (6)	0.0171 (5)	0.0153 (5)	0.0005 (4)	0.0027 (4)	0.0000 (4)
C8	0.0162 (5)	0.0178 (5)	0.0159 (5)	0.0017 (4)	0.0009 (4)	0.0002 (4)
C9	0.0155 (5)	0.0204 (6)	0.0154 (5)	0.0008 (4)	0.0009 (4)	-0.0015 (4)
C10	0.0207 (6)	0.0167 (6)	0.0201 (5)	-0.0003 (4)	0.0006 (4)	-0.0033 (4)
C11	0.0190 (6)	0.0175 (6)	0.0218 (6)	0.0042 (4)	0.0002 (4)	-0.0023 (4)
C12	0.0142 (5)	0.0222 (6)	0.0173 (5)	0.0018 (4)	0.0009 (4)	-0.0012 (4)
C13	0.0141 (5)	0.0283 (6)	0.0243 (6)	0.0055 (5)	0.0010 (4)	-0.0021 (5)
C14	0.0337 (8)	0.0478 (9)	0.0254 (7)	0.0170 (7)	0.0021 (6)	0.0019 (6)
C15	0.0174 (5)	0.0190 (6)	0.0159 (5)	-0.0016 (4)	0.0004 (4)	-0.0016 (4)
C16	0.0142 (5)	0.0212 (6)	0.0138 (5)	-0.0005 (4)	0.0002 (4)	0.0017 (4)
C17	0.0157 (5)	0.0185 (6)	0.0147 (5)	-0.0008 (4)	0.0011 (4)	-0.0008 (4)
C18	0.0157 (5)	0.0198 (6)	0.0149 (5)	-0.0013 (4)	0.0009 (4)	-0.0016 (4)
C19	0.0218 (6)	0.0172 (6)	0.0226 (6)	0.0001 (5)	-0.0017 (5)	-0.0023 (4)
C20	0.0167 (6)	0.0225 (6)	0.0265 (6)	0.0028 (5)	-0.0001 (5)	0.0001 (5)
C21	0.0128 (5)	0.0174 (5)	0.0202 (5)	-0.0003 (4)	0.0000 (4)	-0.0006 (4)
C22	0.0215 (6)	0.0188 (6)	0.0215 (6)	-0.0037 (5)	0.0003 (5)	0.0039 (4)
C23	0.0238 (6)	0.0216 (6)	0.0188 (6)	-0.0006 (5)	-0.0024 (5)	0.0012 (5)
C24	0.0182 (6)	0.0187 (6)	0.0251 (6)	-0.0020 (5)	-0.0001 (5)	-0.0036 (5)
C25	0.0203 (6)	0.0168 (6)	0.0256 (6)	-0.0017 (5)	0.0043 (5)	0.0025 (5)
C26	0.0172 (5)	0.0199 (6)	0.0185 (5)	0.0017 (4)	0.0020 (4)	0.0021 (4)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O1—C16	1.2344 (14)	C11—C12	1.3956 (17)
N1—C12	1.3799 (15)	C11—H11	0.9500
N1—C1	1.3873 (16)	C13—C14	1.5187 (18)
N1—C13	1.4598 (15)	C13—H13A	0.9900
N2—C15	1.2853 (15)	C13—H13B	0.9900
N2—C17	1.3994 (14)	C14—H14A	0.9800
N3—C18	1.3646 (14)	C14—H14B	0.9800
N3—N4	1.4034 (13)	C14—H14C	0.9800
N3—C20	1.4577 (15)	C15—H15	0.9500
N4—C16	1.4028 (14)	C16—C17	1.4511 (16)
N4—C21	1.4237 (14)	C17—C18	1.3682 (16)

C1—C2	1.3933 (17)	C18—C19	1.4861 (16)
C1—C6	1.4134 (16)	C19—H19A	0.9800
C2—C3	1.385 (2)	C19—H19B	0.9800
C2—H2	0.9500	C19—H19C	0.9800
C3—C4	1.397 (2)	C20—H20A	0.9800
C3—H3	0.9500	C20—H20B	0.9800
C4—C5	1.3884 (17)	C20—H20C	0.9800
C4—H4	0.9500	C21—C26	1.3907 (16)
C5—C6	1.3969 (17)	C21—C22	1.3929 (16)
C5—H5	0.9500	C22—C23	1.3895 (16)
C6—C7	1.4484 (16)	C22—H22	0.9500
C7—C8	1.3922 (16)	C23—C24	1.3857 (17)
C7—C12	1.4192 (16)	C23—H23	0.9500
C8—C9	1.3935 (16)	C24—C25	1.3880 (17)
C8—H8	0.9500	C24—H24	0.9500
C9—C10	1.4112 (16)	C25—C26	1.3881 (16)
C9—C15	1.4664 (15)	C25—H25	0.9500
C10—C11	1.3832 (16)	C26—H26	0.9500
C10—H10	0.9500		
C12—N1—C1	108.40 (10)	C14—C13—H13B	109.2
C12—N1—C13	124.84 (10)	H13A—C13—H13B	107.9
C1—N1—C13	126.16 (10)	C13—C14—H14A	109.5
C15—N2—C17	119.32 (10)	C13—C14—H14B	109.5
C18—N3—N4	106.92 (9)	H14A—C14—H14B	109.5
C18—N3—C20	124.62 (10)	C13—C14—H14C	109.5
N4—N3—C20	118.15 (9)	H14A—C14—H14C	109.5
C16—N4—N3	109.60 (9)	H14B—C14—H14C	109.5
C16—N4—C21	124.82 (10)	N2—C15—C9	122.27 (11)
N3—N4—C21	120.86 (9)	N2—C15—H15	118.9
N1—C1—C2	128.85 (11)	C9—C15—H15	118.9
N1—C1—C6	109.44 (10)	O1—C16—N4	123.27 (10)
C2—C1—C6	121.70 (12)	O1—C16—C17	132.33 (11)
C3—C2—C1	117.77 (12)	N4—C16—C17	104.38 (9)
C3—C2—H2	121.1	C18—C17—N2	122.39 (10)
C1—C2—H2	121.1	C18—C17—C16	107.99 (10)
C2—C3—C4	121.29 (12)	N2—C17—C16	129.46 (10)
C2—C3—H3	119.4	N3—C18—C17	110.31 (10)
C4—C3—H3	119.4	N3—C18—C19	121.59 (10)
C5—C4—C3	121.00 (12)	C17—C18—C19	128.09 (10)
C5—C4—H4	119.5	C18—C19—H19A	109.5
C3—C4—H4	119.5	C18—C19—H19B	109.5
C4—C5—C6	118.82 (12)	H19A—C19—H19B	109.5
C4—C5—H5	120.6	C18—C19—H19C	109.5
C6—C5—H5	120.6	H19A—C19—H19C	109.5
C5—C6—C1	119.41 (11)	H19B—C19—H19C	109.5
C5—C6—C7	134.17 (11)	N3—C20—H20A	109.5
C1—C6—C7	106.34 (10)	N3—C20—H20B	109.5
C8—C7—C12	119.64 (10)	H20A—C20—H20B	109.5
C8—C7—C6	134.06 (11)	N3—C20—H20C	109.5

## supplementary materials

---

C12—C7—C6	106.30 (10)	H20A—C20—H20C	109.5
C7—C8—C9	119.39 (11)	H20B—C20—H20C	109.5
C7—C8—H8	120.3	C26—C21—C22	120.59 (11)
C9—C8—H8	120.3	C26—C21—N4	118.08 (10)
C8—C9—C10	119.31 (10)	C22—C21—N4	121.33 (10)
C8—C9—C15	122.61 (10)	C23—C22—C21	119.29 (11)
C10—C9—C15	118.04 (10)	C23—C22—H22	120.4
C11—C10—C9	122.89 (11)	C21—C22—H22	120.4
C11—C10—H10	118.6	C24—C23—C22	120.71 (11)
C9—C10—H10	118.6	C24—C23—H23	119.6
C10—C11—C12	116.75 (11)	C22—C23—H23	119.6
C10—C11—H11	121.6	C23—C24—C25	119.35 (11)
C12—C11—H11	121.6	C23—C24—H24	120.3
N1—C12—C11	128.71 (11)	C25—C24—H24	120.3
N1—C12—C7	109.39 (10)	C26—C25—C24	120.92 (11)
C11—C12—C7	121.87 (10)	C26—C25—H25	119.5
N1—C13—C14	112.18 (10)	C24—C25—H25	119.5
N1—C13—H13A	109.2	C25—C26—C21	119.14 (11)
C14—C13—H13A	109.2	C25—C26—H26	120.4
N1—C13—H13B	109.2	C21—C26—H26	120.4
C18—N3—N4—C16	-9.38 (12)	C8—C7—C12—C11	-3.85 (17)
C20—N3—N4—C16	-156.12 (10)	C6—C7—C12—C11	175.87 (11)
C18—N3—N4—C21	-166.14 (10)	C12—N1—C13—C14	-81.94 (15)
C20—N3—N4—C21	47.12 (15)	C1—N1—C13—C14	88.12 (15)
C12—N1—C1—C2	175.55 (12)	C17—N2—C15—C9	176.64 (10)
C13—N1—C1—C2	4.1 (2)	C8—C9—C15—N2	-4.26 (18)
C12—N1—C1—C6	-3.37 (13)	C10—C9—C15—N2	178.02 (11)
C13—N1—C1—C6	-174.79 (11)	N3—N4—C16—O1	-170.71 (10)
N1—C1—C2—C3	-178.53 (12)	C21—N4—C16—O1	-15.08 (18)
C6—C1—C2—C3	0.28 (18)	N3—N4—C16—C17	7.67 (12)
C1—C2—C3—C4	-0.18 (19)	C21—N4—C16—C17	163.30 (10)
C2—C3—C4—C5	-0.24 (19)	C15—N2—C17—C18	-172.81 (11)
C3—C4—C5—C6	0.56 (18)	C15—N2—C17—C16	1.92 (18)
C4—C5—C6—C1	-0.46 (17)	O1—C16—C17—C18	174.95 (12)
C4—C5—C6—C7	175.81 (12)	N4—C16—C17—C18	-3.22 (12)
N1—C1—C6—C5	179.06 (10)	O1—C16—C17—N2	-0.4 (2)
C2—C1—C6—C5	0.04 (17)	N4—C16—C17—N2	-178.54 (11)
N1—C1—C6—C7	1.84 (13)	N4—N3—C18—C17	7.25 (13)
C2—C1—C6—C7	-177.17 (11)	C20—N3—C18—C17	151.26 (11)
C5—C6—C7—C8	3.4 (2)	N4—N3—C18—C19	-173.40 (10)
C1—C6—C7—C8	179.98 (12)	C20—N3—C18—C19	-29.39 (17)
C5—C6—C7—C12	-176.29 (13)	N2—C17—C18—N3	173.21 (10)
C1—C6—C7—C12	0.32 (12)	C16—C17—C18—N3	-2.51 (13)
C12—C7—C8—C9	2.68 (17)	N2—C17—C18—C19	-6.09 (19)
C6—C7—C8—C9	-176.94 (12)	C16—C17—C18—C19	178.19 (11)
C7—C8—C9—C10	0.78 (16)	C16—N4—C21—C26	62.98 (15)
C7—C8—C9—C15	-176.91 (10)	N3—N4—C21—C26	-143.95 (11)
C8—C9—C10—C11	-3.41 (17)	C16—N4—C21—C22	-116.02 (13)
C15—C9—C10—C11	174.39 (11)	N3—N4—C21—C22	37.05 (16)

## supplementary materials

---

C9—C10—C11—C12	2.29 (18)	C26—C21—C22—C23	-0.13 (18)
C1—N1—C12—C11	-174.53 (12)	N4—C21—C22—C23	178.84 (11)
C13—N1—C12—C11	-2.97 (19)	C21—C22—C23—C24	-0.31 (19)
C1—N1—C12—C7	3.58 (13)	C22—C23—C24—C25	0.22 (19)
C13—N1—C12—C7	175.14 (10)	C23—C24—C25—C26	0.31 (18)
C10—C11—C12—N1	179.23 (11)	C24—C25—C26—C21	-0.74 (18)
C10—C11—C12—C7	1.34 (17)	C22—C21—C26—C25	0.65 (18)
C8—C7—C12—N1	177.90 (10)	N4—C21—C26—C25	-178.36 (10)
C6—C7—C12—N1	-2.39 (13)		

## supplementary materials

---

Fig. 1

