

1,5-Dimethyl-3-propargyl-1*H*-1,5-benzodiazepine-2,4(3*H*,5*H*)-dione

Rachid Dardouri,^a F. Ouazzani Chahdi,^a Natalie Saffon,^b El Mokhtar Essassi^c and Seik Weng Ng^{d*}

^aLaboratoire de Chimie Organique Appliquée, Faculté des Sciences et Techniques, Université Sidi Mohamed Ben Abdallah, Fès, Morocco, ^bService Commun Rayons-X FR2599, Université Paul Sabatier, Bâtiment 2R1, 118 route de Narbonne, Toulouse, France, ^cLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, and ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 21 June 2010; accepted 22 June 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.038; wR factor = 0.105; data-to-parameter ratio = 10.6.

The asymmetric unit of the title compound, $\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_2$, comprises two independent molecules, which slightly differ in the orientation of the propargyl chain. In both molecules, the diazepine ring adopts a boat conformation with the propargyl-bearing C atom as the prow and the C atoms at the ring junction as the stern. The carbonyl O atom of one independent molecule is hydrogen bonded to the acetylenic H atom of the other independent molecule. In the crystal, symmetry-related molecules are linked together by $\text{C}-\text{H} \cdots \text{O}$ hydrogen bonds, forming a ribbon-like structure along the c axis.

Related literature

For a related structure, see: Jabli *et al.* (2009).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_2$	$V = 2457.42(8)\text{ \AA}^3$
$M_r = 242.27$	$Z = 8$
Monoclinic, Cc	Mo $K\alpha$ radiation
$a = 16.0768(3)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 17.1087(3)\text{ \AA}$	$T = 293\text{ K}$
$c = 8.9530(2)\text{ \AA}$	$0.40 \times 0.30 \times 0.05\text{ mm}$
$\beta = 93.701(1)^\circ$	

Data collection

Bruker X8 APEXII area-detector diffractometer	3580 independent reflections
21953 measured reflections	2637 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.105$	$\Delta\rho_{\text{max}} = 0.22\text{ e \AA}^{-3}$
$S = 1.06$	$\Delta\rho_{\text{min}} = -0.19\text{ e \AA}^{-3}$
3580 reflections	
337 parameters	
4 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
C13—H13 \cdots O4	0.93 (3)	2.44 (3)	3.366 (3)	172 (3)
C8—H8 \cdots O2 ⁱ	0.98	2.56	3.536 (3)	175
C19—H19 \cdots O1 ⁱ	0.93	2.49	3.374 (3)	160

Symmetry code: (i) $x, -y + 1, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5110).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jabli, H., Ouazzani Chahdi, F., Garrigues, B., Essassi, E. M. & Ng, S. W. (2009). *Acta Cryst. E65*, o3149.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, Submitted.

supplementary materials

Acta Cryst. (2010). E66, o1797 [doi:10.1107/S1600536810024219]

1,5-Dimethyl-3-propargyl-1*H*-1,5-benzodiazepine-2,4(3*H*,5*H*)-dione

R. Dardouri, F. Ouazzani Chahdi, N. Saffon, E. M. Essassi and S. W. Ng

Comment

We recently reported the crystal structure of 1,5-dibenzyl-3-propargyl-1,5-benzodiazepine-2,4-dione, a compound readily synthesized by reacting the disubstituted 1,5-benzodiazepine-2,4-dione with propargyl bromide (Jabli *et al.*, 2009). The background to the study of such compounds is given in other reports. Replacing the benzyl unit by a methyl unit in the present study gives a compound having a diazepine ring system (Scheme I, Fig. 1). This ring adopts a boat conformation (with the propargyl-bearing C atom as the prow and the fused-ring C atoms as the stern). There are two independent molecules; the acetylenic H-atom of one molecule forms a hydrogen bond to the carbonyl O-atom of the other independent molecule.

Experimental

To a solution of potassium *t*-butoxide (0.42 g, 3.6 mmol) in DMF (15 ml) was added 1,5-dimethyl-1,5-benzodiazepine-2,4-dione (0.50 g, 2.4 mmol) and propargyl bromide (0.26 ml, 2.87 mmol). Stirring was continued for 24 h. The reaction was monitored by thin layer chromatography. On completion of the reaction, the mixture was filtered; crystals were obtained when the solvent was allowed to evaporate.

Refinement

The acetylenic H atoms were located in a difference Fourier map and were refined with C–H distances restrained to 0.93 (1) Å; their U_{iso} parameters were freely refined. The remaining H atoms were placed in calculated positions (C–H = 0.93–0.98 Å) and were included in the refinement in the riding-model approximation, with $U_{iso}(\text{H})$ set to 1.2–1.5 $U_{eq}(\text{C})$. In the absence of significant anomalous scattering effects, 3266 Friedel pairs were averaged.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of the two independent molecules of $\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_2$ at the 50% probability level shown as a hydrogen-bonded dimer; H atoms are drawn as spheres of arbitrary radius. The dashed line denotes a hydrogen bond.

1,5-Dimethyl-3-propargyl-1*H*-1,5-benzodiazepine- 2,4(3*H*,5*H*)-dione

Crystal data

$\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_2$

$F(000) = 1024$

$M_r = 242.27$

$D_x = 1.310 \text{ Mg m}^{-3}$

Monoclinic, Cc

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Hall symbol: C -2yc

Cell parameters from 5448 reflections

supplementary materials

$a = 16.0768 (3) \text{ \AA}$	$\theta = 2.5\text{--}29.4^\circ$
$b = 17.1087 (3) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 8.9530 (2) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 93.701 (1)^\circ$	Plate, colourless
$V = 2457.42 (8) \text{ \AA}^3$	$0.40 \times 0.30 \times 0.05 \text{ mm}$
$Z = 8$	

Data collection

Bruker X8 APEXII area-detector diffractometer	2637 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.049$
graphite	$\theta_{\text{max}} = 30.0^\circ, \theta_{\text{min}} = 2.8^\circ$
φ and ω scans	$h = -22\text{--}22$
21953 measured reflections	$k = -24\text{--}23$
3580 independent reflections	$l = -12\text{--}12$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.038$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.105$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0514P)^2 + 0.5357P]$ where $P = (F_o^2 + 2F_c^2)/3$
3580 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
337 parameters	$\Delta\rho_{\text{max}} = 0.22 \text{ e \AA}^{-3}$
4 restraints	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.50007 (11)	0.33109 (11)	0.5000 (2)	0.0339 (4)
O2	0.34014 (11)	0.41769 (12)	0.21819 (19)	0.0332 (4)
O3	0.99978 (11)	0.63153 (11)	0.9419 (2)	0.0363 (4)
O4	0.78408 (12)	0.58439 (13)	0.74162 (19)	0.0380 (5)
N1	0.39120 (12)	0.33841 (12)	0.6473 (2)	0.0268 (4)
N2	0.26570 (12)	0.39080 (12)	0.4189 (2)	0.0253 (4)
N3	0.91197 (13)	0.68624 (13)	1.1007 (2)	0.0293 (5)
N4	0.75177 (12)	0.65404 (14)	0.9454 (2)	0.0298 (5)
C1	0.31443 (15)	0.37280 (14)	0.6847 (3)	0.0251 (5)
C2	0.29827 (17)	0.38043 (16)	0.8365 (3)	0.0318 (6)
H2	0.3393	0.3670	0.9099	0.038*
C3	0.22235 (19)	0.40758 (17)	0.8778 (3)	0.0382 (6)
H3	0.2124	0.4121	0.9786	0.046*

C4	0.16095 (18)	0.42805 (17)	0.7692 (3)	0.0384 (6)
H4	0.1091	0.4449	0.7969	0.046*
C5	0.17692 (16)	0.42341 (15)	0.6190 (3)	0.0308 (5)
H5	0.1359	0.4385	0.5467	0.037*
C6	0.25334 (15)	0.39661 (13)	0.5746 (3)	0.0240 (5)
C7	0.33523 (15)	0.41709 (14)	0.3543 (3)	0.0242 (5)
C8	0.40776 (14)	0.44186 (14)	0.4628 (2)	0.0232 (5)
H8	0.3882	0.4782	0.5375	0.028*
C9	0.43852 (14)	0.36629 (14)	0.5383 (3)	0.0250 (5)
C10	0.19504 (16)	0.36807 (17)	0.3156 (3)	0.0329 (6)
H10A	0.2141	0.3334	0.2408	0.049*
H10B	0.1711	0.4139	0.2682	0.049*
H10C	0.1537	0.3421	0.3704	0.049*
C11	0.47672 (15)	0.48013 (16)	0.3774 (3)	0.0295 (5)
H11A	0.4954	0.4434	0.3040	0.035*
H11B	0.4540	0.5255	0.3239	0.035*
C12	0.54896 (16)	0.50462 (16)	0.4769 (3)	0.0324 (5)
C13	0.60798 (19)	0.5229 (2)	0.5539 (4)	0.0446 (7)
H13	0.6549 (15)	0.537 (2)	0.614 (4)	0.062 (11)*
C14	0.42070 (18)	0.26755 (16)	0.7274 (3)	0.0373 (6)
H14A	0.4325	0.2277	0.6563	0.056*
H14B	0.3784	0.2493	0.7898	0.056*
H14C	0.4704	0.2794	0.7884	0.056*
C15	0.83776 (15)	0.68259 (15)	1.1784 (3)	0.0279 (5)
C16	0.84147 (18)	0.69589 (16)	1.3331 (3)	0.0343 (6)
H16	0.8919	0.7095	1.3829	0.041*
C17	0.7709 (2)	0.68902 (17)	1.4124 (3)	0.0404 (7)
H17	0.7745	0.6968	1.5154	0.049*
C18	0.6949 (2)	0.67062 (18)	1.3391 (3)	0.0429 (7)
H18	0.6477	0.6658	1.3932	0.052*
C19	0.68919 (17)	0.65948 (16)	1.1858 (3)	0.0357 (6)
H19	0.6378	0.6481	1.1369	0.043*
C20	0.75997 (16)	0.66527 (14)	1.1039 (3)	0.0278 (5)
C21	0.79742 (15)	0.60039 (16)	0.8740 (3)	0.0281 (5)
C22	0.87014 (15)	0.56424 (15)	0.9697 (3)	0.0274 (5)
H22	0.8497	0.5453	1.0640	0.033*
C23	0.93407 (15)	0.62914 (15)	1.0034 (3)	0.0272 (5)
C24	0.67961 (17)	0.68895 (18)	0.8601 (3)	0.0386 (6)
H24A	0.6957	0.7049	0.7634	0.058*
H24B	0.6605	0.7336	0.9131	0.058*
H24C	0.6356	0.6511	0.8484	0.058*
C25	0.90761 (17)	0.49596 (16)	0.8873 (3)	0.0330 (6)
H25A	0.8631	0.4621	0.8477	0.040*
H25B	0.9365	0.5159	0.8034	0.040*
C26	0.96610 (17)	0.45014 (16)	0.9847 (3)	0.0326 (6)
C27	1.01106 (19)	0.41189 (19)	1.0637 (4)	0.0445 (7)
H27	1.043 (2)	0.3804 (19)	1.130 (4)	0.070 (12)*
C28	0.97458 (18)	0.74656 (18)	1.1443 (4)	0.0426 (7)
H28A	1.0095	0.7556	1.0630	0.064*

supplementary materials

H28B	1.0082	0.7291	1.2305	0.064*
H28C	0.9468	0.7942	1.1679	0.064*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0256 (9)	0.0397 (10)	0.0365 (11)	0.0051 (8)	0.0020 (7)	-0.0001 (8)
O2	0.0345 (9)	0.0472 (11)	0.0175 (9)	-0.0022 (8)	-0.0014 (7)	0.0014 (8)
O3	0.0284 (10)	0.0455 (11)	0.0350 (11)	-0.0010 (8)	0.0007 (8)	-0.0021 (9)
O4	0.0363 (10)	0.0558 (13)	0.0210 (10)	-0.0022 (9)	-0.0054 (8)	-0.0026 (8)
N1	0.0276 (10)	0.0311 (10)	0.0210 (10)	0.0018 (8)	-0.0029 (8)	0.0052 (9)
N2	0.0234 (10)	0.0323 (11)	0.0197 (10)	-0.0023 (8)	-0.0022 (8)	-0.0003 (8)
N3	0.0267 (10)	0.0350 (12)	0.0254 (11)	-0.0037 (8)	-0.0038 (8)	-0.0050 (9)
N4	0.0256 (10)	0.0398 (12)	0.0233 (11)	-0.0010 (8)	-0.0030 (8)	0.0021 (9)
C1	0.0275 (12)	0.0277 (12)	0.0202 (11)	-0.0040 (9)	0.0019 (9)	0.0009 (10)
C2	0.0372 (13)	0.0376 (14)	0.0204 (12)	-0.0069 (11)	0.0008 (10)	0.0013 (10)
C3	0.0507 (16)	0.0403 (15)	0.0249 (13)	-0.0065 (12)	0.0139 (12)	-0.0015 (12)
C4	0.0376 (15)	0.0383 (15)	0.0411 (16)	-0.0005 (12)	0.0161 (12)	-0.0007 (13)
C5	0.0272 (12)	0.0342 (13)	0.0312 (13)	-0.0017 (10)	0.0034 (10)	0.0002 (11)
C6	0.0256 (11)	0.0261 (11)	0.0202 (11)	-0.0039 (9)	0.0015 (9)	0.0003 (9)
C7	0.0244 (11)	0.0288 (12)	0.0194 (11)	0.0038 (9)	-0.0003 (8)	0.0019 (9)
C8	0.0243 (11)	0.0293 (12)	0.0162 (10)	-0.0010 (9)	0.0022 (8)	-0.0013 (9)
C9	0.0230 (11)	0.0309 (12)	0.0202 (11)	-0.0016 (9)	-0.0056 (9)	-0.0008 (9)
C10	0.0276 (13)	0.0400 (15)	0.0295 (14)	-0.0048 (11)	-0.0102 (10)	-0.0004 (11)
C11	0.0279 (12)	0.0380 (14)	0.0227 (12)	-0.0040 (10)	0.0017 (9)	0.0037 (10)
C12	0.0308 (13)	0.0374 (13)	0.0296 (13)	-0.0036 (11)	0.0060 (11)	0.0054 (11)
C13	0.0379 (15)	0.0542 (19)	0.0412 (17)	-0.0145 (14)	-0.0016 (13)	0.0030 (14)
C14	0.0414 (15)	0.0365 (15)	0.0335 (14)	0.0051 (12)	-0.0019 (11)	0.0121 (12)
C15	0.0316 (13)	0.0288 (13)	0.0231 (12)	0.0016 (10)	-0.0010 (10)	-0.0013 (10)
C16	0.0438 (15)	0.0345 (14)	0.0238 (13)	0.0031 (11)	-0.0047 (11)	-0.0013 (11)
C17	0.0598 (18)	0.0374 (15)	0.0244 (13)	0.0026 (13)	0.0047 (13)	0.0006 (12)
C18	0.0497 (17)	0.0412 (17)	0.0401 (17)	-0.0037 (13)	0.0199 (14)	0.0010 (13)
C19	0.0337 (14)	0.0384 (15)	0.0356 (15)	-0.0055 (11)	0.0056 (12)	-0.0004 (12)
C20	0.0317 (12)	0.0284 (12)	0.0233 (12)	-0.0028 (10)	0.0011 (10)	0.0004 (10)
C21	0.0254 (11)	0.0374 (13)	0.0210 (12)	-0.0078 (10)	-0.0016 (9)	0.0018 (10)
C22	0.0312 (12)	0.0335 (13)	0.0170 (11)	-0.0003 (10)	-0.0016 (9)	0.0007 (10)
C23	0.0250 (11)	0.0353 (13)	0.0202 (12)	0.0019 (9)	-0.0056 (9)	0.0023 (10)
C24	0.0291 (13)	0.0506 (17)	0.0351 (15)	0.0009 (12)	-0.0066 (11)	0.0030 (13)
C25	0.0422 (14)	0.0367 (14)	0.0197 (12)	-0.0012 (11)	-0.0007 (11)	-0.0029 (10)
C26	0.0341 (13)	0.0362 (14)	0.0279 (13)	-0.0034 (11)	0.0051 (10)	-0.0044 (11)
C27	0.0344 (15)	0.0495 (18)	0.0488 (19)	-0.0012 (13)	-0.0036 (13)	0.0078 (14)
C28	0.0351 (14)	0.0497 (17)	0.0423 (17)	-0.0121 (13)	-0.0030 (12)	-0.0113 (14)

Geometric parameters (\AA , $^\circ$)

O1—C9	1.226 (3)	C11—C12	1.478 (4)
O2—C7	1.226 (3)	C11—H11A	0.97
O3—C23	1.223 (3)	C11—H11B	0.97
O4—C21	1.222 (3)	C12—C13	1.179 (4)

N1—C9	1.361 (3)	C13—H13	0.93 (3)
N1—C1	1.427 (3)	C14—H14A	0.96
N1—C14	1.472 (3)	C14—H14B	0.96
N2—C7	1.367 (3)	C14—H14C	0.96
N2—C6	1.424 (3)	C15—C16	1.401 (3)
N2—C10	1.470 (3)	C15—C20	1.411 (3)
N3—C23	1.371 (3)	C16—C17	1.381 (4)
N3—C15	1.421 (3)	C16—H16	0.93
N3—C28	1.476 (3)	C17—C18	1.385 (4)
N4—C21	1.360 (3)	C17—H17	0.93
N4—C20	1.430 (3)	C18—C19	1.383 (4)
N4—C24	1.473 (3)	C18—H18	0.93
C1—C6	1.406 (3)	C19—C20	1.397 (4)
C1—C2	1.406 (4)	C19—H19	0.93
C2—C3	1.379 (4)	C21—C22	1.535 (3)
C2—H2	0.93	C22—C25	1.526 (4)
C3—C4	1.385 (4)	C22—C23	1.530 (3)
C3—H3	0.93	C22—H22	0.98
C4—C5	1.388 (4)	C24—H24A	0.96
C4—H4	0.93	C24—H24B	0.96
C5—C6	1.393 (4)	C24—H24C	0.96
C5—H5	0.93	C25—C26	1.468 (4)
C7—C8	1.528 (3)	C25—H25A	0.97
C8—C9	1.526 (3)	C25—H25B	0.97
C8—C11	1.534 (3)	C26—C27	1.177 (4)
C8—H8	0.98	C27—H27	0.93 (3)
C10—H10A	0.96	C28—H28A	0.96
C10—H10B	0.96	C28—H28B	0.96
C10—H10C	0.96	C28—H28C	0.96
C9—N1—C1	123.77 (19)	N1—C14—H14A	109.5
C9—N1—C14	117.4 (2)	N1—C14—H14B	109.5
C1—N1—C14	118.8 (2)	H14A—C14—H14B	109.5
C7—N2—C6	124.09 (19)	N1—C14—H14C	109.5
C7—N2—C10	116.2 (2)	H14A—C14—H14C	109.5
C6—N2—C10	118.88 (19)	H14B—C14—H14C	109.5
C23—N3—C15	122.7 (2)	C16—C15—C20	118.8 (2)
C23—N3—C28	117.8 (2)	C16—C15—N3	119.6 (2)
C15—N3—C28	118.7 (2)	C20—C15—N3	121.6 (2)
C21—N4—C20	122.6 (2)	C17—C16—C15	120.7 (2)
C21—N4—C24	117.4 (2)	C17—C16—H16	119.7
C20—N4—C24	118.6 (2)	C15—C16—H16	119.7
C6—C1—C2	119.1 (2)	C16—C17—C18	120.3 (3)
C6—C1—N1	122.0 (2)	C16—C17—H17	119.9
C2—C1—N1	118.8 (2)	C18—C17—H17	119.9
C3—C2—C1	120.8 (2)	C17—C18—C19	120.1 (3)
C3—C2—H2	119.6	C17—C18—H18	120.0
C1—C2—H2	119.6	C19—C18—H18	120.0
C2—C3—C4	120.0 (3)	C18—C19—C20	120.5 (3)
C2—C3—H3	120.0	C18—C19—H19	119.8

supplementary materials

C4—C3—H3	120.0	C20—C19—H19	119.8
C3—C4—C5	119.8 (3)	C19—C20—C15	119.6 (2)
C3—C4—H4	120.1	C19—C20—N4	119.1 (2)
C5—C4—H4	120.1	C15—C20—N4	121.3 (2)
C4—C5—C6	121.1 (3)	O4—C21—N4	122.7 (2)
C4—C5—H5	119.4	O4—C21—C22	122.1 (2)
C6—C5—H5	119.4	N4—C21—C22	115.1 (2)
C5—C6—C1	119.0 (2)	C25—C22—C23	111.7 (2)
C5—C6—N2	118.9 (2)	C25—C22—C21	110.40 (19)
C1—C6—N2	122.1 (2)	C23—C22—C21	107.2 (2)
O2—C7—N2	122.0 (2)	C25—C22—H22	109.2
O2—C7—C8	122.3 (2)	C23—C22—H22	109.2
N2—C7—C8	115.7 (2)	C21—C22—H22	109.2
C9—C8—C7	105.00 (19)	O3—C23—N3	121.9 (2)
C9—C8—C11	111.0 (2)	O3—C23—C22	121.6 (2)
C7—C8—C11	110.36 (19)	N3—C23—C22	116.4 (2)
C9—C8—H8	110.1	N4—C24—H24A	109.5
C7—C8—H8	110.1	N4—C24—H24B	109.5
C11—C8—H8	110.1	H24A—C24—H24B	109.5
O1—C9—N1	121.8 (2)	N4—C24—H24C	109.5
O1—C9—C8	122.4 (2)	H24A—C24—H24C	109.5
N1—C9—C8	115.7 (2)	H24B—C24—H24C	109.5
N2—C10—H10A	109.5	C26—C25—C22	112.3 (2)
N2—C10—H10B	109.5	C26—C25—H25A	109.1
H10A—C10—H10B	109.5	C22—C25—H25A	109.1
N2—C10—H10C	109.5	C26—C25—H25B	109.1
H10A—C10—H10C	109.5	C22—C25—H25B	109.1
H10B—C10—H10C	109.5	H25A—C25—H25B	107.9
C12—C11—C8	112.7 (2)	C27—C26—C25	178.0 (3)
C12—C11—H11A	109.1	C26—C27—H27	176 (3)
C8—C11—H11A	109.1	N3—C28—H28A	109.5
C12—C11—H11B	109.1	N3—C28—H28B	109.5
C8—C11—H11B	109.1	H28A—C28—H28B	109.5
H11A—C11—H11B	107.8	N3—C28—H28C	109.5
C13—C12—C11	178.1 (3)	H28A—C28—H28C	109.5
C12—C13—H13	180 (3)	H28B—C28—H28C	109.5
C9—N1—C1—C6	44.7 (3)	C23—N3—C15—C16	131.1 (3)
C14—N1—C1—C6	-133.7 (3)	C28—N3—C15—C16	-38.5 (3)
C9—N1—C1—C2	-137.7 (2)	C23—N3—C15—C20	-48.5 (3)
C14—N1—C1—C2	43.9 (3)	C28—N3—C15—C20	142.0 (3)
C6—C1—C2—C3	2.9 (4)	C20—C15—C16—C17	2.7 (4)
N1—C1—C2—C3	-174.9 (2)	N3—C15—C16—C17	-176.9 (3)
C1—C2—C3—C4	-0.3 (4)	C15—C16—C17—C18	-1.5 (4)
C2—C3—C4—C5	-1.9 (4)	C16—C17—C18—C19	-0.4 (4)
C3—C4—C5—C6	1.6 (4)	C17—C18—C19—C20	1.1 (4)
C4—C5—C6—C1	1.0 (4)	C18—C19—C20—C15	0.0 (4)
C4—C5—C6—N2	178.2 (2)	C18—C19—C20—N4	-179.5 (3)
C2—C1—C6—C5	-3.2 (3)	C16—C15—C20—C19	-1.9 (4)
N1—C1—C6—C5	174.4 (2)	N3—C15—C20—C19	177.7 (2)

C2—C1—C6—N2	179.8 (2)	C16—C15—C20—N4	177.6 (2)
N1—C1—C6—N2	-2.6 (3)	N3—C15—C20—N4	-2.8 (4)
C7—N2—C6—C5	133.8 (2)	C21—N4—C20—C19	-124.8 (3)
C10—N2—C6—C5	-35.1 (3)	C24—N4—C20—C19	41.7 (3)
C7—N2—C6—C1	-49.2 (3)	C21—N4—C20—C15	55.7 (3)
C10—N2—C6—C1	141.9 (2)	C24—N4—C20—C15	-137.8 (2)
C6—N2—C7—O2	-173.0 (2)	C20—N4—C21—O4	171.5 (2)
C10—N2—C7—O2	-3.8 (3)	C24—N4—C21—O4	4.8 (4)
C6—N2—C7—C8	9.9 (3)	C20—N4—C21—C22	-11.6 (3)
C10—N2—C7—C8	179.0 (2)	C24—N4—C21—C22	-178.2 (2)
O2—C7—C8—C9	-109.3 (3)	O4—C21—C22—C25	-12.3 (3)
N2—C7—C8—C9	67.9 (3)	N4—C21—C22—C25	170.7 (2)
O2—C7—C8—C11	10.4 (3)	O4—C21—C22—C23	109.5 (3)
N2—C7—C8—C11	-172.4 (2)	N4—C21—C22—C23	-67.4 (3)
C1—N1—C9—O1	-173.1 (2)	C15—N3—C23—O3	-177.0 (2)
C14—N1—C9—O1	5.3 (3)	C28—N3—C23—O3	-7.4 (4)
C1—N1—C9—C8	4.2 (3)	C15—N3—C23—C22	6.0 (3)
C14—N1—C9—C8	-177.4 (2)	C28—N3—C23—C22	175.6 (2)
C7—C8—C9—O1	100.1 (2)	C25—C22—C23—O3	14.7 (3)
C11—C8—C9—O1	-19.1 (3)	C21—C22—C23—O3	-106.3 (3)
C7—C8—C9—N1	-77.2 (2)	C25—C22—C23—N3	-168.3 (2)
C11—C8—C9—N1	163.55 (19)	C21—C22—C23—N3	70.7 (3)
C9—C8—C11—C12	-63.5 (3)	C23—C22—C25—C26	72.0 (3)
C7—C8—C11—C12	-179.5 (2)	C21—C22—C25—C26	-168.9 (2)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C13—H13···O4	0.93 (3)	2.44 (3)	3.366 (3)	172 (3)
C8—H8···O2 ⁱ	0.98	2.56	3.536 (3)	175
C19—H19···O1 ⁱ	0.93	2.49	3.374 (3)	160

Symmetry codes: (i) $x, -y+1, z+1/2$.

supplementary materials

Fig. 1

