

## 4,5-Diaza-9H-fluoren-9-imine

Hui Cang,<sup>a</sup> Jin-Xiu Ji,<sup>b</sup> Si-Qing Wang<sup>c</sup> and Jin-Tang Wang<sup>c\*</sup>

<sup>a</sup>Chemical and Biological Engineering College, Yancheng Institute of Technology, Yancheng 224051, People's Republic of China, <sup>b</sup>Research & Development Center, Sinochem Jiangsu Corporation, Nanjing 210005, People's Republic of China, and <sup>c</sup>Department of Applied Chemistry, College of Science, Nanjing University of Technology, Nanjing 210009, People's Republic of China  
Correspondence e-mail: canghui@ycit.edu.cn


Received 14 May 2010; accepted 29 May 2010

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.066;  $wR$  factor = 0.194; data-to-parameter ratio = 11.8.

In the title compound,  $\text{C}_{11}\text{H}_7\text{N}_3$ , the diazafluorene rings are almost coplanar with an r.m.s. deviation of 0.0160 Å. In the crystal structure,  $\text{C}-\text{H}\cdots\text{N}$  hydrogen bonds link molecules into sheets parallel to the  $ab$  plane. Molecules are also stacked regularly along the  $c$  axis by a variety of  $\pi-\pi$  interactions with centroid-centroid distances in the range 3.527 (2)–3.908 (2) Å.

### Related literature

For the use of the title compound in synthesizing complexes with interesting photochemical properties and for the synthesis, see: Wang & Rillema (1997). For reference bond-length data, see: Allen *et al.* (1987).



### Experimental

#### Crystal data

 $\text{C}_{11}\text{H}_7\text{N}_3$ 
 $M_r = 181.20$ 

Monoclinic,  $P2_1/c$   
 $a = 10.008$  (2) Å  
 $b = 12.407$  (3) Å  
 $c = 6.8140$  (14) Å  
 $\beta = 99.74$  (3)°  
 $V = 833.9$  (3) Å<sup>3</sup>

$Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.09$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.30 \times 0.10 \times 0.10$  mm

#### Data collection

Enraf–Nonius CAD-4 diffractometer  
 Absorption correction:  $\psi$  scan (North *et al.*, 1968)  
 $T_{\min} = 0.973$ ,  $T_{\max} = 0.991$   
 1638 measured reflections

1503 independent reflections  
 1010 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.022$   
 3 standard reflections every 200 reflections  
 intensity decay: none

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.066$   
 $wR(F^2) = 0.194$   
 $S = 1.06$   
 1503 reflections  
 127 parameters

40 restraints  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.28$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.21$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C}11-\text{H}11A\cdots\text{N}3^i$	0.93	2.45	3.382 (4)	178
$\text{C}4-\text{H}4A\cdots\text{N}1^ii$	0.93	2.69	3.536 (4)	152

Symmetry codes: (i)  $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x + 2, -y + 1, -z + 1$ .

Data collection: *CAD-4 Software* (Enraf–Nonius, 1985); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Center of Testing and Analysis, Nanjing University, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5007).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.  
 Enraf–Nonius (1985). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.  
 Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.  
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Wang, Y. X. & Rillema, D. P. (1997). *Tetrahedron*, **37**, 12377–12390.

**supplementary materials**

*Acta Cryst.* (2010). E66, o1802 [ doi:10.1107/S1600536810020441 ]

## 4,5-Diaza-9*H*-fluoren-9-imine

H. Cang, J.-X. Ji, S.-Q. Wang and J.-T. Wang

### Comment

4,5-diazafluorene-9-imine is one of the important ligands, being utilized to synthesize complexes with interesting photochemical properties (Wang & Rillema, 1997). Here we report the crystal structure of the title compound, (I).

The molecular structure of (I) is shown in Fig. 1, and the selected geometric parameters are given in Table 1. The bond lengths and angles (Table 1) are within normal ranges (Allen *et al.*, 1987). The diazafluorene rings are almost coplanar with an rms deviation 0.0160 Å.

In the crystal structure C—H...N hydrogen bonds link molecules into sheets parallel to the *ab* plane, Table 1. An extensive system of  $\pi$ - $\pi$  contacts stacks molecules in an obverse fashion down the *c* axis, Fig. 2, with Cg1...Cg1 = 3.876 (2) %Å, Cg2...Cg2 = 3.572 (2) %Å, Cg(3)...Cg3 = 3.908 (2) and Cg1...Cg2 3.776 (2) Å and 3.863 (2) Å. Symmetry operations *x*, 1/2-*y*, 1/2+*z*, and *x*, 1/2-*y*, -1/2+*z*; Cg1, Cg2 and Cg3 are the centroids of the C2,C3,C5,C7,C8; N2,C1,C2,C3,C4,C5 and N3,C6,C7,C10,C11 rings, respectively.

### Experimental

The title compound was synthesized by a method reported in literature (Wang & Rillema, 1997). Crystals were obtained by dissolving the compound (2.0 g, 11.0 mmol) in ethyl acetate(50 ml), and evaporating the solvent slowly at room temperature for about 5 d.

### Refinement

H atoms were positioned geometrically, with N—H = 0.75 and C—H = 0.93Å for aromatic C—H, and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C/N})$ , where  $x = 1.2$  for aromatic H and  $x = 1.5$  for the N—H.

### Figures




Fig. 1. A drawing of the title molecular structure, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. A packing diagram for (I). Hydrogen bonds are drawn as dashed lines.

## 4,5-Diaza-9H-fluoren-9-imine

### Crystal data

$C_{11}H_7N_3$

$M_r = 181.20$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 10.008$  (2) Å

$b = 12.407$  (3) Å

$c = 6.8140$  (14) Å

$\beta = 99.74$  (3)°

$V = 833.9$  (3) Å<sup>3</sup>

$Z = 4$

$F(000) = 376$

$D_x = 1.443$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 25 reflections

$\theta = 10\text{--}13^\circ$

$\mu = 0.09$  mm<sup>-1</sup>

$T = 293$  K

Block, colourless

$0.30 \times 0.10 \times 0.10$  mm

### Data collection

Enraf-Nonius CAD-4  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

$\omega/2\theta$  scans

Absorption correction:  $\psi$  scan  
(North *et al.*, 1968)

$T_{\min} = 0.973$ ,  $T_{\max} = 0.991$

1638 measured reflections

1503 independent reflections

1010 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\max} = 25.3^\circ$ ,  $\theta_{\min} = 2.1^\circ$

$h = -11 \rightarrow 11$

$k = 0 \rightarrow 14$

$l = 0 \rightarrow 8$

3 standard reflections every 200 reflections

intensity decay: none

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.066$

$wR(F^2) = 0.194$

$S = 1.06$

1503 reflections

127 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0948P)^2 + 0.5792P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.28$  e Å<sup>-3</sup>

40 restraints

$$\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.8062 (3)	0.5097 (2)	0.3404 (5)	0.0530 (8)
H1	0.7491	0.5497	0.3246	0.079*
N2	0.7922 (3)	0.1271 (2)	0.3022 (4)	0.0488 (8)
N3	0.5012 (3)	0.2137 (2)	0.2363 (4)	0.0455 (7)
C1	0.9253 (4)	0.1124 (3)	0.3376 (5)	0.0540 (10)
H1A	0.9567	0.0418	0.3391	0.065*
C2	0.7547 (3)	0.2304 (2)	0.3049 (5)	0.0410 (8)
C3	0.8433 (3)	0.3171 (2)	0.3340 (4)	0.0389 (8)
C4	0.9804 (4)	0.2998 (3)	0.3749 (5)	0.0508 (9)
H4A	1.0425	0.3559	0.4027	0.061*
C5	1.0202 (4)	0.1925 (3)	0.3720 (6)	0.0545 (10)
H5A	1.1119	0.1751	0.3935	0.065*
C6	0.6122 (3)	0.2709 (2)	0.2711 (4)	0.0360 (7)
C7	0.6173 (3)	0.3850 (2)	0.2811 (5)	0.0422 (8)
C8	0.7604 (3)	0.4188 (2)	0.3239 (5)	0.0441 (8)
C9	0.3856 (3)	0.2703 (3)	0.2096 (5)	0.0496 (9)
H9A	0.3044	0.2324	0.1837	0.060*
C10	0.3791 (3)	0.3820 (3)	0.2181 (6)	0.0546 (10)
H10A	0.2955	0.4167	0.2002	0.066*
C11	0.4973 (3)	0.4412 (3)	0.2531 (5)	0.0505 (9)
H11A	0.4959	0.5161	0.2576	0.061*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0453 (17)	0.0273 (15)	0.083 (2)	-0.0112 (13)	0.0014 (15)	0.0041 (14)
N2	0.0498 (18)	0.0401 (16)	0.0546 (17)	0.0120 (14)	0.0033 (13)	0.0013 (13)
N3	0.0466 (17)	0.0278 (14)	0.0614 (17)	-0.0041 (12)	0.0071 (13)	-0.0015 (12)
C1	0.058 (2)	0.041 (2)	0.063 (2)	0.0158 (18)	0.0088 (17)	0.0032 (17)
C2	0.0445 (18)	0.0317 (17)	0.0454 (18)	0.0036 (14)	0.0032 (14)	-0.0001 (14)
C3	0.0430 (18)	0.0370 (17)	0.0380 (15)	-0.0060 (14)	0.0105 (13)	0.0004 (13)

## supplementary materials

---

C4	0.045 (2)	0.046 (2)	0.059 (2)	-0.0079 (16)	0.0011 (16)	0.0065 (16)
C5	0.0399 (19)	0.065 (3)	0.058 (2)	0.0084 (18)	0.0070 (15)	0.0018 (18)
C6	0.0387 (16)	0.0285 (16)	0.0390 (16)	-0.0005 (13)	0.0011 (12)	-0.0013 (12)
C7	0.0478 (19)	0.0246 (16)	0.0516 (18)	-0.0031 (14)	0.0013 (14)	0.0001 (13)
C8	0.0497 (19)	0.0282 (16)	0.0536 (19)	-0.0097 (15)	0.0062 (15)	-0.0024 (14)
C9	0.0339 (18)	0.045 (2)	0.068 (2)	-0.0037 (15)	0.0030 (15)	-0.0008 (17)
C10	0.045 (2)	0.040 (2)	0.079 (3)	0.0097 (17)	0.0094 (17)	-0.0004 (18)
C11	0.056 (2)	0.0250 (17)	0.069 (2)	0.0054 (16)	0.0049 (17)	0.0010 (16)

### *Geometric parameters (Å, °)*

N1—C8	1.216 (4)	C4—C5	1.391 (5)
N1—H1	0.7500	C4—H4A	0.9300
N2—C1	1.325 (4)	C5—H5A	0.9300
N2—C2	1.336 (4)	C6—C7	1.417 (4)
N3—C6	1.305 (4)	C7—C11	1.373 (4)
N3—C9	1.340 (4)	C7—C8	1.473 (4)
C1—C5	1.367 (5)	C9—C10	1.389 (5)
C1—H1A	0.9300	C9—H9A	0.9300
C2—C3	1.386 (4)	C10—C11	1.379 (5)
C2—C6	1.493 (4)	C10—H10A	0.9300
C3—C4	1.371 (5)	C11—H11A	0.9300
C3—C8	1.505 (4)		
C8—N1—H1	109.5	N3—C6—C7	125.1 (3)
C1—N2—C2	113.9 (3)	N3—C6—C2	127.3 (3)
C6—N3—C9	115.4 (3)	C7—C6—C2	107.6 (3)
N2—C1—C5	125.4 (3)	C11—C7—C6	118.5 (3)
N2—C1—H1A	117.3	C11—C7—C8	132.9 (3)
C5—C1—H1A	117.3	C6—C7—C8	108.6 (3)
N2—C2—C3	124.9 (3)	N1—C8—C7	128.4 (3)
N2—C2—C6	125.7 (3)	N1—C8—C3	125.3 (3)
C3—C2—C6	109.4 (3)	C7—C8—C3	106.3 (2)
C4—C3—C2	120.1 (3)	N3—C9—C10	124.3 (3)
C4—C3—C8	131.7 (3)	N3—C9—H9A	117.9
C2—C3—C8	108.0 (3)	C10—C9—H9A	117.9
C3—C4—C5	115.2 (3)	C11—C10—C9	119.5 (3)
C3—C4—H4A	122.4	C11—C10—H10A	120.2
C5—C4—H4A	122.4	C9—C10—H10A	120.2
C1—C5—C4	120.4 (3)	C7—C11—C10	117.3 (3)
C1—C5—H5A	119.8	C7—C11—H11A	121.3
C4—C5—H5A	119.8	C10—C11—H11A	121.3
C2—N2—C1—C5	-0.8 (5)	N3—C6—C7—C11	0.2 (5)
C1—N2—C2—C3	2.3 (5)	C2—C6—C7—C11	179.6 (3)
C1—N2—C2—C6	-179.7 (3)	N3—C6—C7—C8	179.7 (3)
N2—C2—C3—C4	-4.0 (5)	C2—C6—C7—C8	-0.9 (4)
C6—C2—C3—C4	177.8 (3)	C11—C7—C8—N1	-1.9 (6)
N2—C2—C3—C8	179.3 (3)	C6—C7—C8—N1	178.7 (4)
C6—C2—C3—C8	1.1 (3)	C11—C7—C8—C3	-179.1 (3)
C2—C3—C4—C5	3.7 (5)	C6—C7—C8—C3	1.5 (4)

C8—C3—C4—C5	179.5 (3)	C4—C3—C8—N1	5.0 (6)
N2—C1—C5—C4	1.0 (6)	C2—C3—C8—N1	-178.9 (3)
C3—C4—C5—C1	-2.4 (5)	C4—C3—C8—C7	-177.8 (3)
C9—N3—C6—C7	-0.2 (5)	C2—C3—C8—C7	-1.6 (3)
C9—N3—C6—C2	-179.5 (3)	C6—N3—C9—C10	-0.4 (5)
N2—C2—C6—N3	1.0 (5)	N3—C9—C10—C11	1.0 (6)
C3—C2—C6—N3	179.3 (3)	C6—C7—C11—C10	0.4 (5)
N2—C2—C6—C7	-178.4 (3)	C8—C7—C11—C10	-179.0 (3)
C3—C2—C6—C7	-0.1 (3)	C9—C10—C11—C7	-0.9 (5)

*Hydrogen-bond geometry* ( $\text{\AA}$ ,  $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C11—H11A $\cdots$ N3 <sup>i</sup>	0.93	2.45	3.382 (4)	178
C4—H4A $\cdots$ N1 <sup>ii</sup>	0.93	2.69	3.536 (4)	152

Symmetry codes: (i)  $-x+1, y+1/2, -z+1/2$ ; (ii)  $-x+2, -y+1, -z+1$ .

Fig. 1






Fig. 2

