

Methyl 5-chloro-2-[*N*-(3-ethoxy-carbonylpropyl)-4-methylbenzene-sulfonamido]benzoate

Bin Wang,^{a,b} Ru Jia,^a Ya-Bin Shi,^{a,b} Fei-Fei He^{a,b} and Hai-Bo Wang^{a*}

^aCollege of Food Science and Light Industry, Nanjing University of Technology, Xinmofan Road No.5 Nanjing, Nanjing 210009, People's Republic of China, and

^bCollege of Science, Nanjing University of Technology, Xinmofan Road No.5

Nanjing, Nanjing 210009, People's Republic of China

Correspondence e-mail: wanghaibo@njut.edu.cn


Received 7 June 2010; accepted 19 June 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.009\text{ \AA}$; R factor = 0.063; wR factor = 0.114; data-to-parameter ratio = 15.2.

In the title compound, $\text{C}_{21}\text{H}_{24}\text{ClNO}_6\text{S}$, the benzene rings are oriented at a dihedral angles of $41.6(2)^\circ$. In the crystal structure, weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ interactions link the molecules.

Related literature

For the preparation of the title compound, see: Kondo *et al.* (1999). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{24}\text{ClNO}_6\text{S}$

$M_r = 453.92$

Orthorhombic, $P2_12_12_1$

$a = 9.1480(18)\text{ \AA}$

$b = 10.742(2)\text{ \AA}$

$c = 23.258(5)\text{ \AA}$

$V = 2285.5(8)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.29\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.30 \times 0.10 \times 0.05\text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer

Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.917$, $T_{\max} = 0.985$

4623 measured reflections

4130 independent reflections
2134 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.072$

3 standard reflections every 200 reflections
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.063$
 $wR(F^2) = 0.114$

$S = 0.92$

4130 reflections
271 parameters
H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.19\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.21\text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
1748 Friedel pairs
Flack parameter: 0.03 (12)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C11—H11A…O3 ⁱ	0.93	2.55	3.275 (6)	135
C14—H14A…O2 ⁱⁱ	0.96	2.60	3.338 (7)	134
C17—H17A…O2 ⁱⁱⁱ	0.93	2.59	3.414 (8)	148

Symmetry codes: (i) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, y + 1, z$; (iii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1989); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2221).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Enraf–Nonius (1989). *CAD-4 EXPRESS*. Enraf–Nonius, Delft. The Netherlands.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Kondo, K., Ogawa, H., Yamashita, H. & Miyamoto, H. (1999). *Bioorg. Med. Chem.* **7**, 1743–1754.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A24*, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o1819 [doi:10.1107/S1600536810023792]

Methyl 5-chloro-2-[*N*-(3-ethoxycarbonylpropyl)-4-methylbenzenesulfonamido]benzoate

B. Wang, R. Jia, Y.-B. Shi, F.-F. He and H.-B. Wang

Comment

Quinolines are a major class of alkaloids and play an important role in the fields of natural products and medicinal chemistry. The title compound, (I), is a useful intermediate. (Kondo *et al.*, 1999). In the molecule of the title compound (Fig. 1), the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. The intramolecular C-H···O hydrogen bond (Table 1) results in the formation of a five-membered ring C (C6/H6A/O4/S/N). Rings A(C7-C12) and B(C15-C20) are planar with maximum deviations of 0.011 (4) Å for C9 and -0.021 (5) Å for C16, respectively. The dihedral angle between these rings is 41.6 (2) °. In the crystal structure, intermolecular weak C-H..O interactions link the molecules to form a stable structure.

Experimental

The title compound, (I) was prepared by the literature method (Kondo *et al.*, 1999). Crystals suitable for X-ray analysis were obtained by slow evaporation of an methanol solution.

Refinement

H atoms were positioned geometrically, with C-H = 0.93, 0.98 and 0.96 Å for aromatic, methine and methyl H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N})$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

Methyl 5-chloro-2-[*N*-(3-ethoxycarbonylpropyl)- 4-methylbenzenesulfonamido]benzoate

Crystal data

$\text{C}_{21}\text{H}_{24}\text{ClNO}_6\text{S}$	$D_x = 1.319 \text{ Mg m}^{-3}$
$M_r = 453.92$	Melting point: 353 K
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 25 reflections
$a = 9.1480 (18) \text{ \AA}$	$\theta = 8\text{--}12^\circ$
$b = 10.742 (2) \text{ \AA}$	$\mu = 0.29 \text{ mm}^{-1}$
$c = 23.258 (5) \text{ \AA}$	$T = 296 \text{ K}$

supplementary materials

$V = 2285.5 (8) \text{ \AA}^3$ Needle, colorless
 $Z = 4$ $0.30 \times 0.10 \times 0.05 \text{ mm}$
 $F(000) = 952$

Data collection

Enraf–Nonius CAD-4 diffractometer
Radiation source: fine-focus sealed tube
graphite
 $\omega/2\theta$ scans
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.917$, $T_{\max} = 0.985$
4623 measured reflections
4130 independent reflections
2134 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.072$
 $\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 1.8^\circ$
 $h = -10 \rightarrow 0$
 $k = -12 \rightarrow 0$
 $l = -27 \rightarrow 27$
3 standard reflections every 200 reflections
intensity decay: 1%

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.063$ H-atom parameters constrained
 $wR(F^2) = 0.114$ $w = 1/[\sigma^2(F_o^2) + (0.025P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $S = 0.92$ $(\Delta/\sigma)_{\max} < 0.001$
4130 reflections $\Delta\rho_{\max} = 0.19 \text{ e \AA}^{-3}$
271 parameters $\Delta\rho_{\min} = -0.21 \text{ e \AA}^{-3}$
0 restraints Absolute structure: Flack (1983), 1748 Friedel pairs
Primary atom site location: structure-invariant direct Flack parameter: 0.03 (12)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S	0.32206 (15)	0.97963 (12)	0.21916 (6)	0.0523 (4)

Cl	-0.34814 (14)	1.01035 (14)	0.35104 (7)	0.0753 (5)
O1	0.5372 (6)	0.6826 (5)	0.4389 (3)	0.130 (2)
O2	0.3409 (6)	0.5726 (4)	0.4257 (2)	0.1119 (19)
O3	0.2927 (4)	1.1049 (3)	0.23835 (15)	0.0605 (11)
O4	0.4681 (4)	0.9398 (3)	0.21051 (16)	0.0668 (12)
O5	0.1790 (4)	1.2067 (3)	0.35132 (15)	0.0537 (9)
O6	0.3165 (4)	1.0402 (3)	0.37290 (16)	0.0651 (11)
N	0.2482 (4)	0.8844 (3)	0.26778 (18)	0.0459 (12)
C1	0.6798 (12)	0.6647 (8)	0.5254 (4)	0.165 (4)
H1A	0.6820	0.6204	0.5613	0.247*
H1B	0.6714	0.7524	0.5328	0.247*
H1C	0.7684	0.6487	0.5045	0.247*
C2	0.5609 (11)	0.6252 (9)	0.4933 (4)	0.145 (4)
H2A	0.4737	0.6372	0.5163	0.174*
H2B	0.5713	0.5363	0.4869	0.174*
C3	0.4202 (8)	0.6527 (6)	0.4104 (3)	0.077 (2)
C4	0.4113 (6)	0.7278 (5)	0.3548 (3)	0.0671 (17)
H4B	0.3982	0.8152	0.3641	0.080*
H4C	0.5023	0.7193	0.3338	0.080*
C5	0.2883 (6)	0.6854 (5)	0.3181 (2)	0.0570 (16)
H5A	0.1971	0.6994	0.3384	0.068*
H5B	0.2978	0.5966	0.3116	0.068*
C6	0.2816 (6)	0.7504 (4)	0.2610 (2)	0.0553 (16)
H6A	0.3746	0.7411	0.2415	0.066*
H6B	0.2069	0.7118	0.2374	0.066*
C7	0.1045 (6)	0.9196 (4)	0.2868 (2)	0.0452 (13)
C8	0.0844 (5)	1.0121 (4)	0.3272 (2)	0.0412 (12)
C9	-0.0552 (5)	1.0403 (5)	0.3460 (2)	0.0543 (15)
H9A	-0.0678	1.1043	0.3725	0.065*
C10	-0.1762 (6)	0.9764 (4)	0.3267 (2)	0.0549 (14)
C11	-0.1565 (6)	0.8815 (5)	0.2868 (3)	0.0610 (16)
H11A	-0.2360	0.8361	0.2734	0.073*
C12	-0.0171 (6)	0.8557 (5)	0.2676 (3)	0.0562 (16)
H12A	-0.0045	0.7927	0.2405	0.067*
C13	0.2085 (6)	1.0834 (4)	0.3535 (2)	0.0476 (13)
C14	0.2962 (6)	1.2856 (5)	0.3736 (3)	0.080 (2)
H14A	0.2676	1.3714	0.3709	0.120*
H14B	0.3834	1.2724	0.3514	0.120*
H14C	0.3147	1.2648	0.4131	0.120*
C15	0.2247 (6)	0.9560 (5)	0.1556 (2)	0.0558 (15)
C16	0.0987 (6)	1.0246 (6)	0.1445 (3)	0.0737 (18)
H16A	0.0677	1.0842	0.1708	0.088*
C17	0.0199 (7)	1.0052 (7)	0.0952 (3)	0.082 (2)
H17A	-0.0607	1.0549	0.0870	0.098*
C18	0.0601 (7)	0.9126 (7)	0.0581 (3)	0.074 (2)
C19	0.1794 (8)	0.8414 (6)	0.0698 (3)	0.078 (2)
H19A	0.2045	0.7768	0.0451	0.094*
C20	0.2628 (7)	0.8637 (6)	0.1175 (2)	0.0659 (18)
H20A	0.3460	0.8160	0.1242	0.079*

supplementary materials

C21	-0.0298 (8)	0.8910 (8)	0.0048 (3)	0.132 (3)
H21A	-0.1074	0.9511	0.0032	0.198*
H21B	-0.0705	0.8087	0.0059	0.198*
H21C	0.0311	0.8995	-0.0285	0.198*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S	0.0493 (8)	0.0417 (7)	0.0659 (10)	-0.0006 (8)	0.0099 (8)	0.0019 (8)
Cl	0.0481 (8)	0.0653 (10)	0.1124 (13)	-0.0118 (8)	0.0152 (9)	-0.0006 (10)
O1	0.139 (5)	0.116 (4)	0.134 (5)	-0.032 (4)	-0.076 (4)	0.041 (4)
O2	0.134 (5)	0.089 (4)	0.113 (4)	-0.045 (4)	-0.014 (4)	0.026 (3)
O3	0.084 (3)	0.0296 (18)	0.068 (3)	0.002 (2)	0.012 (2)	-0.0045 (17)
O4	0.054 (2)	0.063 (3)	0.083 (3)	-0.006 (2)	0.017 (2)	-0.006 (2)
O5	0.046 (2)	0.0352 (18)	0.080 (3)	-0.0061 (18)	-0.001 (2)	-0.0094 (19)
O6	0.056 (2)	0.051 (2)	0.088 (3)	0.009 (2)	-0.016 (2)	-0.007 (2)
N	0.055 (3)	0.031 (2)	0.051 (3)	0.000 (2)	0.014 (2)	0.001 (2)
C1	0.192 (12)	0.155 (9)	0.148 (9)	-0.042 (9)	-0.011 (9)	-0.002 (7)
C2	0.163 (10)	0.136 (8)	0.138 (9)	-0.013 (8)	-0.045 (8)	0.033 (7)
C3	0.087 (6)	0.056 (4)	0.087 (5)	-0.010 (4)	-0.028 (5)	-0.004 (4)
C4	0.070 (4)	0.041 (3)	0.090 (5)	0.008 (3)	-0.011 (4)	0.007 (4)
C5	0.057 (4)	0.039 (3)	0.075 (4)	0.002 (3)	-0.005 (3)	0.004 (3)
C6	0.068 (4)	0.029 (3)	0.070 (4)	0.013 (3)	0.007 (3)	-0.001 (3)
C7	0.046 (3)	0.033 (3)	0.056 (4)	0.000 (3)	0.005 (3)	0.007 (3)
C8	0.034 (3)	0.030 (3)	0.060 (3)	-0.008 (3)	0.003 (2)	0.004 (3)
C9	0.046 (3)	0.040 (3)	0.077 (4)	-0.004 (3)	-0.006 (3)	0.006 (3)
C10	0.050 (3)	0.039 (3)	0.076 (4)	-0.007 (3)	0.002 (3)	0.003 (3)
C11	0.052 (4)	0.051 (3)	0.080 (4)	0.000 (3)	-0.011 (4)	0.003 (3)
C12	0.057 (4)	0.039 (3)	0.072 (5)	-0.005 (3)	-0.004 (3)	-0.002 (3)
C13	0.053 (4)	0.040 (3)	0.050 (3)	0.001 (3)	0.000 (3)	-0.002 (3)
C14	0.068 (4)	0.050 (3)	0.122 (6)	-0.032 (4)	0.002 (4)	-0.019 (4)
C15	0.059 (4)	0.050 (3)	0.058 (4)	-0.004 (3)	0.010 (3)	0.000 (3)
C16	0.067 (4)	0.082 (5)	0.073 (5)	0.036 (4)	0.004 (4)	-0.009 (4)
C17	0.063 (4)	0.102 (6)	0.081 (5)	0.018 (5)	0.001 (4)	0.017 (5)
C18	0.051 (4)	0.107 (6)	0.065 (5)	-0.003 (4)	0.010 (4)	-0.001 (5)
C19	0.084 (5)	0.082 (5)	0.069 (5)	-0.010 (5)	0.030 (4)	-0.017 (4)
C20	0.081 (5)	0.068 (4)	0.049 (4)	0.029 (4)	0.013 (4)	-0.005 (3)
C21	0.113 (7)	0.215 (10)	0.068 (5)	-0.005 (7)	-0.021 (5)	-0.025 (6)

Geometric parameters (\AA , $^\circ$)

S—O4	1.417 (4)	C7—C8	1.380 (6)
S—O3	1.443 (3)	C7—C12	1.382 (7)
S—N	1.668 (4)	C8—C9	1.383 (6)
S—C15	1.745 (6)	C8—C13	1.500 (7)
Cl—C10	1.711 (5)	C9—C10	1.377 (6)
O1—C3	1.299 (7)	C9—H9A	0.9300
O1—C2	1.424 (9)	C10—C11	1.390 (7)
O2—C3	1.181 (7)	C11—C12	1.380 (7)

O5—C13	1.353 (5)	C11—H11A	0.9300
O5—C14	1.461 (6)	C12—H12A	0.9300
O6—C13	1.181 (6)	C14—H14A	0.9600
N—C7	1.437 (6)	C14—H14B	0.9600
N—C6	1.480 (5)	C14—H14C	0.9600
C1—C2	1.387 (10)	C15—C20	1.375 (7)
C1—H1A	0.9600	C15—C16	1.392 (7)
C1—H1B	0.9600	C16—C17	1.371 (8)
C1—H1C	0.9600	C16—H16A	0.9300
C2—H2A	0.9700	C17—C18	1.367 (8)
C2—H2B	0.9700	C17—H17A	0.9300
C3—C4	1.525 (8)	C18—C19	1.360 (8)
C4—C5	1.484 (7)	C18—C21	1.504 (8)
C4—H4B	0.9700	C19—C20	1.368 (8)
C4—H4C	0.9700	C19—H19A	0.9300
C5—C6	1.501 (7)	C20—H20A	0.9300
C5—H5A	0.9700	C21—H21A	0.9600
C5—H5B	0.9700	C21—H21B	0.9600
C6—H6A	0.9700	C21—H21C	0.9600
C6—H6B	0.9700		
O4—S—O3	120.1 (2)	C7—C8—C13	123.0 (4)
O4—S—N	107.0 (2)	C9—C8—C13	117.3 (5)
O3—S—N	106.7 (2)	C10—C9—C8	122.0 (5)
O4—S—C15	108.5 (3)	C10—C9—H9A	119.0
O3—S—C15	107.6 (2)	C8—C9—H9A	119.0
N—S—C15	106.2 (2)	C9—C10—C11	118.6 (5)
C3—O1—C2	118.1 (7)	C9—C10—Cl	121.6 (4)
C13—O5—C14	114.1 (4)	C11—C10—Cl	119.7 (4)
C7—N—C6	118.5 (4)	C12—C11—C10	118.9 (5)
C7—N—S	114.7 (3)	C12—C11—H11A	120.5
C6—N—S	116.1 (3)	C10—C11—H11A	120.5
C2—C1—H1A	109.5	C11—C12—C7	122.6 (5)
C2—C1—H1B	109.5	C11—C12—H12A	118.7
H1A—C1—H1B	109.5	C7—C12—H12A	118.7
C2—C1—H1C	109.5	O6—C13—O5	124.5 (5)
H1A—C1—H1C	109.5	O6—C13—C8	126.0 (5)
H1B—C1—H1C	109.5	O5—C13—C8	109.5 (5)
C1—C2—O1	117.8 (9)	O5—C14—H14A	109.5
C1—C2—H2A	107.9	O5—C14—H14B	109.5
O1—C2—H2A	107.9	H14A—C14—H14B	109.5
C1—C2—H2B	107.9	O5—C14—H14C	109.5
O1—C2—H2B	107.9	H14A—C14—H14C	109.5
H2A—C2—H2B	107.2	H14B—C14—H14C	109.5
O2—C3—O1	122.2 (7)	C20—C15—C16	118.2 (6)
O2—C3—C4	127.4 (7)	C20—C15—S	121.4 (5)
O1—C3—C4	110.2 (6)	C16—C15—S	120.1 (5)
C5—C4—C3	111.5 (5)	C17—C16—C15	120.6 (6)
C5—C4—H4B	109.3	C17—C16—H16A	119.7
C3—C4—H4B	109.3	C15—C16—H16A	119.7

supplementary materials

C5—C4—H4C	109.3	C18—C17—C16	119.8 (6)
C3—C4—H4C	109.3	C18—C17—H17A	120.1
H4B—C4—H4C	108.0	C16—C17—H17A	120.1
C4—C5—C6	113.4 (5)	C19—C18—C17	119.9 (7)
C4—C5—H5A	108.9	C19—C18—C21	121.1 (7)
C6—C5—H5A	108.9	C17—C18—C21	119.0 (7)
C4—C5—H5B	108.9	C18—C19—C20	120.8 (6)
C6—C5—H5B	108.9	C18—C19—H19A	119.6
H5A—C5—H5B	107.7	C20—C19—H19A	119.6
N—C6—C5	111.5 (4)	C19—C20—C15	120.5 (6)
N—C6—H6A	109.3	C19—C20—H20A	119.8
C5—C6—H6A	109.3	C15—C20—H20A	119.8
N—C6—H6B	109.3	C18—C21—H21A	109.5
C5—C6—H6B	109.3	C18—C21—H21B	109.5
H6A—C6—H6B	108.0	H21A—C21—H21B	109.5
C8—C7—C12	118.1 (5)	C18—C21—H21C	109.5
C8—C7—N	121.4 (4)	H21A—C21—H21C	109.5
C12—C7—N	120.4 (5)	H21B—C21—H21C	109.5
C7—C8—C9	119.7 (5)		
O4—S—N—C7	173.3 (4)	C9—C10—C11—C12	-0.8 (8)
O3—S—N—C7	43.6 (4)	Cl—C10—C11—C12	179.9 (4)
C15—S—N—C7	-71.0 (4)	C10—C11—C12—C7	0.8 (9)
O4—S—N—C6	-42.5 (4)	C8—C7—C12—C11	0.5 (8)
O3—S—N—C6	-172.2 (4)	N—C7—C12—C11	176.5 (5)
C15—S—N—C6	73.2 (4)	C14—O5—C13—O6	2.0 (8)
C3—O1—C2—C1	-175.0 (9)	C14—O5—C13—C8	-177.4 (4)
C2—O1—C3—O2	-6.6 (12)	C7—C8—C13—O6	-47.9 (8)
C2—O1—C3—C4	177.6 (7)	C9—C8—C13—O6	130.6 (6)
O2—C3—C4—C5	-2.1 (10)	C7—C8—C13—O5	131.6 (5)
O1—C3—C4—C5	173.4 (6)	C9—C8—C13—O5	-49.9 (6)
C3—C4—C5—C6	-175.8 (5)	O4—S—C15—C20	28.1 (5)
C7—N—C6—C5	-69.4 (6)	O3—S—C15—C20	159.4 (4)
S—N—C6—C5	147.8 (4)	N—S—C15—C20	-86.6 (5)
C4—C5—C6—N	-65.9 (6)	O4—S—C15—C16	-157.0 (4)
C6—N—C7—C8	137.4 (5)	O3—S—C15—C16	-25.7 (5)
S—N—C7—C8	-79.3 (5)	N—S—C15—C16	88.2 (5)
C6—N—C7—C12	-38.6 (7)	C20—C15—C16—C17	-3.8 (9)
S—N—C7—C12	104.8 (5)	S—C15—C16—C17	-178.8 (5)
C12—C7—C8—C9	-1.6 (7)	C15—C16—C17—C18	4.0 (10)
N—C7—C8—C9	-177.7 (5)	C16—C17—C18—C19	-1.0 (10)
C12—C7—C8—C13	176.8 (5)	C16—C17—C18—C21	178.6 (6)
N—C7—C8—C13	0.8 (7)	C17—C18—C19—C20	-2.1 (10)
C7—C8—C9—C10	1.7 (8)	C21—C18—C19—C20	178.3 (6)
C13—C8—C9—C10	-176.9 (5)	C18—C19—C20—C15	2.2 (9)
C8—C9—C10—C11	-0.4 (8)	C16—C15—C20—C19	0.7 (8)
C8—C9—C10—Cl	178.9 (4)	S—C15—C20—C19	175.7 (5)

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C6—H6A···O4	0.97	2.41	2.903 (6)	111
C11—H11A···O3 ⁱ	0.93	2.55	3.275 (6)	135
C14—H14A···O2 ⁱⁱ	0.96	2.60	3.338 (7)	134
C17—H17A···O2 ⁱⁱⁱ	0.93	2.59	3.414 (8)	148

Symmetry codes: (i) $-x, y-1/2, -z+1/2$; (ii) $x, y+1, z$; (iii) $-x, y+1/2, -z+1/2$.

supplementary materials

Fig. 1

