

A monoclinic modification of 2-[(1,3-benzothiazol-2-yl)iminomethyl]phenol

Abdullah M. Asiri,^a Salman A. Khan,^a Kong Wai Tan^b and Seik Weng Ng^{b*}

^aChemistry Department, Faculty of Science, King Abdul Aziz University, PO Box 80203, Jeddah 21589, Saudi Arabia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 16 June 2010; accepted 19 June 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.040; wR factor = 0.110; data-to-parameter ratio = 15.8.

In the title Schiff base, $\text{C}_{14}\text{H}_{10}\text{N}_2\text{OS}$, the azomethine double bond is in an *E* configuration; the benzothiazolyl ring (r.m.s. deviation = 0.007 Å) is coplanar with the phenylene ring (r.m.s. deviation = 0.007 Å), the two rings being slightly bent at 2.6 (1)°. The hydroxy H atom forms an intramolecular hydrogen bond to the imino group. The bond dimensions of the monoclinic modification are similar to those of the orthorhombic modification [Liu *et al.* (2009). *Acta Cryst. E65*, o738].

Related literature

For an orthorhombic modification of this structure, see: Liu *et al.* (2009).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{10}\text{N}_2\text{OS}$

$M_r = 254.30$

Monoclinic, Pn
 $a = 8.6391 (4)\text{ \AA}$
 $b = 6.2313 (4)\text{ \AA}$
 $c = 11.4459 (8)\text{ \AA}$
 $\beta = 108.893 (1)$ °
 $V = 582.97 (6)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.26\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.14 \times 0.13 \times 0.08\text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.964$, $T_{\max} = 0.979$

5307 measured reflections
2599 independent reflections
2512 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.110$
 $S = 1.05$
2599 reflections
164 parameters
2 restraints

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.40\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
1242 Friedel pairs
Flack parameter: 0.27 (8)

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1···N1	0.87	1.73	2.550 (2)	156

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdul Aziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2043).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Liu, S.-Q., Bi, C.-F., Chen, L.-Y. & Fan, Y.-H. (2009). *Acta Cryst. E65*, o738.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.

supplementary materials

Acta Cryst. (2010). E66, o1826 [doi:10.1107/S1600536810023755]

A monoclinic modification of 2-[(1,3-benzothiazol-2-yl)iminomethyl]phenol

A. M. Asiri, S. A. Khan, K. W. Tan and S. W. Ng

Comment

The orthorhombic modification of 2-[(1,3-benzothiazol-2-yl)iminomethyl]phenol (Scheme I) is a flat molecule having a (calculated) density of 1.409 g ml^{-1} (Liu *et al.*, 2009). In the monoclinic modification, the packing is more compact (calculated density 1.449 g ml^{-1}). The benzisotholyl ring [r.m.s. deviation 0.007 \AA] is co-planar with the phenylene ring [r.m.s. deviation 0.007 \AA], the two rings being slightly bent by $2.6(1)^\circ$ only. The hydroxy H-atom forms an intramolecular hydrogen bond to the imino group (Fig. 1). The r.m.s. deviation of the non-hydrogen atoms for the least-squares plane is 0.034 \AA , and all deviations deviate by 0.002 \AA only.

Experimental

2-Aminobenzothiazole (0.50 g, 4.4 mol) and salicylaldehyde (0.66 g, 4.4 mol) were heated in methanol (15 ml) for 5 h. Yellowish-orange crystals deposited when the solution was set aside for a day.

Refinement

Carbon-bound H-atoms were placed in calculated positions [$\text{C-H } 0.95 \text{ \AA}$, $U(\text{H}) 1.2U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation. The hydroxy H-atom was located in a difference Fourier map, but attempts to refine it even with a distance restraint led to a small temperature factor. The position and temperature factor were not refined. The structure is a racemic twin, the explicit refinement of the Flack parameter from 1242 Friedel pairs gave a value of $0.27(8)$.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{14}\text{H}_{10}\text{N}_2\text{OS}$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

2-[(1,3-benzothiazol-2-yl)iminomethyl]phenol

Crystal data

$\text{C}_{14}\text{H}_{10}\text{N}_2\text{OS}$

$F(000) = 264$

$M_r = 254.30$

$D_x = 1.449 \text{ Mg m}^{-3}$

Monoclinic, Pn

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Hall symbol: P -2yac

Cell parameters from 3896 reflections

$a = 8.6391(4) \text{ \AA}$

$\theta = 2.6\text{--}28.3^\circ$

supplementary materials

$b = 6.2313 (4) \text{ \AA}$	$\mu = 0.26 \text{ mm}^{-1}$
$c = 11.4459 (8) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 108.893 (1)^\circ$	Prism, orange
$V = 582.97 (6) \text{ \AA}^3$	$0.14 \times 0.13 \times 0.08 \text{ mm}$
$Z = 2$	

Data collection

Bruker SMART APEX diffractometer	2599 independent reflections
Radiation source: fine-focus sealed tube graphite	2512 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.029$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.6^\circ$
$T_{\text{min}} = 0.964, T_{\text{max}} = 0.979$	$h = -11 \rightarrow 10$
5307 measured reflections	$k = -8 \rightarrow 8$
	$l = -14 \rightarrow 14$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.110$	$w = 1/[\sigma^2(F_o^2) + (0.0831P)^2 + 0.0645P]$
	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.05$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2599 reflections	$\Delta\rho_{\text{max}} = 0.40 \text{ e \AA}^{-3}$
164 parameters	$\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$
2 restraints	Absolute structure: Flack (1983), 1242 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.27 (8)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.50007 (6)	0.06429 (7)	0.49993 (5)	0.01864 (15)
O1	0.7048 (2)	0.5684 (3)	0.89741 (16)	0.0277 (4)
H1	0.6924	0.4623	0.8462	0.042*
N1	0.6095 (2)	0.3254 (3)	0.70823 (16)	0.0186 (4)
N2	0.7613 (2)	0.0261 (3)	0.69409 (17)	0.0193 (4)
C1	0.5700 (3)	0.6867 (4)	0.84466 (19)	0.0195 (4)
C2	0.5445 (3)	0.8696 (4)	0.9059 (2)	0.0222 (4)
H2	0.6227	0.9096	0.9825	0.027*
C3	0.4058 (3)	0.9936 (4)	0.8561 (2)	0.0226 (5)
H3	0.3895	1.1183	0.8985	0.027*
C4	0.2898 (3)	0.9363 (3)	0.7441 (2)	0.0228 (5)
H4	0.1937	1.0201	0.7108	0.027*

C5	0.3153 (3)	0.7571 (4)	0.68168 (19)	0.0214 (4)
H5	0.2366	0.7193	0.6049	0.026*
C6	0.4554 (3)	0.6303 (3)	0.72981 (19)	0.0175 (4)
C7	0.4806 (3)	0.4437 (3)	0.6638 (2)	0.0188 (4)
H7	0.4019	0.4071	0.5868	0.023*
C8	0.6349 (3)	0.1448 (3)	0.64566 (19)	0.0184 (4)
C9	0.7588 (3)	-0.1475 (3)	0.61665 (18)	0.0178 (4)
C10	0.8777 (3)	-0.3090 (4)	0.6428 (2)	0.0228 (4)
H10	0.9681	-0.3056	0.7170	0.027*
C11	0.8612 (3)	-0.4729 (4)	0.5589 (2)	0.0249 (5)
H11	0.9414	-0.5832	0.5757	0.030*
C12	0.7290 (3)	-0.4798 (4)	0.4498 (2)	0.0229 (5)
H12	0.7212	-0.5939	0.3931	0.027*
C13	0.6081 (3)	-0.3219 (4)	0.4225 (2)	0.0213 (4)
H13	0.5170	-0.3277	0.3488	0.026*
C14	0.6257 (3)	-0.1556 (3)	0.5070 (2)	0.0190 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0181 (2)	0.0172 (2)	0.0187 (2)	-0.0003 (2)	0.00322 (17)	-0.00262 (19)
O1	0.0216 (9)	0.0314 (10)	0.0232 (9)	0.0063 (7)	-0.0024 (7)	-0.0064 (6)
N1	0.0193 (8)	0.0166 (8)	0.0183 (8)	-0.0016 (7)	0.0040 (7)	-0.0015 (7)
N2	0.0222 (10)	0.0148 (8)	0.0191 (9)	-0.0020 (7)	0.0042 (7)	-0.0008 (7)
C1	0.0181 (10)	0.0220 (10)	0.0181 (9)	-0.0008 (8)	0.0051 (8)	-0.0004 (8)
C2	0.0235 (11)	0.0215 (10)	0.0211 (10)	-0.0032 (9)	0.0066 (9)	-0.0052 (8)
C3	0.0265 (12)	0.0193 (10)	0.0263 (11)	-0.0017 (9)	0.0144 (9)	-0.0023 (8)
C4	0.0228 (12)	0.0211 (11)	0.0255 (11)	0.0054 (8)	0.0095 (9)	0.0051 (8)
C5	0.0197 (10)	0.0245 (10)	0.0190 (9)	0.0002 (9)	0.0048 (8)	0.0025 (8)
C6	0.0179 (10)	0.0178 (9)	0.0164 (9)	-0.0011 (8)	0.0050 (8)	-0.0012 (8)
C7	0.0173 (10)	0.0200 (11)	0.0177 (10)	-0.0029 (8)	0.0035 (8)	-0.0002 (7)
C8	0.0196 (10)	0.0168 (10)	0.0182 (9)	-0.0025 (8)	0.0051 (8)	0.0001 (7)
C9	0.0221 (10)	0.0148 (9)	0.0175 (9)	-0.0025 (8)	0.0077 (8)	-0.0021 (8)
C10	0.0228 (11)	0.0209 (10)	0.0249 (10)	0.0016 (9)	0.0081 (8)	0.0032 (8)
C11	0.0245 (12)	0.0224 (11)	0.0311 (12)	0.0032 (9)	0.0136 (10)	0.0047 (9)
C12	0.0282 (13)	0.0182 (10)	0.0276 (12)	-0.0036 (9)	0.0166 (10)	-0.0058 (9)
C13	0.0205 (10)	0.0224 (11)	0.0208 (10)	-0.0043 (9)	0.0063 (8)	-0.0018 (8)
C14	0.0178 (10)	0.0156 (9)	0.0249 (10)	0.0012 (8)	0.0088 (8)	0.0022 (8)

Geometric parameters (\AA , $^\circ$)

S1—C14	1.733 (2)	C4—H4	0.9500
S1—C8	1.770 (2)	C5—C6	1.400 (3)
O1—C1	1.345 (3)	C5—H5	0.9500
O1—H1	0.8672	C6—C7	1.442 (3)
N1—C7	1.294 (3)	C7—H7	0.9500
N1—C8	1.389 (3)	C9—C10	1.399 (3)
N2—C8	1.286 (3)	C9—C14	1.402 (3)
N2—C9	1.394 (3)	C10—C11	1.378 (3)

supplementary materials

C1—C2	1.393 (3)	C10—H10	0.9500
C1—C6	1.410 (3)	C11—C12	1.393 (4)
C2—C3	1.383 (4)	C11—H11	0.9500
C2—H2	0.9500	C12—C13	1.395 (3)
C3—C4	1.394 (4)	C12—H12	0.9500
C3—H3	0.9500	C13—C14	1.392 (3)
C4—C5	1.381 (3)	C13—H13	0.9500
C14—S1—C8	88.21 (11)	N1—C7—H7	119.7
C1—O1—H1	102.4	C6—C7—H7	119.7
C7—N1—C8	121.20 (18)	N2—C8—N1	119.83 (19)
C8—N2—C9	109.65 (18)	N2—C8—S1	116.81 (16)
O1—C1—C2	118.56 (19)	N1—C8—S1	123.36 (17)
O1—C1—C6	121.8 (2)	N2—C9—C10	124.3 (2)
C2—C1—C6	119.7 (2)	N2—C9—C14	115.84 (19)
C3—C2—C1	120.4 (2)	C10—C9—C14	119.9 (2)
C3—C2—H2	119.8	C11—C10—C9	118.8 (2)
C1—C2—H2	119.8	C11—C10—H10	120.6
C2—C3—C4	120.3 (2)	C9—C10—H10	120.6
C2—C3—H3	119.8	C10—C11—C12	121.1 (2)
C4—C3—H3	119.8	C10—C11—H11	119.4
C5—C4—C3	119.7 (2)	C12—C11—H11	119.4
C5—C4—H4	120.2	C13—C12—C11	121.1 (2)
C3—C4—H4	120.2	C13—C12—H12	119.5
C4—C5—C6	121.0 (2)	C11—C12—H12	119.5
C4—C5—H5	119.5	C14—C13—C12	117.7 (2)
C6—C5—H5	119.5	C14—C13—H13	121.2
C5—C6—C1	118.87 (19)	C12—C13—H13	121.2
C5—C6—C7	120.47 (19)	C13—C14—C9	121.5 (2)
C1—C6—C7	120.7 (2)	C13—C14—S1	129.07 (18)
N1—C7—C6	120.7 (2)	C9—C14—S1	109.47 (16)
O1—C1—C2—C3	−178.7 (2)	C14—S1—C8—N2	1.30 (18)
C6—C1—C2—C3	1.4 (3)	C14—S1—C8—N1	−178.28 (18)
C1—C2—C3—C4	0.2 (4)	C8—N2—C9—C10	−179.1 (2)
C2—C3—C4—C5	−1.2 (4)	C8—N2—C9—C14	0.2 (3)
C3—C4—C5—C6	0.6 (3)	N2—C9—C10—C11	179.7 (2)
C4—C5—C6—C1	0.9 (3)	C14—C9—C10—C11	0.4 (3)
C4—C5—C6—C7	−180.0 (2)	C9—C10—C11—C12	−0.1 (3)
O1—C1—C6—C5	178.2 (2)	C10—C11—C12—C13	−0.6 (4)
C2—C1—C6—C5	−1.9 (3)	C11—C12—C13—C14	1.1 (3)
O1—C1—C6—C7	−0.9 (3)	C12—C13—C14—C9	−0.8 (3)
C2—C1—C6—C7	179.0 (2)	C12—C13—C14—S1	179.15 (17)
C8—N1—C7—C6	179.8 (2)	N2—C9—C14—C13	−179.33 (19)
C5—C6—C7—N1	−179.6 (2)	C10—C9—C14—C13	0.0 (3)
C1—C6—C7—N1	−0.5 (3)	N2—C9—C14—S1	0.7 (2)
C9—N2—C8—N1	178.51 (18)	C10—C9—C14—S1	−179.89 (17)
C9—N2—C8—S1	−1.1 (2)	C8—S1—C14—C13	179.0 (2)
C7—N1—C8—N2	−177.6 (2)	C8—S1—C14—C9	−1.04 (16)
C7—N1—C8—S1	2.0 (3)		

Hydrogen-bond geometry (Å, °)

$D\text{---H}\cdots A$	$D\text{---H}$	$H\cdots A$	$D\cdots A$	$D\text{---H}\cdots A$
O1—H1…N1	0.87	1.73	2.550 (2)	156

supplementary materials

Fig. 1

