

2-(7-Methoxy-1-naphthyl)acetonitrile

Wen-bin Wei,^{a,b} Ru Jia,^a Jie Sun^a and Hai-Bo Wang^{a*}

^aCollege of Food Science and Light Industry, Nanjing University of Technology, Xinmofan Road No.5 Nanjing, Nanjing 210009, People's Republic of China, and

^bCollege of Science, Nanjing University of Technology, Xinmofan Road No.5 Nanjing, Nanjing 210009, People's Republic of China

Correspondence e-mail: wanghaibo@njut.edu.cn


Received 18 June 2010; accepted 23 June 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.059; wR factor = 0.163; data-to-parameter ratio = 13.9.

The molecule of the title compound, $\text{C}_{13}\text{H}_{11}\text{NO}$, is almost planar (r.m.s. deviation = 0.013 Å), apart from the cyanide group, for which the C and N atoms deviate from the mean plane of the other atoms by 0.341 (3) and 0.571 (4) Å, respectively. In the crystal, weak aromatic $\pi-\pi$ stacking [centroid–centroid distance = 3.758 (3) Å] may help to stabilize the structure.

Related literature

For background to the use of naphthylethyl acetonitrile as an intermediate for the synthesis of *N*-naphthylethyl amide derivatives, see: Depreux & Lesieur (1994). For further synthetic details, see: Yous & Andrieux (1992).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{11}\text{NO}$
 $M_r = 197.23$
Monoclinic, $P2_1/n$
 $a = 7.5110 (15)\text{ \AA}$
 $b = 9.6170 (19)\text{ \AA}$
 $c = 14.731 (3)\text{ \AA}$
 $\beta = 101.03 (3)^\circ$

$V = 1044.4 (4)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.08\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.30 \times 0.20 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4
diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.976$, $T_{\max} = 0.992$
1971 measured reflections

1897 independent reflections
1045 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.011$
3 standard reflections every 200
reflections
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.163$
 $S = 1.00$
1897 reflections

136 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.14\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.14\text{ e \AA}^{-3}$

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON*.

The authors thank the Center of Testing and Analysis of Nanjing University for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5505).

References

- Depreux, P. & Lesieur, D. (1994). *J. Med. Chem.* **37**, 3231–3239.
Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
Yous, S. & Andrieux, J. (1992). *J. Med. Chem.* **35**, 1484–1486.

supplementary materials

Acta Cryst. (2010). E66, o1868 [doi:10.1107/S1600536810024372]

2-(7-Methoxy-1-naphthyl)acetonitrile

W. Wei, R. Jia, J. Sun and H.-B. Wang

Comment

Naphthylethyl acetonitrile is an important pharmaceutical intermediate for synthesizing *N*-naphthylethyl amide derivatives which was evaluated as melatonin receptor ligands (Depreux & Lesieur, 1994). We report herein the crystal structure of the title compound, (I).

In the molecule of the title compound (Fig 1), the bond lengths and angles are within normal ranges. Rings A (C3—C7/C12), B (C8—C12) are, of course, planar, and they are oriented at dihedral angle A/B= 1.10 (3) °. So, they are nearly co-planar. No classical hydrogen bond was found in the molecule. The π — π contacts between the naphthalene rings, $Cg1—Cg2^i$ [symmetry codes: $-x, 1 - y, 1 - z$, where $Cg1$ and $Cg2$ are centroids of the rings A (C3—C7/C10/C12), and B (C8—C12), respectively] may further stabilize the structure, with centroid-centroid distances of 3.758 (3) Å.

Experimental

(7-Methoxy-1-naphthyl)acetic acid was reacted with thionyl chloride in CHCl_3 , and the crude acid chloride was treated with aqueous ammonia to produce (7-Methoxy-1-naphthyl)acetamide. Dehydration of this amide with trifluoroacetic anhydride in THF at 273 K gave the title compound (Yous & Andrieux, 1992). Crystals suitable for X-ray analysis were obtained by slow evaporation of a methanol solution (yield; 66%, m.p. 353 K).

Refinement

H atoms were positioned geometrically, with C—H = 0.93 Å for aromatic H and C—H = 0.96 and 0.97 Å for methyl and methylene H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{O})$, where $x = 1.5$ for OH H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. View of the title compound with displacement ellipsoids for non-H atoms drawn at the 50% probability level.

supplementary materials




Fig. 2. A packing diagram of title molecule.

2-(7-Methoxy-1-naphthyl)acetonitrile

Crystal data

C ₁₃ H ₁₁ NO	$F(000) = 416$
$M_r = 197.23$	$D_x = 1.254 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Melting point: 353 K
Hall symbol: -P 2yn	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.5110 (15) \text{ \AA}$	Cell parameters from 25 reflections
$b = 9.6170 (19) \text{ \AA}$	$\theta = 9\text{--}13^\circ$
$c = 14.731 (3) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$\beta = 101.03 (3)^\circ$	$T = 293 \text{ K}$
$V = 1044.4 (4) \text{ \AA}^3$	Block, colorless
$Z = 4$	$0.30 \times 0.20 \times 0.10 \text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer	1045 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube graphite	$R_{\text{int}} = 0.011$
$\omega/2\theta$ scans	$\theta_{\text{max}} = 25.3^\circ, \theta_{\text{min}} = 2.5^\circ$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$h = -9 \rightarrow 8$
$T_{\text{min}} = 0.976, T_{\text{max}} = 0.992$	$k = -11 \rightarrow 0$
1971 measured reflections	$l = 0 \rightarrow 17$
1897 independent reflections	3 standard reflections every 200 reflections intensity decay: 1%

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.059$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.163$	H-atom parameters constrained
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.069P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
1897 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
136 parameters	$\Delta\rho_{\text{max}} = 0.14 \text{ e \AA}^{-3}$

0 restraints

$\Delta\rho_{\min} = -0.14 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O	0.8616 (3)	-0.2547 (2)	0.13823 (13)	0.0718 (6)
N	0.7349 (5)	0.1342 (3)	-0.3297 (2)	0.1076 (12)
C1	0.7327 (4)	0.0786 (3)	-0.2623 (2)	0.0691 (9)
C2	0.7316 (4)	0.0085 (3)	-0.17484 (17)	0.0565 (7)
H2A	0.8471	-0.0380	-0.1550	0.068*
H2B	0.6376	-0.0620	-0.1842	0.068*
C3	0.6992 (3)	0.1069 (3)	-0.09873 (18)	0.0484 (7)
C4	0.6435 (4)	0.2412 (3)	-0.1174 (2)	0.0622 (8)
H4A	0.6228	0.2727	-0.1782	0.075*
C5	0.6171 (4)	0.3319 (3)	-0.0472 (3)	0.0716 (9)
H5A	0.5794	0.4227	-0.0613	0.086*
C6	0.6466 (4)	0.2874 (3)	0.0411 (2)	0.0685 (9)
H6A	0.6293	0.3486	0.0876	0.082*
C7	0.7028 (4)	0.1504 (3)	0.06461 (19)	0.0552 (8)
C8	0.7346 (4)	0.1027 (3)	0.1570 (2)	0.0675 (9)
H8A	0.7200	0.1635	0.2042	0.081*
C9	0.7857 (4)	-0.0299 (4)	0.1779 (2)	0.0685 (9)
H9A	0.8060	-0.0596	0.2391	0.082*
C10	0.8086 (4)	-0.1236 (3)	0.10760 (19)	0.0564 (7)
C11	0.7816 (3)	-0.0820 (3)	0.01827 (18)	0.0495 (7)
H11A	0.7979	-0.1448	-0.0275	0.059*
C12	0.7285 (3)	0.0571 (3)	-0.00618 (18)	0.0471 (7)
C13	0.8857 (4)	-0.3548 (3)	0.0708 (2)	0.0723 (9)
H13A	0.9233	-0.4415	0.1007	0.108*
H13B	0.9768	-0.3229	0.0380	0.108*
H13C	0.7733	-0.3677	0.0280	0.108*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O	0.0897 (16)	0.0689 (14)	0.0570 (12)	0.0037 (12)	0.0148 (11)	0.0088 (11)

supplementary materials

N	0.177 (4)	0.085 (2)	0.062 (2)	-0.005 (2)	0.024 (2)	0.0025 (17)
C1	0.092 (2)	0.064 (2)	0.0507 (18)	-0.0084 (18)	0.0111 (16)	-0.0041 (16)
C2	0.0636 (19)	0.0525 (17)	0.0549 (17)	-0.0034 (14)	0.0150 (14)	-0.0016 (14)
C3	0.0441 (15)	0.0478 (16)	0.0553 (17)	-0.0072 (13)	0.0142 (13)	-0.0041 (13)
C4	0.065 (2)	0.0532 (18)	0.070 (2)	-0.0012 (15)	0.0164 (15)	0.0030 (16)
C5	0.074 (2)	0.0490 (18)	0.097 (3)	0.0035 (16)	0.0285 (19)	-0.0043 (18)
C6	0.073 (2)	0.057 (2)	0.084 (2)	-0.0069 (16)	0.0352 (18)	-0.0222 (17)
C7	0.0506 (17)	0.0552 (19)	0.0636 (19)	-0.0093 (14)	0.0210 (14)	-0.0135 (15)
C8	0.074 (2)	0.073 (2)	0.062 (2)	-0.0102 (18)	0.0293 (16)	-0.0217 (17)
C9	0.078 (2)	0.082 (2)	0.0499 (17)	-0.0113 (19)	0.0234 (16)	-0.0027 (17)
C10	0.0572 (18)	0.0583 (18)	0.0552 (18)	-0.0043 (14)	0.0141 (14)	0.0006 (15)
C11	0.0491 (16)	0.0508 (17)	0.0519 (17)	-0.0058 (13)	0.0182 (13)	-0.0061 (13)
C12	0.0384 (15)	0.0485 (16)	0.0565 (17)	-0.0098 (12)	0.0142 (12)	-0.0069 (13)
C13	0.078 (2)	0.0579 (19)	0.079 (2)	0.0039 (16)	0.0087 (18)	0.0028 (17)

Geometric parameters (\AA , $^\circ$)

O—C10	1.372 (3)	C6—H6A	0.9300
O—C13	1.419 (3)	C7—C8	1.413 (4)
N—C1	1.130 (4)	C7—C12	1.417 (3)
C1—C2	1.456 (4)	C8—C9	1.350 (4)
C2—C3	1.522 (3)	C8—H8A	0.9300
C2—H2A	0.9700	C9—C10	1.407 (4)
C2—H2B	0.9700	C9—H9A	0.9300
C3—C4	1.369 (4)	C10—C11	1.353 (4)
C3—C12	1.422 (3)	C11—C12	1.422 (4)
C4—C5	1.396 (4)	C11—H11A	0.9300
C4—H4A	0.9300	C13—H13A	0.9600
C5—C6	1.347 (4)	C13—H13B	0.9600
C5—H5A	0.9300	C13—H13C	0.9600
C6—C7	1.406 (4)		
C10—O—C13	117.4 (2)	C8—C7—C12	118.8 (3)
N—C1—C2	179.1 (4)	C9—C8—C7	120.9 (3)
C1—C2—C3	113.2 (2)	C9—C8—H8A	119.5
C1—C2—H2A	108.9	C7—C8—H8A	119.5
C3—C2—H2A	108.9	C8—C9—C10	120.4 (3)
C1—C2—H2B	108.9	C8—C9—H9A	119.8
C3—C2—H2B	108.9	C10—C9—H9A	119.8
H2A—C2—H2B	107.8	C11—C10—O	124.9 (3)
C4—C3—C12	119.7 (3)	C11—C10—C9	120.6 (3)
C4—C3—C2	121.6 (3)	O—C10—C9	114.5 (3)
C12—C3—C2	118.7 (2)	C10—C11—C12	120.5 (3)
C3—C4—C5	121.5 (3)	C10—C11—H11A	119.8
C3—C4—H4A	119.2	C12—C11—H11A	119.8
C5—C4—H4A	119.2	C7—C12—C3	118.3 (3)
C6—C5—C4	119.8 (3)	C7—C12—C11	118.7 (2)
C6—C5—H5A	120.1	C3—C12—C11	123.0 (2)
C4—C5—H5A	120.1	O—C13—H13A	109.5
C5—C6—C7	121.4 (3)	O—C13—H13B	109.5

supplementary materials

C5—C6—H6A	119.3	H13A—C13—H13B	109.5
C7—C6—H6A	119.3	O—C13—H13C	109.5
C6—C7—C8	121.9 (3)	H13A—C13—H13C	109.5
C6—C7—C12	119.3 (3)	H13B—C13—H13C	109.5

supplementary materials

Fig. 1




Fig. 2

