

## catena-Poly[[tris(4-fluorobenzyl)tin(IV)]- $\mu$ -2-[(piperidin-1-yl)carbothioylsulfanyl]-acetato- $\kappa^2$ O:O']

Chun Thy Keng, Kong Mun Lo and Seik Weng Ng\*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 14 July 2010; accepted 15 July 2010

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å; disorder in main residue;  $R$  factor = 0.034;  $wR$  factor = 0.077; data-to-parameter ratio = 17.5.

Adjacent units of the title polymeric complex,  $[\text{Sn}(\text{C}_7\text{H}_6\text{F})_3(\text{C}_8\text{H}_{12}\text{NO}_2\text{S}_2)]$ , are bridged by the carboxylate ion into a helical chain running along the  $b$  axis. The Sn(IV) atom shows a distorted  $trans$ - $\text{C}_3\text{SnO}_2$  trigonal-bipyramidal coordination and is displaced by 0.113 (2) Å out of the  $\text{C}_3\text{Sn}$  girdle in the direction of the covalently bonded O atom. The ring is disordered of two positions with an occupancy of 0.631 (4) for the major occupied site.

### Related literature

For a comment on the repeat distance of carboxylate-bridged trialkyltin carboxylates, see: Ng *et al.* (1989).



### Experimental

#### Crystal data

$[\text{Sn}(\text{C}_7\text{H}_6\text{F})_3(\text{C}_8\text{H}_{12}\text{NO}_2\text{S}_2)]$   
 $M_r = 664.35$   
Monoclinic,  $P2_1/n$   
 $a = 11.8016$  (7) Å  
 $b = 10.4572$  (6) Å  
 $c = 23.0334$  (13) Å  
 $\beta = 94.013$  (1)°

$V = 2835.6$  (3) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 1.10$  mm<sup>-1</sup>  
 $T = 100$  K  
0.30 × 0.25 × 0.20 mm

#### Data collection

Bruker SMART APEX  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.735$ ,  $T_{\max} = 0.811$

25437 measured reflections  
6498 independent reflections  
5328 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.038$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.077$   
 $S = 1.05$   
6498 reflections  
371 parameters

17 restraints  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.50$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -1.19$  e Å<sup>-3</sup>

**Table 1**

Selected geometric parameters (Å, °).

Sn1—C1	2.144 (3)	Sn1—O1	2.175 (2)
Sn1—C8	2.155 (3)	Sn1—O2 <sup>i</sup>	2.339 (2)
Sn1—C15	2.158 (3)		
O1—Sn1—O2 <sup>i</sup>	170.59 (7)		

Symmetry code: (i)  $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *PUBLICIF* (Westrip, 2010).

We thank the University of Malaya (RG020/09AFR) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5300).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Ng, S. W., Chen, W. & Kumar Das, V. G. (1989). *J. Organomet. Chem.* **346**, 59–64.  
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

**supplementary materials**

*Acta Cryst.* (2010). E66, m978 [ doi:10.1107/S1600536810028321 ]

***catena*-Poly[[tris(4-fluorobenzyl)tin(IV)]- $\mu$ -2-[(piperidin-1-yl)carbothioylsulfanyl]acetato- $\kappa^2$ O:O']**

**C. T. Keng, K. M. Lo and S. W. Ng**

### Comment

Trialkyltin carboxylates generally adopt linear carboxylate-bridged motifs; the repeat distance of such polymers is independent of the nature of the carboxylate group. This feature of trialkyltin carboxylates is also borne out in tris(4-fluorobenzyl)tin (*N*-pentamethylenecarbamothio)sulfanylacetate (Scheme I), whose repeat distance, *i.e.*, half the *b*-axis, of 5.23 Å, falls within the range of repeat distances reported (Ng *et al.*, 1989). The tin atom shows *trans*-C<sub>3</sub>SnO<sub>2</sub> trigonal bipyramidal coordination; the tin atom is displaced by 0.113 (2) Å out of the C<sub>3</sub>Sn girdle in the direction of the covalently-bonded oxygen atom. The O-Sn-O angle is close to linearity (Table 1).

### Experimental

Tris(4-fluorobenzyl)tin hydroxide was prepared by hydrolysis of tris(4-fluorobenzyl)tin chloride in dilute sodium hydroxide. The hydroxide (1.0 g, 2.2 mmol) and *N*-piperinyldithiocarbamylacetic acid (0.48 g, 2.2 mmol) were heated in ethanol for an hour. The solution was filtered and then set aside for the growth of crystals.

### Refinement

Hydrogen atoms were placed in calculated positions (C–H 0.95–0.99 Å) and were included in the refinement in the riding model approximation, with  $U(\text{H})$  set to  $1.2U_{\text{eq}}(\text{C})$ .

The carboxylate ion is disordered in the C–N(CH<sub>2</sub>)<sub>5</sub> portion only; the sulfur atoms are ordered. Pairs of carbon-sulfur single-bond distances, carbon-sulfur double-bond distances, carbon<sub>sulfur</sub>-nitrogen distances, carbon<sub>methylene</sub>-nitrogen distances and carbon-carbon (methylene) distances were restrained to within 0.01 Å of each other. The displacement parameters of the minor (primed) atoms were restrained to be similar to those of the major (unprimed) atoms. The two C(S<sub>2</sub>)–N(C<sub>2</sub>) fragments were restrained to be within 0.01 Å of a plane. The pairs of non-bonded sulfur<sub>single bond</sub>-nitrogen and sulfur<sub>double bond</sub>-nitrogen distances were also restrained, though with a much tighter restraint.

The final difference Fourier map had a hole in the vicinity of O1.

### Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of a portion of polymeric [Sn(C<sub>7</sub>H<sub>6</sub>F)<sub>3</sub>(C<sub>8</sub>H<sub>12</sub>NO<sub>2</sub>S<sub>2</sub>)]<sub>n</sub>; ellipsoids are drawn at the 70% probability level and H atoms are of arbitrary radius. The disorder is not shown. Symmetry code is given in Table 1.

## catena-Poly[[tris(4-fluorobenzyl)tin(IV)]- $\mu$ -2-[(piperidin-1-yl)carbothioylsulfanyl]acetato- $\kappa^2$ O:O']

### Crystal data

[Sn(C <sub>7</sub> H <sub>6</sub> F) <sub>3</sub> (C <sub>8</sub> H <sub>12</sub> NO <sub>2</sub> S <sub>2</sub> )]	$F(000) = 1344$
$M_r = 664.35$	$D_x = 1.556 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 9852 reflections
$a = 11.8016 (7) \text{ \AA}$	$\theta = 2.4\text{--}28.3^\circ$
$b = 10.4572 (6) \text{ \AA}$	$\mu = 1.10 \text{ mm}^{-1}$
$c = 23.0334 (13) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 94.013 (1)^\circ$	Prism, colorless
$V = 2835.6 (3) \text{ \AA}^3$	$0.30 \times 0.25 \times 0.20 \text{ mm}$
$Z = 4$	

### Data collection

Bruker SMART APEX diffractometer	6498 independent reflections
Radiation source: fine-focus sealed tube graphite	5328 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.038$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 27.5^\circ$ , $\theta_{\text{min}} = 1.8^\circ$
$T_{\text{min}} = 0.735$ , $T_{\text{max}} = 0.811$	$h = -15 \rightarrow 15$
25437 measured reflections	$k = -13 \rightarrow 13$
	$l = -29 \rightarrow 29$

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.034$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.077$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0254P)^2 + 5.3214P]$
6498 reflections	where $P = (F_o^2 + 2F_c^2)/3$
371 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
17 restraints	$\Delta\rho_{\text{max}} = 0.50 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -1.19 \text{ e \AA}^{-3}$

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Sn1	0.710443 (15)	0.543605 (18)	0.737119 (9)	0.01476 (6)	
S1	0.57734 (6)	0.05444 (7)	0.62134 (3)	0.01883 (15)	

S2	0.68919 (7)	0.25659 (8)	0.55248 (4)	0.02778 (18)	
F1	0.20300 (17)	0.7560 (2)	0.79264 (11)	0.0494 (6)	
F2	0.2292 (2)	0.5457 (3)	0.56053 (13)	0.0778 (10)	
F3	1.0169 (2)	0.7386 (2)	0.52170 (10)	0.0508 (6)	
O1	0.61265 (18)	0.39691 (19)	0.68910 (10)	0.0219 (5)	
O2	0.71042 (17)	0.22277 (19)	0.71464 (9)	0.0199 (4)	
N1	0.7199 (2)	0.0057 (3)	0.54087 (13)	0.0219 (9)	0.631 (4)
N1'	0.6619 (3)	0.0131 (4)	0.52047 (19)	0.0219 (9)	0.37
C1	0.6237 (2)	0.5024 (3)	0.81369 (13)	0.0203 (6)	
H1A	0.6141	0.4088	0.8174	0.024*	
H1B	0.6699	0.5330	0.8485	0.024*	
C2	0.5100 (2)	0.5657 (3)	0.81084 (13)	0.0194 (6)	
C3	0.4159 (3)	0.5111 (3)	0.78054 (14)	0.0244 (7)	
H3	0.4231	0.4294	0.7632	0.029*	
C4	0.3121 (3)	0.5734 (4)	0.77516 (16)	0.0318 (8)	
H4	0.2479	0.5347	0.7551	0.038*	
C5	0.3044 (3)	0.6923 (4)	0.79952 (16)	0.0316 (8)	
C6	0.3936 (3)	0.7503 (3)	0.83053 (16)	0.0306 (8)	
H6	0.3853	0.8322	0.8475	0.037*	
C7	0.4965 (3)	0.6853 (3)	0.83628 (14)	0.0236 (7)	
H7	0.5590	0.7233	0.8580	0.028*	
C8	0.6459 (3)	0.6860 (3)	0.67581 (14)	0.0212 (6)	
H8A	0.6363	0.7671	0.6971	0.025*	
H8B	0.7037	0.7011	0.6474	0.025*	
C9	0.5355 (2)	0.6548 (3)	0.64269 (13)	0.0192 (6)	
C10	0.4343 (3)	0.7070 (4)	0.65835 (15)	0.0309 (8)	
H10	0.4356	0.7684	0.6888	0.037*	
C11	0.3307 (3)	0.6717 (5)	0.63062 (18)	0.0463 (11)	
H11	0.2617	0.7076	0.6420	0.056*	
C12	0.3303 (3)	0.5845 (5)	0.58683 (19)	0.0475 (11)	
C13	0.4278 (3)	0.5319 (4)	0.56833 (16)	0.0375 (9)	
H13	0.4253	0.4725	0.5370	0.045*	
C14	0.5303 (3)	0.5679 (3)	0.59670 (14)	0.0249 (7)	
H14	0.5988	0.5324	0.5845	0.030*	
C15	0.8782 (2)	0.4800 (3)	0.71871 (13)	0.0187 (6)	
H15A	0.9318	0.4967	0.7529	0.022*	
H15B	0.8769	0.3867	0.7114	0.022*	
C16	0.9180 (2)	0.5479 (3)	0.66665 (13)	0.0183 (6)	
C17	0.8872 (3)	0.5034 (3)	0.61081 (14)	0.0242 (7)	
H17	0.8426	0.4280	0.6062	0.029*	
C18	0.9200 (3)	0.5665 (3)	0.56162 (15)	0.0324 (8)	
H18	0.8983	0.5354	0.5237	0.039*	
C19	0.9847 (3)	0.6754 (3)	0.56939 (16)	0.0323 (8)	
C20	1.0173 (3)	0.7237 (3)	0.62352 (16)	0.0287 (7)	
H20	1.0617	0.7994	0.6277	0.034*	
C21	0.9836 (2)	0.6587 (3)	0.67181 (14)	0.0222 (6)	
H21	1.0058	0.6906	0.7095	0.027*	
C22	0.6302 (2)	0.2774 (3)	0.68712 (12)	0.0163 (6)	
C23	0.5386 (2)	0.2056 (3)	0.65075 (13)	0.0189 (6)	

## supplementary materials

H23A	0.5114	0.2615	0.6180	0.023*	
H23B	0.4738	0.1915	0.6751	0.023*	
C24	0.6704 (3)	0.1026 (4)	0.56691 (15)	0.0166 (13)	0.631 (4)
C25	0.7999 (3)	0.0258 (5)	0.49474 (17)	0.0273 (10)	0.631 (4)
H25A	0.8750	-0.0103	0.5077	0.033*	0.631 (4)
H25B	0.8097	0.1187	0.4884	0.033*	0.631 (4)
C26	0.7582 (5)	-0.0356 (5)	0.4389 (2)	0.0273 (11)	0.631 (4)
H26A	0.8159	-0.0255	0.4100	0.033*	0.631 (4)
H26B	0.6878	0.0076	0.4234	0.033*	0.631 (4)
C27	0.7341 (5)	-0.1788 (5)	0.4475 (2)	0.0275 (14)	0.631 (4)
H27A	0.6973	-0.2150	0.4112	0.033*	0.631 (4)
H27B	0.8064	-0.2249	0.4566	0.033*	0.631 (4)
C28	0.6562 (5)	-0.1965 (5)	0.4974 (2)	0.0292 (11)	0.631 (4)
H28A	0.5804	-0.1607	0.4857	0.035*	0.631 (4)
H28B	0.6470	-0.2889	0.5050	0.035*	0.631 (4)
C29	0.7039 (3)	-0.1307 (4)	0.55303 (17)	0.0224 (9)	0.631 (4)
H29A	0.6506	-0.1411	0.5840	0.027*	0.631 (4)
H29B	0.7774	-0.1699	0.5666	0.027*	0.631 (4)
C24'	0.6478 (3)	0.1059 (6)	0.5589 (3)	0.029 (3)	0.369 (4)
C25'	0.7185 (5)	0.0395 (8)	0.4665 (3)	0.0273 (10)	0.37
H25C	0.7472	0.1285	0.4672	0.033*	0.369 (4)
H25D	0.6632	0.0302	0.4324	0.033*	0.369 (4)
C26'	0.8152 (7)	-0.0511 (8)	0.4610 (4)	0.0273 (11)	0.37
H26C	0.8744	-0.0349	0.4927	0.033*	0.369 (4)
H26D	0.8491	-0.0365	0.4235	0.033*	0.369 (4)
C27'	0.7748 (9)	-0.1904 (9)	0.4642 (4)	0.0275 (14)	0.37
H27C	0.7178	-0.2082	0.4315	0.033*	0.369 (4)
H27D	0.8398	-0.2491	0.4609	0.033*	0.369 (4)
C28'	0.7223 (7)	-0.2116 (8)	0.5220 (4)	0.0292 (11)	0.37
H28C	0.6947	-0.3007	0.5244	0.035*	0.369 (4)
H28D	0.7801	-0.1972	0.5546	0.035*	0.369 (4)
C29'	0.6248 (5)	-0.1199 (6)	0.5267 (3)	0.0224 (9)	0.37
H29C	0.5644	-0.1395	0.4959	0.027*	0.369 (4)
H29D	0.5925	-0.1308	0.5649	0.027*	0.369 (4)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Sn1	0.01313 (9)	0.01198 (9)	0.01910 (10)	0.00144 (8)	0.00059 (7)	0.00057 (8)
S1	0.0223 (3)	0.0147 (3)	0.0196 (4)	-0.0040 (3)	0.0021 (3)	-0.0018 (3)
S2	0.0371 (4)	0.0163 (4)	0.0304 (5)	-0.0072 (3)	0.0054 (4)	0.0006 (3)
F1	0.0236 (10)	0.0570 (15)	0.0692 (16)	0.0201 (10)	0.0158 (10)	0.0259 (13)
F2	0.0422 (14)	0.109 (2)	0.076 (2)	-0.0369 (16)	-0.0362 (13)	0.0325 (18)
F3	0.0663 (16)	0.0503 (14)	0.0387 (13)	-0.0086 (12)	0.0230 (12)	0.0114 (11)
O1	0.0226 (11)	0.0145 (10)	0.0276 (12)	0.0009 (8)	-0.0042 (9)	-0.0019 (9)
O2	0.0204 (10)	0.0144 (10)	0.0238 (11)	-0.0009 (8)	-0.0049 (8)	0.0033 (9)
N1	0.029 (2)	0.0191 (16)	0.018 (2)	-0.0064 (18)	0.0059 (16)	-0.0010 (15)
N1'	0.029 (2)	0.0191 (16)	0.018 (2)	-0.0064 (18)	0.0059 (16)	-0.0010 (15)

C1	0.0197 (14)	0.0210 (14)	0.0204 (15)	-0.0020 (11)	0.0023 (12)	0.0036 (12)
C2	0.0194 (14)	0.0190 (15)	0.0200 (15)	-0.0025 (11)	0.0038 (11)	0.0052 (12)
C3	0.0199 (15)	0.0256 (16)	0.0275 (17)	-0.0024 (12)	0.0013 (13)	0.0032 (13)
C4	0.0148 (15)	0.046 (2)	0.034 (2)	-0.0019 (14)	0.0005 (13)	0.0118 (16)
C5	0.0184 (15)	0.038 (2)	0.040 (2)	0.0098 (14)	0.0113 (14)	0.0173 (16)
C6	0.0315 (18)	0.0217 (16)	0.041 (2)	0.0036 (13)	0.0173 (16)	0.0063 (15)
C7	0.0202 (15)	0.0235 (16)	0.0282 (17)	-0.0050 (12)	0.0091 (13)	0.0018 (13)
C8	0.0239 (15)	0.0146 (14)	0.0244 (16)	0.0028 (12)	-0.0032 (12)	0.0069 (12)
C9	0.0203 (14)	0.0169 (14)	0.0201 (15)	0.0011 (11)	-0.0004 (12)	0.0063 (12)
C10	0.0258 (16)	0.039 (2)	0.0282 (18)	0.0087 (15)	0.0018 (14)	0.0041 (15)
C11	0.0193 (17)	0.074 (3)	0.046 (2)	0.0072 (18)	-0.0001 (16)	0.018 (2)
C12	0.0280 (19)	0.068 (3)	0.044 (2)	-0.0182 (19)	-0.0180 (17)	0.022 (2)
C13	0.050 (2)	0.033 (2)	0.0270 (18)	-0.0134 (18)	-0.0155 (16)	0.0071 (16)
C14	0.0317 (17)	0.0224 (17)	0.0198 (16)	0.0013 (13)	-0.0029 (13)	0.0035 (12)
C15	0.0153 (13)	0.0154 (15)	0.0257 (16)	0.0013 (11)	0.0031 (11)	-0.0023 (12)
C16	0.0139 (12)	0.0161 (13)	0.0251 (15)	0.0020 (11)	0.0029 (11)	-0.0029 (12)
C17	0.0246 (15)	0.0193 (14)	0.0286 (17)	-0.0020 (12)	0.0016 (13)	-0.0034 (13)
C18	0.0383 (19)	0.034 (2)	0.0250 (18)	-0.0007 (15)	0.0039 (15)	-0.0038 (15)
C19	0.0365 (19)	0.0315 (19)	0.0306 (19)	0.0003 (15)	0.0145 (15)	0.0069 (15)
C20	0.0232 (16)	0.0233 (16)	0.041 (2)	-0.0021 (13)	0.0110 (14)	0.0004 (15)
C21	0.0175 (14)	0.0209 (15)	0.0284 (17)	-0.0005 (12)	0.0036 (12)	-0.0066 (13)
C22	0.0183 (13)	0.0146 (14)	0.0162 (14)	0.0004 (11)	0.0016 (11)	-0.0016 (11)
C23	0.0197 (14)	0.0130 (14)	0.0237 (16)	0.0012 (11)	-0.0007 (12)	-0.0037 (12)
C24	0.025 (3)	0.014 (3)	0.012 (3)	-0.007 (2)	0.004 (2)	-0.001 (2)
C25	0.031 (2)	0.026 (2)	0.026 (2)	-0.007 (2)	0.0096 (17)	-0.001 (2)
C26	0.033 (3)	0.033 (2)	0.016 (3)	-0.006 (3)	0.0051 (18)	-0.002 (2)
C27	0.029 (4)	0.032 (2)	0.022 (3)	-0.007 (3)	0.001 (2)	-0.009 (2)
C28	0.035 (3)	0.025 (2)	0.028 (3)	-0.011 (2)	0.010 (2)	-0.007 (2)
C29	0.033 (2)	0.018 (2)	0.017 (2)	-0.0037 (19)	0.0049 (16)	-0.0008 (17)
C24'	0.034 (6)	0.023 (7)	0.030 (7)	-0.003 (5)	-0.006 (5)	-0.001 (5)
C25'	0.031 (2)	0.026 (2)	0.026 (2)	-0.007 (2)	0.0096 (17)	-0.001 (2)
C26'	0.033 (3)	0.033 (2)	0.016 (3)	-0.006 (3)	0.0051 (18)	-0.002 (2)
C27'	0.029 (4)	0.032 (2)	0.022 (3)	-0.007 (3)	0.001 (2)	-0.009 (2)
C28'	0.035 (3)	0.025 (2)	0.028 (3)	-0.011 (2)	0.010 (2)	-0.007 (2)
C29'	0.033 (2)	0.018 (2)	0.017 (2)	-0.0037 (19)	0.0049 (16)	-0.0008 (17)

*Geometric parameters (Å, °)*

Sn1—C1	2.144 (3)	C13—H13	0.9500
Sn1—C8	2.155 (3)	C14—H14	0.9500
Sn1—C15	2.158 (3)	C15—C16	1.498 (4)
Sn1—O1	2.175 (2)	C15—H15A	0.9900
Sn1—O2 <sup>i</sup>	2.339 (2)	C15—H15B	0.9900
S1—C24'	1.793 (7)	C16—C17	1.392 (4)
S1—C23	1.792 (3)	C16—C21	1.394 (4)
S1—C24	1.796 (5)	C17—C18	1.390 (5)
S2—C24'	1.659 (7)	C17—H17	0.9500
S2—C24	1.662 (5)	C18—C19	1.376 (5)
F1—C5	1.369 (4)	C18—H18	0.9500

## supplementary materials

---

F2—C12	1.362 (4)	C19—C20	1.375 (5)
F3—C19	1.358 (4)	C20—C21	1.385 (5)
O1—C22	1.268 (3)	C20—H20	0.9500
O2—C22	1.241 (3)	C21—H21	0.9500
O2—Sn1 <sup>ii</sup>	2.339 (2)	C22—C23	1.518 (4)
N1—C24	1.333 (6)	C23—H23A	0.9900
N1—C29	1.468 (5)	C23—H23B	0.9900
N1—C25	1.485 (5)	C25—C26	1.490 (6)
N1'—C24'	1.332 (8)	C25—H25A	0.9900
N1'—C29'	1.468 (7)	C25—H25B	0.9900
N1'—C25'	1.479 (7)	C26—C27	1.540 (6)
C1—C2	1.494 (4)	C26—H26A	0.9900
C1—H1A	0.9900	C26—H26B	0.9900
C1—H1B	0.9900	C27—C28	1.530 (6)
C2—C3	1.392 (4)	C27—H27A	0.9900
C2—C7	1.395 (4)	C27—H27B	0.9900
C3—C4	1.386 (4)	C28—C29	1.527 (6)
C3—H3	0.9500	C28—H28A	0.9900
C4—C5	1.369 (5)	C28—H28B	0.9900
C4—H4	0.9500	C29—H29A	0.9900
C5—C6	1.371 (5)	C29—H29B	0.9900
C6—C7	1.389 (4)	C25'—C26'	1.495 (8)
C6—H6	0.9500	C25'—H25C	0.9900
C7—H7	0.9500	C25'—H25D	0.9900
C8—C9	1.499 (4)	C26'—C27'	1.535 (9)
C8—H8A	0.9900	C26'—H26C	0.9900
C8—H8B	0.9900	C26'—H26D	0.9900
C9—C10	1.384 (4)	C27'—C28'	1.525 (8)
C9—C14	1.394 (4)	C27'—H27C	0.9900
C10—C11	1.389 (5)	C27'—H27D	0.9900
C10—H10	0.9500	C28'—C29'	1.507 (8)
C11—C12	1.359 (6)	C28'—H28C	0.9900
C11—H11	0.9500	C28'—H28D	0.9900
C12—C13	1.370 (6)	C29'—H29C	0.9900
C13—C14	1.386 (5)	C29'—H29D	0.9900
C1—Sn1—C8	120.72 (12)	F3—C19—C18	118.8 (3)
C1—Sn1—C15	126.65 (11)	C20—C19—C18	122.6 (3)
C8—Sn1—C15	111.80 (12)	C19—C20—C21	118.1 (3)
C1—Sn1—O1	90.56 (10)	C19—C20—H20	121.0
C8—Sn1—O1	90.20 (10)	C21—C20—H20	121.0
C15—Sn1—O1	98.28 (10)	C20—C21—C16	121.9 (3)
C1—Sn1—O2 <sup>i</sup>	88.16 (10)	C20—C21—H21	119.1
C8—Sn1—O2 <sup>i</sup>	82.46 (10)	C16—C21—H21	119.1
C15—Sn1—O2 <sup>i</sup>	89.94 (9)	O2—C22—O1	123.7 (3)
O1—Sn1—O2 <sup>i</sup>	170.59 (7)	O2—C22—C23	122.8 (3)
C24'—S1—C23	100.5 (2)	O1—C22—C23	113.3 (2)
C23—S1—C24	101.60 (18)	C22—C23—S1	117.0 (2)
C22—O1—Sn1	129.16 (19)	C22—C23—H23A	108.0



C22—O2—Sn1 <sup>ii</sup>	151.17 (19)	S1—C23—H23A	108.0
C24—N1—C29	125.9 (4)	C22—C23—H23B	108.0
C24—N1—C25	122.3 (4)	S1—C23—H23B	108.0
C29—N1—C25	111.8 (4)	H23A—C23—H23B	107.3
C24'—N1'—C29'	125.0 (5)	N1—C24—S2	125.3 (3)
C24'—N1'—C25'	120.6 (6)	N1—C24—S1	114.2 (3)
C29'—N1'—C25'	114.5 (5)	S2—C24—S1	120.5 (3)
C2—C1—Sn1	110.7 (2)	N1—C25—C26	111.6 (3)
C2—C1—H1A	109.5	N1—C25—H25A	109.3
Sn1—C1—H1A	109.5	C26—C25—H25A	109.3
C2—C1—H1B	109.5	N1—C25—H25B	109.3
Sn1—C1—H1B	109.5	C26—C25—H25B	109.3
H1A—C1—H1B	108.1	H25A—C25—H25B	108.0
C3—C2—C7	117.9 (3)	C25—C26—C27	111.1 (4)
C3—C2—C1	121.5 (3)	C25—C26—H26A	109.4
C7—C2—C1	120.5 (3)	C27—C26—H26A	109.4
C4—C3—C2	121.4 (3)	C25—C26—H26B	109.4
C4—C3—H3	119.3	C27—C26—H26B	109.4
C2—C3—H3	119.3	H26A—C26—H26B	108.0
C5—C4—C3	118.3 (3)	C28—C27—C26	109.9 (4)
C5—C4—H4	120.9	C28—C27—H27A	109.7
C3—C4—H4	120.9	C26—C27—H27A	109.7
C4—C5—F1	118.5 (3)	C28—C27—H27B	109.7
C4—C5—C6	123.0 (3)	C26—C27—H27B	109.7
F1—C5—C6	118.5 (3)	H27A—C27—H27B	108.2
C5—C6—C7	117.8 (3)	C29—C28—C27	111.7 (4)
C5—C6—H6	121.1	C29—C28—H28A	109.3
C7—C6—H6	121.1	C27—C28—H28A	109.3
C6—C7—C2	121.5 (3)	C29—C28—H28B	109.3
C6—C7—H7	119.2	C27—C28—H28B	109.3
C2—C7—H7	119.2	H28A—C28—H28B	107.9
C9—C8—Sn1	116.0 (2)	N1—C29—C28	108.8 (3)
C9—C8—H8A	108.3	N1—C29—H29A	109.9
Sn1—C8—H8A	108.3	C28—C29—H29A	109.9
C9—C8—H8B	108.3	N1—C29—H29B	109.9
Sn1—C8—H8B	108.3	C28—C29—H29B	109.9
H8A—C8—H8B	107.4	H29A—C29—H29B	108.3
C10—C9—C14	117.6 (3)	N1'—C24'—S2	125.5 (5)
C10—C9—C8	121.1 (3)	N1'—C24'—S1	113.6 (5)
C14—C9—C8	121.2 (3)	S2—C24'—S1	120.9 (4)
C9—C10—C11	121.5 (4)	N1'—C25'—C26'	110.2 (5)
C9—C10—H10	119.2	N1'—C25'—H25C	109.6
C11—C10—H10	119.2	C26'—C25'—H25C	109.6
C12—C11—C10	118.5 (4)	N1'—C25'—H25D	109.6
C12—C11—H11	120.8	C26'—C25'—H25D	109.6
C10—C11—H11	120.8	H25C—C25'—H25D	108.1
C11—C12—F2	119.1 (4)	C25'—C26'—C27'	110.9 (8)
C11—C12—C13	122.8 (3)	C25'—C26'—H26C	109.5
F2—C12—C13	118.1 (4)	C27'—C26'—H26C	109.5

## supplementary materials

C12—C13—C14	117.9 (4)	C25'—C26'—H26D	109.5
C12—C13—H13	121.0	C27'—C26'—H26D	109.5
C14—C13—H13	121.0	H26C—C26'—H26D	108.1
C13—C14—C9	121.7 (3)	C28'—C27'—C26'	109.0 (8)
C13—C14—H14	119.2	C28'—C27'—H27C	109.9
C9—C14—H14	119.2	C26'—C27'—H27C	109.9
C16—C15—Sn1	110.79 (19)	C28'—C27'—H27D	109.9
C16—C15—H15A	109.5	C26'—C27'—H27D	109.9
Sn1—C15—H15A	109.5	H27C—C27'—H27D	108.3
C16—C15—H15B	109.5	C29'—C28'—C27'	109.2 (8)
Sn1—C15—H15B	109.5	C29'—C28'—H28C	109.8
H15A—C15—H15B	108.1	C27'—C28'—H28C	109.8
C17—C16—C21	117.7 (3)	C29'—C28'—H28D	109.8
C17—C16—C15	120.3 (3)	C27'—C28'—H28D	109.8
C21—C16—C15	122.0 (3)	H28C—C28'—H28D	108.3
C18—C17—C16	121.7 (3)	N1'—C29'—C28'	111.2 (5)
C18—C17—H17	119.2	N1'—C29'—H29C	109.4
C16—C17—H17	119.2	C28'—C29'—H29C	109.4
C19—C18—C17	118.1 (3)	N1'—C29'—H29D	109.4
C19—C18—H18	121.0	C28'—C29'—H29D	109.4
C17—C18—H18	121.0	H29C—C29'—H29D	108.0
F3—C19—C20	118.6 (3)		
C1—Sn1—O1—C22	-83.6 (3)	C19—C20—C21—C16	-0.3 (5)
C8—Sn1—O1—C22	155.7 (3)	C17—C16—C21—C20	0.2 (4)
C15—Sn1—O1—C22	43.6 (3)	C15—C16—C21—C20	-177.9 (3)
C8—Sn1—C1—C2	10.2 (3)	Sn1 <sup>ii</sup> —O2—C22—O1	149.7 (3)
C15—Sn1—C1—C2	178.85 (19)	Sn1 <sup>ii</sup> —O2—C22—C23	-26.1 (6)
O1—Sn1—C1—C2	-80.4 (2)	Sn1—O1—C22—O2	0.8 (4)
O2 <sup>i</sup> —Sn1—C1—C2	90.3 (2)	Sn1—O1—C22—C23	177.03 (18)
Sn1—C1—C2—C3	82.4 (3)	O2—C22—C23—S1	-27.3 (4)
Sn1—C1—C2—C7	-94.3 (3)	O1—C22—C23—S1	156.4 (2)
C7—C2—C3—C4	0.6 (5)	C24'—S1—C23—C22	-80.7 (2)
C1—C2—C3—C4	-176.3 (3)	C24—S1—C23—C22	-70.5 (2)
C2—C3—C4—C5	1.3 (5)	C29—N1—C24—S2	-179.80 (10)
C3—C4—C5—F1	177.8 (3)	C25—N1—C24—S2	0.15 (17)
C3—C4—C5—C6	-2.2 (5)	C29—N1—C24—S1	0.16 (15)
C4—C5—C6—C7	1.2 (5)	C25—N1—C24—S1	-179.89 (8)
F1—C5—C6—C7	-178.9 (3)	C24'—S2—C24—N1	100.1 (16)
C5—C6—C7—C2	0.9 (5)	C24'—S2—C24—S1	-79.9 (16)
C3—C2—C7—C6	-1.7 (4)	C24'—S1—C24—N1	-100.3 (15)
C1—C2—C7—C6	175.2 (3)	C23—S1—C24—N1	174.76 (15)
C1—Sn1—C8—C9	-67.4 (3)	C24'—S1—C24—S2	79.7 (15)
C15—Sn1—C8—C9	122.4 (2)	C23—S1—C24—S2	-5.27 (16)
O1—Sn1—C8—C9	23.4 (2)	C24—N1—C25—C26	-119.8 (4)
O2 <sup>i</sup> —Sn1—C8—C9	-150.7 (2)	C29—N1—C25—C26	60.2 (4)
Sn1—C8—C9—C10	101.0 (3)	N1—C25—C26—C27	-55.1 (6)
Sn1—C8—C9—C14	-76.1 (3)	C25—C26—C27—C28	52.1 (6)
C14—C9—C10—C11	1.9 (5)	C26—C27—C28—C29	-53.8 (6)

C8—C9—C10—C11	-175.3 (3)	C24—N1—C29—C28	120.1 (4)
C9—C10—C11—C12	-0.6 (6)	C25—N1—C29—C28	-59.9 (3)
C10—C11—C12—F2	177.8 (4)	C27—C28—C29—N1	57.6 (5)
C10—C11—C12—C13	-1.1 (6)	C29'—N1'—C24'—S2	-179.98 (10)
C11—C12—C13—C14	1.4 (6)	C25'—N1'—C24'—S2	0.02 (17)
F2—C12—C13—C14	-177.6 (3)	C29'—N1'—C24'—S1	0.01 (15)
C12—C13—C14—C9	0.0 (5)	C25'—N1'—C24'—S1	-179.99 (8)
C10—C9—C14—C13	-1.6 (5)	C24—S2—C24'—N1'	-98.2 (16)
C8—C9—C14—C13	175.6 (3)	C24—S2—C24'—S1	81.8 (16)
C1—Sn1—C15—C16	-169.27 (19)	C23—S1—C24'—N1'	-164.49 (17)
C8—Sn1—C15—C16	0.3 (2)	C24—S1—C24'—N1'	98.5 (15)
O1—Sn1—C15—C16	93.8 (2)	C23—S1—C24'—S2	15.51 (17)
O2 <sup>i</sup> —Sn1—C15—C16	-81.6 (2)	C24—S1—C24'—S2	-81.5 (15)
Sn1—C15—C16—C17	-84.4 (3)	C24'—N1'—C25'—C26'	125.9 (6)
Sn1—C15—C16—C21	93.6 (3)	C29'—N1'—C25'—C26'	-54.1 (6)
C21—C16—C17—C18	-0.1 (5)	N1'—C25'—C26'—C27'	55.3 (9)
C15—C16—C17—C18	178.0 (3)	C25'—C26'—C27'—C28'	-58.9 (10)
C16—C17—C18—C19	0.1 (5)	C26'—C27'—C28'—C29'	58.6 (10)
C17—C18—C19—F3	-179.7 (3)	C24'—N1'—C29'—C28'	-124.7 (6)
C17—C18—C19—C20	-0.3 (5)	C25'—N1'—C29'—C28'	55.3 (6)
F3—C19—C20—C21	179.8 (3)	C27'—C28'—C29'—N1'	-56.7 (9)
C18—C19—C20—C21	0.4 (5)		

Symmetry codes: (i)  $-x+3/2, y+1/2, -z+3/2$ ; (ii)  $-x+3/2, y-1/2, -z+3/2$ .

Fig. 1

