

Potassium clavulanate

Kotaro Fujii,^a Kazuyuki Toyota,^a Akiko Sekine,^a Hidehiro Uekusa,^{a*} Ilma Nugrahani,^b Sukmadjaja Asyarie,^b N. Sundani Soewandhi^b and Slamet Ibrahim^b

^aDepartment of Chemistry and Materials Science, Tokyo Institute of Technology, Ookayama, Meguro, Tokyo 152-8551, Japan, and ^bSchool of Pharmacy, Institut Teknologi Bandung, Ganesha 10, Bandung, 40312, Indonesia
Correspondence e-mail: uekusa@cms.titech.ac.jp


Received 4 July 2010; accepted 14 July 2010

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.044; wR factor = 0.118; data-to-parameter ratio = 15.6.

The title salt, $\text{K}^+\cdot\text{C}_8\text{H}_8\text{NO}_5^-$ [systematic name: potassium (*2R,5R,Z*)-3-(2-hydroxyethylidene)-7-oxo-4-oxa-1-azabicyclo-[3.2.0]heptane-2-carboxylate], a widely used β -lactam antibiotic, is usually chemically unstable even in the solid state owing to its tendency to be hydrolysed. In the crystal structure, the potassium cations are arranged along the *a* axis, forming interactions to the carboxylate and hydroxy groups, resulting in one-dimensional ionic columns. These columns are arranged along the *b* axis, connected by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a layer in the *ab* plane.

Related literature

For the pharmacological activity of clavulanic acid and potassium clavulanate, see: Bird *et al.* (1982); Mayer & Deckwer (1996); Navarro (2005). For the hydrolysis properties of clavulanic acid and potassium clavulanate, see: Bersanetti *et al.* (2005); Brethauer *et al.* (2008); Haginaka *et al.* (1985); Hickey *et al.* (2007); Saudagar *et al.* (2008).



Experimental

Crystal data

$\text{K}^+\cdot\text{C}_8\text{H}_8\text{NO}_5^-$	$V = 934.1(2)\text{ \AA}^3$
$M_r = 237.25$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 4.3453(6)\text{ \AA}$	$\mu = 0.57\text{ mm}^{-1}$
$b = 7.8191(11)\text{ \AA}$	$T = 173\text{ K}$
$c = 27.491(3)\text{ \AA}$	$0.24 \times 0.04 \times 0.01\text{ mm}$

Data collection

Rigaku R-AXIS RAPID IP area-detector diffractometer	9047 measured reflections
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	2138 independent reflections
$T_{\min} = 0.876$, $T_{\max} = 0.994$	1433 reflections with $I > 2\sigma(I)$
$R_{\text{int}} = 0.088$	$R_{\text{int}} = 0.088$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$	$\Delta\rho_{\text{max}} = 0.52\text{ e \AA}^{-3}$
$wR(F^2) = 0.118$	$\Delta\rho_{\text{min}} = -0.56\text{ e \AA}^{-3}$
$S = 1.12$	Absolute structure: Flack (1983), 839 Friedel pairs
2138 reflections	Flack parameter: -0.05 (9)
137 parameters	
H-atom parameters constrained	

Table 1
Selected bond lengths (\AA).

O2—K1 ⁱ	2.773 (3)	O3—K1	2.786 (3)
O2—K1 ⁱⁱ	2.799 (3)	O4—K1 ⁱⁱ	2.818 (4)
O2—K1	2.827 (3)	O4—K1 ⁱⁱⁱ	2.865 (4)

Symmetry codes: (i) $x + 1, y, z$; (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, -z + 2$; (iii) $x + \frac{3}{2}, -y + \frac{3}{2}, -z + 2$.

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O4—H4A \cdots O3 ^{iv}	0.84	1.90	2.673 (5)	153

Symmetry code: (iv) $x + 1, y + 1, z$.

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *PROCESS-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Grateful thanks are given to the High Directorate of the Education Ministry of Indonesia for a research grant and PT Tempo Scan Pacific Indonesia for the supply of the material. KF was supported by a Grant-in-Aid for JSPS Fellows (Tokyo Institute of Technology G-COE Program: Education and the Research Centre for Emergence of New Molecular Chemistry).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2689).

References

- Bersanetti, P. A., Almeida, R. M. R. G., Barboza, M., Araújo, M. L. G. C. & Hokka, C. O. (2005). *Biochem. Eng. J.* **23**, 31–36.
- Bird, A. E., Bellis, J. M. & Basson, B. C. (1982). *Analyst*, **107**, 1241–1245.
- Brethauer, S., Held, M. & Panke, S. (2008). *Biotechnol. Bioeng.* **100**, 439–447.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Haginaka, J., Yasuda, H., Uno, T. & Nakagawa, T. (1985). *Chem. Pharm. Bull.* **33**, 218–224.
- Hickey, M. B., Peterson, M. L., Manas, E. S., Alvarez, J., Haeffner, F. & Almarsson, O. (2007). *J. Pharm. Sci.* **96**, 1090–1099.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.

metal-organic compounds

- Mayer, A. F. & Deckwer, W. D. (1996). *Appl. Microbiol. Biotechnol.* **45**, 41–46.
Navarro, A. S. (2005). *Clin. Pharmacokinet.* **44**, 1097–1115.
Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Saudagar, P. S., Survase, S. A. & Singhal, R. S. (2008). *Biotechnol. Adv.* **26**, 335–351.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64** 112–122.

supplementary materials

Acta Cryst. (2010). E66, m985-m986 [doi:10.1107/S1600536810027984]

Potassium clavulanate

K. Fujii, K. Toyota, A. Sekine, H. Uekusa, I. Nugrahani, S. Asyarie, N. S. Soewandhi and S. Ibrahim

Comment

Many pathogenic bacteria secrete β -lactamases as a defense mechanism against β -lactam antibiotics. Because such β -lactamases have the potential to inactivate β -lactam antibiotics, inhibitors for these β -lactamases are clinically very important. Clavulanic acid (CA) is a powerful naturally obtained inhibitor for bacterial β -lactamases produced by the organism *Streptomyces clavuligerus*. Although CA itself can act as an β -lactam antibiotic and is active against a wide spectrum of Gram-positive and Gram-negative bacteria (Mayer & Deckwer, 1996), it is much more effective as a drug in combination with β -lactamase-sensitive penicillins, such as amoxicillin. In that situation, CA protects the β -lactam ring of the amoxicillin from hydrolysis and can maintain its activity against β -lactamase producing bacteria (Bird *et al.*, 1982). The CA potassium salt is widely used as a drug in injectable and solid form, especially combined with amoxicillin sodium and amoxicillin trihydrate (Navarro, 2005).

In this context, an understanding of the structure of CA is important in order to establish its ability to form molecular interactions. Unfortunately, CA is chemically unstable as are the other β -lactam antibiotics, being very sensitive to pH, temperature, and humidity *via* the hydrolysis degradation mechanism; (Bersanetti *et al.*, 2005; Hickey *et al.*, 2007; Saudagar *et al.*, 2008). The decomposition is also self-catalyzed (Brethauer *et al.* 2008; Haginaka *et al.* 1985) and there have been some difficulties in obtaining a single crystals of CA. Therefore, until now there has been no report of a crystal structure of CA. In this study, single crystals of potassium clavulanate were successfully obtained by a low-temperature crystallization process and the crystal structure was determined.

In the molecular structure of potassium clavulanate, Fig. 1, the C5–O2 and C5–O3 distances of 1.262 (5) \AA and 1.256 (5) \AA , respectively, indicate that the negative charge of the carboxylate group is delocalised. The potassium cation is surrounded by six oxygen atoms, three O2, one O3 and two O4, derived from four different clavulanate anions. The selected bond lengths around the potassium cation are listed in Table 1. These interactions are infinitely linked along the *a* axis and lead to an ionic (hydrophilic) column structure. These columns are connected by intermolecular O–H \cdots O hydrogen bonds formed between O4-hydroxyl groups and carboxylate-O2 atoms, and form a hydrophobic layer in the *ab* plane; Fig. 2. By contrast, the remaining hydrophobic groups (*i.e.* bicyclo groups) form a hydrophobic layer so that the crystal structure comprises alternating hydrophilic and hydrophilic regions.

Experimental

The single crystals of potassium clavulanate were grown at a low temperature in order to prevent decomposition. After the compound was dissolved into an 8:2 mixture of methanol/water, a few drops of 1-propanol were added to the solution and the solution was kept at 235 K for a few days. The crystal used in the analysis was immediately covered with inert oil in order to prevent the decomposition through contact with atmospheric water vapor.

supplementary materials

Refinement

The O- and C-bound H atoms were geometrically placed ($O-H = 0.84 \text{ \AA}$ and $C-H = 0.95-1.00 \text{ \AA}$) and refined as riding with $U_{iso}(H) = 1.2U_{eq}(\text{carrier atom})$. The absolute structure was assigned according to the known configuration of the acid, an assignment confirmed by the refinement of the Flack parameter (Flack, 1983).

Figures




Fig. 1. Molecular structure of potassium clavulanate showing numbering scheme and displacement ellipsoids at the 50% probability .



Fig. 2. Crystal packing viewed along the a axis. The hydrogen bonds are shown with red dotted lines.

potassium (2*R*,5*R*,*Z*)- 3-(2-hydroxyethylidene)-7-oxo-4-oxa-1-azabicyclo[3.2.0]heptane-2-carboxylate

Crystal data

$K^+\cdot C_8H_8NO_5^-$	$F(000) = 488$
$M_r = 237.25$	$D_x = 1.687 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation, $\lambda = 0.71075 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 9047 reflections
$a = 4.3453 (6) \text{ \AA}$	$\theta = 3.0-27.5^\circ$
$b = 7.8191 (11) \text{ \AA}$	$\mu = 0.57 \text{ mm}^{-1}$
$c = 27.491 (3) \text{ \AA}$	$T = 173 \text{ K}$
$V = 934.1 (2) \text{ \AA}^3$	Platet, colorless
$Z = 4$	$0.24 \times 0.04 \times 0.01 \text{ mm}$

Data collection

Rigaku R-AXIS RAPID IP area-detector diffractometer	2138 independent reflections
Radiation source: rotating anode, Rigaku UltraX18 graphite	1433 reflections with $I > 2\sigma(I)$
ω scan	$R_{int} = 0.088$
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	$\theta_{\max} = 27.5^\circ, \theta_{\min} = 3.0^\circ$
$T_{\min} = 0.876, T_{\max} = 0.994$	$h = -5 \rightarrow 5$
9047 measured reflections	$k = -10 \rightarrow 10$
	$l = -35 \rightarrow 35$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	H-atom parameters constrained
$wR(F^2) = 0.118$	$w = 1/[\sigma^2(F_o^2) + (0.0305P)^2 + 0.8726P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.12$	$(\Delta/\sigma)_{\max} = 0.001$
2138 reflections	$\Delta\rho_{\max} = 0.52 \text{ e \AA}^{-3}$
137 parameters	$\Delta\rho_{\min} = -0.56 \text{ e \AA}^{-3}$
0 restraints	Absolute structure: Flack (1983), 839 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: -0.05 (9)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.2101 (11)	0.3967 (6)	1.20066 (15)	0.0297 (11)
C2	1.1168 (11)	0.5309 (6)	1.23813 (13)	0.0276 (9)
H2A	1.2901	0.5798	1.2569	0.033*
H2B	0.9447	0.4964	1.2596	0.033*
C3	1.0199 (11)	0.6369 (6)	1.19340 (14)	0.0250 (10)
H3	0.8027	0.6790	1.1942	0.030*
C4	1.2417 (9)	0.5363 (6)	1.11851 (13)	0.0192 (8)
H4	1.4273	0.4611	1.1163	0.023*
C5	1.0676 (9)	0.5213 (6)	1.06982 (13)	0.0201 (8)
C6	1.3526 (11)	0.7148 (5)	1.13197 (12)	0.0218 (9)
C7	1.5398 (10)	0.8175 (6)	1.10754 (14)	0.0245 (10)
H7	1.6169	0.7752	1.0775	0.029*
C8	1.6419 (11)	0.9914 (5)	1.12198 (13)	0.0269 (9)
H8A	1.5056	1.0382	1.1476	0.032*
H8B	1.8551	0.9881	1.1346	0.032*
O1	1.3545 (10)	0.2661 (4)	1.20053 (11)	0.0458 (9)
O2	1.1415 (8)	0.6217 (4)	1.03630 (9)	0.0258 (7)

supplementary materials

O3	0.8682 (9)	0.4058 (4)	1.06647 (10)	0.0318 (7)
O4	1.6262 (9)	1.0947 (4)	1.07905 (10)	0.0321 (8)
H4A	1.6958	1.1926	1.0851	0.038*
O5	1.2401 (7)	0.7604 (4)	1.17783 (9)	0.0266 (7)
N1	1.0715 (8)	0.4860 (5)	1.16197 (10)	0.0242 (8)
K1	0.6334 (2)	0.52404 (13)	0.97836 (3)	0.0284 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.036 (3)	0.027 (2)	0.026 (2)	-0.001 (2)	0.001 (2)	0.0026 (19)
C2	0.029 (2)	0.035 (2)	0.0185 (16)	-0.001 (3)	-0.0026 (18)	0.0049 (18)
C3	0.026 (2)	0.025 (2)	0.024 (2)	0.0001 (19)	0.0021 (19)	0.0033 (18)
C4	0.0183 (18)	0.018 (2)	0.0211 (17)	-0.0025 (18)	-0.0009 (15)	0.0023 (17)
C5	0.017 (2)	0.022 (2)	0.0214 (17)	-0.0007 (18)	0.0031 (15)	-0.0040 (17)
C6	0.020 (2)	0.028 (2)	0.0174 (17)	0.001 (2)	-0.0018 (18)	-0.0019 (16)
C7	0.027 (2)	0.026 (2)	0.0204 (18)	-0.0043 (19)	0.0037 (18)	0.0005 (17)
C8	0.030 (2)	0.025 (2)	0.0255 (18)	-0.003 (2)	-0.002 (2)	0.0008 (18)
O1	0.070 (3)	0.033 (2)	0.0340 (16)	0.020 (2)	0.006 (2)	0.0063 (15)
O2	0.0265 (16)	0.0296 (17)	0.0213 (12)	-0.0041 (16)	-0.0002 (13)	0.0045 (12)
O3	0.0366 (19)	0.0304 (17)	0.0285 (14)	-0.0127 (16)	-0.0071 (15)	0.0038 (13)
O4	0.046 (2)	0.0201 (15)	0.0304 (15)	-0.0063 (17)	-0.0020 (17)	0.0034 (12)
O5	0.0328 (18)	0.0248 (17)	0.0221 (13)	-0.0045 (13)	0.0064 (13)	-0.0013 (13)
N1	0.032 (2)	0.0236 (19)	0.0174 (14)	0.0030 (17)	0.0036 (14)	0.0043 (14)
K1	0.0242 (4)	0.0362 (5)	0.0248 (4)	0.0008 (5)	-0.0007 (4)	0.0031 (4)

Geometric parameters (\AA , $^\circ$)

C1—O1	1.199 (6)	C7—H7	0.9500
C1—N1	1.408 (5)	C8—O4	1.432 (5)
C1—C2	1.525 (6)	C8—H8A	0.9900
C2—C3	1.542 (5)	C8—H8B	0.9900
C2—H2A	0.9900	O4—H4A	0.8400
C2—H2B	0.9900	O2—K1 ⁱ	2.773 (3)
C3—O5	1.425 (5)	O2—K1 ⁱⁱ	2.799 (3)
C3—N1	1.480 (5)	O2—K1	2.827 (3)
C3—H3	1.0000	O3—K1	2.786 (3)
C4—N1	1.459 (5)	O4—K1 ⁱⁱ	2.818 (4)
C4—C6	1.522 (6)	O4—K1 ⁱⁱⁱ	2.865 (4)
C4—C5	1.542 (5)	O4—H4A	0.8400
C4—H4	1.0000	K1—O2 ^{iv}	2.773 (3)
C5—O2	1.252 (5)	K1—O2 ^v	2.799 (3)
C5—O3	1.256 (5)	K1—O4 ^v	2.818 (4)
C6—C7	1.325 (6)	K1—O4 ^{vi}	2.865 (4)
C6—O5	1.398 (4)	K1—C7 ^v	3.198 (4)
C7—C8	1.485 (6)		
O1—C1—N1	130.1 (4)	C1—N1—C4	122.4 (4)

O1—C1—C2	136.7 (4)	C1—N1—C3	91.1 (3)
N1—C1—C2	93.2 (3)	C4—N1—C3	109.9 (3)
C1—C2—C3	84.5 (3)	O2 ^{iv} —K1—O3	82.78 (10)
C1—C2—H2A	114.6	O2 ^{iv} —K1—O2 ^v	79.62 (9)
C3—C2—H2A	114.6	O3—K1—O2 ^v	116.68 (9)
C1—C2—H2B	114.6	O2 ^{iv} —K1—O4 ^v	176.73 (10)
C3—C2—H2B	114.6	O3—K1—O4 ^v	95.70 (10)
H2A—C2—H2B	111.7	O2 ^v —K1—O4 ^v	103.64 (9)
O5—C3—N1	105.3 (3)	O2 ^{iv} —K1—O2	101.77 (9)
O5—C3—C2	114.9 (4)	O3—K1—O2	46.69 (9)
N1—C3—C2	89.7 (3)	O2 ^v —K1—O2	78.70 (9)
O5—C3—H3	114.7	O4 ^v —K1—O2	79.11 (9)
N1—C3—H3	114.7	O2 ^{iv} —K1—O4 ^{vi}	79.21 (9)
C2—C3—H3	114.7	O3—K1—O4 ^{vi}	130.76 (10)
N1—C4—C6	102.0 (3)	O2 ^v —K1—O4 ^{vi}	104.54 (9)
N1—C4—C5	116.2 (3)	O4 ^v —K1—O4 ^{vi}	99.74 (9)
C6—C4—C5	115.9 (3)	O2—K1—O4 ^{vi}	176.76 (10)
N1—C4—H4	107.4	O2 ^{iv} —K1—C5	90.29 (9)
C6—C4—H4	107.4	O3—K1—C5	23.47 (10)
C5—C4—H4	107.4	O2 ^v —K1—C5	96.56 (11)
O2—C5—O3	125.0 (3)	O4 ^v —K1—C5	89.41 (10)
O2—C5—C4	117.7 (4)	O2—K1—C5	23.45 (9)
O3—C5—C4	117.2 (3)	O4 ^{vi} —K1—C5	154.16 (10)
O2—C5—K1	64.0 (2)	O2 ^{iv} —K1—C7 ^v	137.98 (11)
O3—C5—K1	62.1 (2)	O3—K1—C7 ^v	124.63 (12)
C4—C5—K1	171.2 (3)	O2 ^v —K1—C7 ^v	60.23 (10)
O2—C5—K1 ⁱ	44.8 (2)	O4 ^v —K1—C7 ^v	45.13 (10)
O3—C5—K1 ⁱ	115.7 (3)	O2—K1—C7 ^v	83.12 (10)
C4—C5—K1 ⁱ	106.1 (2)	O4 ^{vi} —K1—C7 ^v	98.25 (10)
K1—C5—K1 ⁱ	81.24 (8)	C5—K1—C7 ^v	105.17 (11)
C7—C6—O5	121.1 (4)	O2 ^{iv} —K1—C8 ^{vi}	89.53 (10)
C7—C6—C4	128.8 (4)	O3—K1—C8 ^{vi}	154.09 (11)
O5—C6—C4	110.0 (3)	O2 ^v —K1—C8 ^{vi}	85.89 (9)
C6—C7—C8	127.1 (4)	O4 ^v —K1—C8 ^{vi}	90.62 (10)
C6—C7—K1 ⁱⁱ	105.7 (3)	O2—K1—C8 ^{vi}	158.79 (11)
C8—C7—K1 ⁱⁱ	90.4 (2)	O4 ^{vi} —K1—C8 ^{vi}	23.49 (9)
C6—C7—H7	116.5	C5—K1—C8 ^{vi}	177.48 (12)
C8—C7—H7	116.5	C7 ^v —K1—C8 ^{vi}	76.57 (11)
K1 ⁱⁱ —C7—H7	71.9	O2 ^{iv} —K1—C5 ^{iv}	18.57 (9)
O4—C8—C7	106.4 (3)	O3—K1—C5 ^{iv}	68.43 (10)
O4—C8—K1 ⁱⁱⁱ	52.9 (2)	O2 ^v —K1—C5 ^{iv}	96.76 (10)

supplementary materials

C7—C8—K1 ⁱⁱⁱ	86.5 (2)	O4 ^v —K1—C5 ^{iv}	158.42 (10)
O4—C8—K1 ⁱⁱ	49.2 (2)	O2—K1—C5 ^{iv}	98.32 (9)
C7—C8—K1 ⁱⁱ	64.7 (2)	O4 ^{vi} —K1—C5 ^{iv}	81.62 (9)
K1 ⁱⁱⁱ —C8—K1 ⁱⁱ	76.39 (8)	C5—K1—C5 ^{iv}	81.24 (8)
O4—C8—H8A	110.5	C7 ^v —K1—C5 ^{iv}	156.37 (11)
C7—C8—H8A	110.5	C8 ^{vi} —K1—C5 ^{iv}	97.88 (10)
K1 ⁱⁱⁱ —C8—H8A	160.2	O2 ^{iv} —K1—C8 ^v	159.90 (10)
K1 ⁱⁱ —C8—H8A	101.1	O3—K1—C8 ^v	115.98 (12)
O4—C8—H8B	110.5	O2 ^v —K1—C8 ^v	85.05 (9)
C7—C8—H8B	110.5	O4 ^v —K1—C8 ^v	22.62 (9)
K1 ⁱⁱⁱ —C8—H8B	72.8	O2—K1—C8 ^v	87.75 (10)
K1 ⁱⁱ —C8—H8B	149.1	O4 ^{vi} —K1—C8 ^v	92.28 (10)
H8A—C8—H8B	108.6	C5—K1—C8 ^v	104.40 (10)
C5—O2—K1 ⁱ	116.6 (3)	C7 ^v —K1—C8 ^v	24.83 (10)
C5—O2—K1 ⁱⁱ	136.3 (3)	C8 ^{vi} —K1—C8 ^v	76.39 (8)
K1 ⁱ —O2—K1 ⁱⁱ	101.51 (10)	C5 ^{iv} —K1—C8 ^v	173.89 (10)
C5—O2—K1	92.6 (2)	O2 ^{iv} —K1—K1 ⁱⁱ	90.24 (7)
K1 ⁱ —O2—K1	101.77 (9)	O3—K1—K1 ⁱⁱ	81.25 (7)
K1 ⁱⁱ —O2—K1	100.16 (10)	O2 ^v —K1—K1 ⁱⁱ	39.02 (7)
C5—O3—K1	94.5 (2)	O4 ^v —K1—K1 ⁱⁱ	92.38 (7)
C8—O4—K1 ⁱⁱ	108.2 (2)	O2—K1—K1 ⁱⁱ	39.68 (6)
C8—O4—K1 ⁱⁱⁱ	103.6 (3)	O4 ^{vi} —K1—K1 ⁱⁱ	143.56 (7)
K1 ⁱⁱ —O4—K1 ⁱⁱⁱ	99.74 (9)	C5—K1—K1 ⁱⁱ	58.85 (8)
C8—O4—H4A	109.5	C7 ^v —K1—K1 ⁱⁱ	66.94 (8)
K1 ⁱⁱ —O4—H4A	133.2	C8 ^{vi} —K1—K1 ⁱⁱ	123.66 (8)
K1 ⁱⁱⁱ —O4—H4A	97.3	C5 ^{iv} —K1—K1 ⁱⁱ	99.19 (7)
C6—O5—C3	109.4 (3)	C8 ^v —K1—K1 ⁱⁱ	85.90 (8)
O1—C1—C2—C3	168.2 (6)	C5—O2—K1—O3	-6.0 (2)
N1—C1—C2—C3	-9.2 (3)	K1 ⁱ —O2—K1—O3	111.90 (16)
C1—C2—C3—O5	-98.0 (4)	K1 ⁱⁱ —O2—K1—O3	-143.96 (16)
C1—C2—C3—N1	8.8 (3)	C5—O2—K1—O2 ^v	138.9 (2)
N1—C4—C5—O2	151.2 (4)	K1 ⁱ —O2—K1—O2 ^v	-103.25 (11)
C6—C4—C5—O2	31.4 (5)	K1 ⁱⁱ —O2—K1—O2 ^v	0.89 (9)
N1—C4—C5—O3	-30.8 (6)	C5—O2—K1—O4 ^v	-114.7 (3)
C6—C4—C5—O3	-150.6 (4)	K1 ⁱ —O2—K1—O4 ^v	3.21 (10)
N1—C4—C5—K1	52 (2)	K1 ⁱⁱ —O2—K1—O4 ^{vi}	107.35 (10)
C6—C4—C5—K1	-67.5 (19)	C5—O2—K1—O4 ^{vi}	-45.1 (17)
N1—C4—C5—K1 ⁱ	-161.8 (3)	K1 ⁱ —O2—K1—O4 ^{vi}	72.8 (16)
C6—C4—C5—K1 ⁱ	78.4 (4)	K1 ⁱⁱ —O2—K1—O4 ^{vi}	177 (45)
N1—C4—C6—C7	174.2 (4)	K1 ⁱ —O2—K1—C5	117.9 (3)
C5—C4—C6—C7	-58.6 (6)	K1 ⁱⁱ —O2—K1—C5	-138.0 (3)

N1—C4—C6—O5	-2.6 (4)	C5—O2—K1—C7 ^v	-160.2 (3)
C5—C4—C6—O5	124.7 (4)	K1 ⁱ —O2—K1—C7 ^v	-42.33 (11)
O5—C6—C7—C8	-3.7 (7)	K1 ⁱⁱ —O2—K1—C7 ^v	61.82 (11)
C4—C6—C7—C8	179.9 (4)	C5—O2—K1—C8 ^{vi}	-176.9 (3)
O5—C6—C7—K1 ⁱⁱ	-106.5 (4)	K1 ⁱ —O2—K1—C8 ^{vi}	-59.1 (3)
C4—C6—C7—K1 ⁱⁱ	77.1 (5)	K1 ⁱⁱ —O2—K1—C8 ^{vi}	45.1 (3)
C6—C7—C8—O4	-137.0 (5)	C5—O2—K1—C5 ^{iv}	43.6 (2)
K1 ⁱⁱ —C7—C8—O4	-26.9 (3)	K1 ⁱ —O2—K1—C5 ^{iv}	161.47 (10)
C6—C7—C8—K1 ⁱⁱⁱ	173.3 (5)	K1 ⁱⁱ —O2—K1—C5 ^{iv}	-94.39 (11)
K1 ⁱⁱ —C7—C8—K1 ⁱⁱⁱ	-76.57 (10)	C5—O2—K1—C8 ^v	-135.7 (3)
C6—C7—C8—K1 ⁱⁱ	-110.1 (5)	K1 ⁱ —O2—K1—C8 ^v	-17.85 (11)
O3—C5—O2—K1 ⁱ	-92.7 (4)	K1 ⁱⁱ —O2—K1—C8 ^v	86.29 (10)
C4—C5—O2—K1 ⁱ	85.1 (4)	C5—O2—K1—K1 ⁱⁱ	138.0 (3)
K1—C5—O2—K1 ⁱ	-104.54 (19)	K1 ⁱ —O2—K1—K1 ⁱⁱ	-104.15 (11)
O3—C5—O2—K1 ⁱⁱ	119.4 (4)	O2—C5—K1—O2 ^{iv}	-120.1 (3)
C4—C5—O2—K1 ⁱⁱ	-62.7 (5)	O3—C5—K1—O2 ^{iv}	70.9 (3)
K1—C5—O2—K1 ⁱⁱ	107.6 (3)	C4—C5—K1—O2 ^{iv}	-16.8 (18)
K1 ⁱ —C5—O2—K1 ⁱⁱ	-147.8 (5)	K1 ⁱ —C5—K1—O2 ^{iv}	-163.73 (9)
O3—C5—O2—K1	11.8 (5)	O2—C5—K1—O3	169.1 (4)
C4—C5—O2—K1	-170.3 (3)	C4—C5—K1—O3	-87.6 (18)
K1 ⁱ —C5—O2—K1	104.54 (19)	K1 ⁱ —C5—K1—O3	125.4 (3)
O2—C5—O3—K1	-12.0 (5)	O2—C5—K1—O2 ^v	-40.5 (2)
C4—C5—O3—K1	170.1 (3)	O3—C5—K1—O2 ^v	150.5 (3)
K1 ⁱ —C5—O3—K1	-63.5 (2)	C4—C5—K1—O2 ^v	62.8 (18)
C7—C8—O4—K1 ⁱⁱ	32.7 (4)	K1 ⁱ —C5—K1—O2 ^v	-84.14 (9)
K1 ⁱⁱⁱ —C8—O4—K1 ⁱⁱ	105.23 (18)	O2—C5—K1—O4 ^v	63.2 (2)
C7—C8—O4—K1 ⁱⁱⁱ	-72.5 (3)	O3—C5—K1—O4 ^v	-105.9 (3)
K1 ⁱⁱ —C8—O4—K1 ⁱⁱⁱ	-105.23 (18)	C4—C5—K1—O4 ^v	166.5 (18)
C7—C6—O5—C3	173.7 (4)	K1 ⁱ —C5—K1—O4 ^v	19.52 (9)
C4—C6—O5—C3	-9.2 (4)	O3—C5—K1—O2	-169.1 (4)
N1—C3—O5—C6	16.9 (4)	C4—C5—K1—O2	103.3 (19)
C2—C3—O5—C6	113.9 (4)	K1 ⁱ —C5—K1—O2	-43.7 (2)
O1—C1—N1—C4	-53.6 (7)	O2—C5—K1—O4 ^{vi}	174.7 (2)
C2—C1—N1—C4	124.1 (4)	O3—C5—K1—O4 ^{vi}	5.7 (4)
O1—C1—N1—C3	-168.1 (6)	C4—C5—K1—O4 ^{vi}	-82.0 (19)
C2—C1—N1—C3	9.6 (3)	K1 ⁱ —C5—K1—O4 ^{vi}	131.1 (2)
C6—C4—N1—C1	-91.7 (5)	O2—C5—K1—C7 ^v	20.4 (3)
C5—C4—N1—C1	141.3 (4)	O3—C5—K1—C7 ^v	-148.7 (3)
C6—C4—N1—C3	13.0 (4)	C4—C5—K1—C7 ^v	123.7 (18)
C5—C4—N1—C3	-114.0 (4)	K1 ⁱ —C5—K1—C7 ^v	-23.27 (12)
O5—C3—N1—C1	106.3 (3)	O2—C5—K1—C8 ^{vi}	154 (2)

supplementary materials

C2—C3—N1—C1	−9.5 (3)	O3—C5—K1—C8 ^{vi}	−15 (2)
O5—C3—N1—C4	−18.9 (4)	C4—C5—K1—C8 ^{vi}	−103 (3)
C2—C3—N1—C4	−134.7 (3)	K1 ⁱ —C5—K1—C8 ^{vi}	110 (2)
C5—O3—K1—O2 ^{iv}	−107.7 (3)	O2—C5—K1—C8 ^{iv}	−136.3 (2)
C5—O3—K1—O2 ^v	−33.2 (3)	O3—C5—K1—C8 ^{iv}	54.6 (3)
C5—O3—K1—O4 ^v	75.2 (3)	C4—C5—K1—C8 ^{iv}	−33.0 (19)
C5—O3—K1—O2	6.0 (2)	K1 ⁱ —C5—K1—C8 ^{iv}	180.0
C5—O3—K1—O4 ^{vi}	−176.7 (2)	O2—C5—K1—C8 ^v	46.1 (3)
C5—O3—K1—C7 ^v	37.6 (3)	O3—C5—K1—C8 ^v	−123.0 (3)
C5—O3—K1—C8 ^{vi}	178.5 (2)	C4—C5—K1—C8 ^v	149.4 (18)
C5—O3—K1—C5 ^{iv}	−120.0 (3)	K1 ⁱ —C5—K1—C8 ^v	2.41 (12)
C5—O3—K1—C8 ^v	64.7 (3)	O2—C5—K1—K1 ⁱⁱ	−30.0 (2)
C5—O3—K1—K1 ⁱⁱ	−16.4 (2)	O3—C5—K1—K1 ⁱⁱ	161.0 (3)
C5—O2—K1—O2 ^{iv}	62.1 (3)	C4—C5—K1—K1 ⁱⁱ	73.4 (18)
K1 ⁱ —O2—K1—O2 ^{iv}	180.0	K1 ⁱ —C5—K1—K1 ⁱⁱ	−73.62 (7)
K1 ⁱⁱ —O2—K1—O2 ^{iv}	−75.85 (11)		


Symmetry codes: (i) $x+1, y, z$; (ii) $x+1/2, -y+3/2, -z+2$; (iii) $x+3/2, -y+3/2, -z+2$; (iv) $x-1, y, z$; (v) $x-1/2, -y+3/2, -z+2$; (vi) $x-3/2, -y+3/2, -z+2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O4—H4A \cdots O3 ^{vii}	0.84	1.90	2.673 (5)	153

Symmetry codes: (vii) $x+1, y+1, z$.

Fig. 1



supplementary materials

Fig. 2

