

## Diethyl 2,2'-(biphenyl-2,2'-diyldioxy)-diacetate

Qamar Ali,<sup>a</sup> Itrat Anis,<sup>b</sup> Donald VanDerveer<sup>c</sup> and Muhammad Raza Shah<sup>a\*</sup>

<sup>a</sup>H.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan, <sup>b</sup>Department of Chemistry, University of Karachi, Karachi 75270, Pakistan, and <sup>c</sup>Chemistry Department, Clemson University, Clemson, SC 29634-0973, USA  
Correspondence e-mail: raza\_shahm@yahoo.com


Received 16 June 2010; accepted 7 July 2010

Key indicators: single-crystal X-ray study;  $T = 153\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.039;  $wR$  factor = 0.099; data-to-parameter ratio = 13.2.

In the title compound,  $\text{C}_{20}\text{H}_{22}\text{O}_6$ , the mean planes through the benzene rings make a dihedral angle of  $59.82(7)^\circ$  with each other. Weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  interactions together with  $\pi-\pi$  stacking interactions [centroid–centroid distance =  $3.830(1)\text{ \AA}$ ] between benzene rings are observed in the crystal packing.

### Related literature

For related structures, see: Rabnawaz *et al.* (2008); Ali *et al.* (2008); Ibad *et al.* (2008).



### Experimental

#### Crystal data

$\text{C}_{20}\text{H}_{22}\text{O}_6$

$M_r = 358.38$

Triclinic,  $P\bar{1}$   
 $a = 7.4683(13)\text{ \AA}$   
 $b = 10.8189(15)\text{ \AA}$   
 $c = 12.050(2)\text{ \AA}$   
 $\alpha = 104.733(17)^\circ$   
 $\beta = 95.05(2)^\circ$   
 $\gamma = 106.897(14)^\circ$

$V = 887.1(3)\text{ \AA}^3$   
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.10\text{ mm}^{-1}$   
 $T = 153\text{ K}$   
 $0.58 \times 0.53 \times 0.24\text{ mm}$

#### Data collection

Rigaku AFC-8S diffractometer  
Absorption correction: multi-scan  
(*REQAB*; Jacobson, 1998)  
 $T_{\min} = 0.945$ ,  $T_{\max} = 0.977$

6640 measured reflections  
3104 independent reflections  
2787 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.014$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$   
 $wR(F^2) = 0.099$   
 $S = 1.04$   
3104 reflections

235 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.20\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C4—H4b $\cdots$ O1 <sup>i</sup>	0.99	2.50	3.472 (2)	168 (1)

Symmetry code: (i)  $-x + 1, -y + 1, -z$ .

Data collection: *CrystalClear* (Rigaku/MSC, 2006); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors thank the Pakistan Science Foundation for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZ2221).

### References

- Ali, Q., Ibad, F., Shah, M. R. & VanDerveer, D. (2008). *Acta Cryst. E* **64**, o1408.  
Ibad, F., Mustafa, A., Shah, M. R. & VanDerveer, D. (2008). *Acta Cryst. E* **64**, o1130–o1131.  
Jacobson, R. (1998). *REQAB*. Molecular Structure Corporation, The Woodlands, Texas, USA.  
Rabnawaz, M., Ali, Q., Shah, M. R. & Singh, K. (2008). *Acta Cryst. E* **64**, o1909.  
Rigaku/MSC (2006). *CrystalClear* MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

## **supplementary materials**

*Acta Cryst.* (2010). E66, o1984 [doi:10.1107/S1600536810026863]

## Diethyl 2,2'-(biphenyl-2,2'-diyldioxy)diacetate

**Q. Ali, I. Anis, D. VanDerveer and M. R. Shah**

### Comment

In the continuation of our work on biphenyl derivatives (Rabnawaz *et al.*, 2008) we report here the synthesis and crystal structure of the title compound, I. The molecular structure and atom-numbering scheme of the title compound are shown in Fig. 1. The molecules are essentially non-planar, with a dihedral angle of 59.82 (7) $^{\circ}$  between the mean planes through the benzene rings. This is slightly more than the corresponding tert butyl ester analogue of I (Ali *et al.*, 2008), due to the presence of the sterically less crowded ethyl ester group. The key C=O and C—O bond distances are in agreement with those observed in the related hydrazide structure (Ibad *et al.*, 2008). NMR data show that the molecule has a non-crystallographic two-fold rotation axis.

### Experimental

$\text{K}_2\text{CO}_3$  (414 mg, 3 mmol) and 2,2'-dihydroxybiphenyl (186 mg, 1 mmol) in 15 ml of acetone were stirred for 10 minutes, followed by addition of ethyl bromoacetate (314 mg, 3 mmol). The reaction mixture was stirred at room temperature for three hours. The solvent was evaporated under reduced pressure and the residue was dissolved in a mixture of water (50 ml) and dichloromethane (50 ml). The aqueous layer was extracted three times with dichloromethane. The combined organic phases were evaporated under reduced pressure and the solid residue was dissolved in ethanol. Slow evaporation of ethanol gave colorless crystals (736 mg) in 80% yield.

### Refinement

The H atoms were geometrically placed and treated as riding atoms with C—H = 0.96 Å, and  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}$ (parent C-atom) except for methyl H atoms where  $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}$ (methyl C-atom).

### Figures




Fig. 1. Molecular structure of (I) with atom labels and 30% probability displacement ellipsoids.

## Diethyl 2,2'-(biphenyl-2,2'-diyldioxy)diacetate

### Crystal data

$\text{C}_{20}\text{H}_{22}\text{O}_6$	$Z = 2$
$M_r = 358.38$	$F(000) = 380$
Triclinic, $P\bar{1}$	$D_x = 1.342 \text{ Mg m}^{-3}$

# supplementary materials

---

$a = 7.4683 (13)$ Å	Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
$b = 10.8189 (15)$ Å	Cell parameters from 2971 reflections
$c = 12.050 (2)$ Å	$\theta = 3.1\text{--}26.3^\circ$
$\alpha = 104.733 (17)^\circ$	$\mu = 0.10 \text{ mm}^{-1}$
$\beta = 95.05 (2)^\circ$	$T = 153$ K
$\gamma = 106.897 (14)^\circ$	Chip, colorless
$V = 887.1 (3)$ Å <sup>3</sup>	$0.58 \times 0.53 \times 0.24$ mm

## Data collection

Rigaku AFC-8S diffractometer	3104 independent reflections
Radiation source: normal-focus sealed tube graphite	2787 reflections with $I > 2\sigma(I)$
Detector resolution: 14.6306 pixels mm <sup>-1</sup>	$R_{\text{int}} = 0.014$
$\omega$ scans	$\theta_{\text{max}} = 25.1^\circ, \theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan ( <i>REQAB</i> ; Jacobson, 1998)	$h = -8 \rightarrow 8$
$T_{\text{min}} = 0.945, T_{\text{max}} = 0.977$	$k = -12 \rightarrow 12$
6640 measured reflections	$l = -14 \rightarrow 12$

## Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.039$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.099$	H-atom parameters constrained
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0493P)^2 + 0.3462P]$ where $P = (F_o^2 + 2F_c^2)/3$
3104 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
235 parameters	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.20 \text{ e \AA}^{-3}$

## Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.26152 (13)	0.49153 (10)	-0.09823 (8)	0.0281 (2)
O2	0.10844 (14)	0.63977 (10)	-0.03410 (9)	0.0289 (3)
O3	0.37964 (13)	0.73993 (9)	0.17431 (8)	0.0229 (2)
O4	0.64269 (14)	1.13019 (10)	0.42800 (8)	0.0270 (2)
O5	0.71807 (19)	1.39716 (12)	0.43450 (10)	0.0455 (3)
O6	0.74896 (14)	1.45726 (9)	0.62997 (8)	0.0284 (2)
C1	0.2426 (2)	0.36891 (16)	-0.29568 (13)	0.0315 (3)
H1A	0.1754	0.3436	-0.3759	0.047*
H1B	0.2329	0.2872	-0.2723	0.047*
H1C	0.3768	0.4191	-0.2912	0.047*
C2	0.1555 (2)	0.45549 (19)	-0.21611 (13)	0.0407 (4)
H2A	0.1631	0.5380	-0.2395	0.049*
H2B	0.0201	0.4056	-0.2194	0.049*
C3	0.22147 (18)	0.58407 (13)	-0.01637 (12)	0.0213 (3)
C4	0.33426 (18)	0.60535 (13)	0.10190 (11)	0.0218 (3)
H4A	0.2593	0.5424	0.1406	0.026*
H4B	0.4532	0.5843	0.0911	0.026*
C5	0.50568 (17)	0.83979 (13)	0.14036 (11)	0.0197 (3)
C6	0.51573 (18)	0.97228 (13)	0.19636 (11)	0.0196 (3)
C7	0.64460 (19)	1.07739 (13)	0.16831 (12)	0.0239 (3)
H7A	0.6553	1.1682	0.2068	0.029*
C8	0.75716 (19)	1.05198 (14)	0.08549 (12)	0.0261 (3)
H8A	0.8428	1.1247	0.0667	0.031*
C9	0.74388 (19)	0.92018 (15)	0.03053 (12)	0.0267 (3)
H9A	0.8203	0.9023	-0.0266	0.032*
C10	0.61985 (19)	0.81358 (14)	0.05811 (12)	0.0244 (3)
H10A	0.6129	0.7232	0.0210	0.029*
C11	0.38246 (18)	1.00240 (13)	0.27616 (11)	0.0207 (3)
C12	0.18639 (19)	0.95132 (13)	0.23529 (12)	0.0242 (3)
H12A	0.1391	0.8899	0.1593	0.029*
C13	0.0593 (2)	0.98855 (14)	0.30363 (13)	0.0277 (3)
H13A	-0.0734	0.9537	0.2745	0.033*
C14	0.1289 (2)	1.07714 (14)	0.41479 (13)	0.0297 (3)
H14A	0.0430	1.1041	0.4615	0.036*
C15	0.3225 (2)	1.12712 (14)	0.45896 (13)	0.0281 (3)
H15A	0.3685	1.1868	0.5358	0.034*
C16	0.44882 (19)	1.08923 (13)	0.39009 (12)	0.0223 (3)
C17	0.7184 (2)	1.23201 (14)	0.53605 (12)	0.0284 (3)
H17A	0.8479	1.2336	0.5641	0.034*
H17B	0.6384	1.2107	0.5943	0.034*
C18	0.72672 (19)	1.36982 (14)	0.52501 (12)	0.0263 (3)
C19	0.7734 (2)	1.59703 (14)	0.63189 (14)	0.0349 (4)
H19A	0.6748	1.5994	0.5726	0.042*
H19B	0.8998	1.6391	0.6141	0.042*
C20	0.7560 (2)	1.67243 (15)	0.75077 (14)	0.0352 (4)

## supplementary materials

---

H20A	0.7719	1.7665	0.7541	0.053*
H20B	0.8544	1.6699	0.8088	0.053*
H20C	0.6303	1.6303	0.7674	0.053*

### *Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0283 (5)	0.0326 (6)	0.0214 (5)	0.0160 (4)	0.0003 (4)	-0.0010 (4)
O2	0.0284 (5)	0.0264 (5)	0.0338 (6)	0.0133 (4)	0.0033 (4)	0.0075 (4)
O3	0.0279 (5)	0.0169 (5)	0.0224 (5)	0.0059 (4)	0.0092 (4)	0.0032 (4)
O4	0.0303 (5)	0.0263 (5)	0.0210 (5)	0.0131 (4)	0.0000 (4)	-0.0019 (4)
O5	0.0766 (9)	0.0358 (6)	0.0258 (6)	0.0188 (6)	0.0108 (6)	0.0107 (5)
O6	0.0422 (6)	0.0211 (5)	0.0213 (5)	0.0123 (4)	0.0035 (4)	0.0036 (4)
C1	0.0309 (7)	0.0337 (8)	0.0238 (8)	0.0074 (6)	0.0037 (6)	0.0020 (6)
C2	0.0376 (8)	0.0587 (11)	0.0225 (8)	0.0246 (8)	-0.0030 (6)	-0.0013 (7)
C3	0.0195 (6)	0.0172 (6)	0.0270 (7)	0.0054 (5)	0.0068 (5)	0.0058 (5)
C4	0.0249 (7)	0.0167 (6)	0.0235 (7)	0.0077 (5)	0.0054 (5)	0.0039 (5)
C5	0.0204 (6)	0.0194 (6)	0.0174 (6)	0.0047 (5)	0.0021 (5)	0.0046 (5)
C6	0.0210 (6)	0.0202 (6)	0.0170 (6)	0.0075 (5)	0.0013 (5)	0.0041 (5)
C7	0.0269 (7)	0.0203 (7)	0.0225 (7)	0.0067 (5)	0.0021 (5)	0.0047 (5)
C8	0.0241 (7)	0.0269 (7)	0.0254 (7)	0.0039 (5)	0.0052 (5)	0.0093 (6)
C9	0.0247 (7)	0.0315 (7)	0.0212 (7)	0.0068 (6)	0.0082 (5)	0.0044 (6)
C10	0.0257 (7)	0.0221 (7)	0.0219 (7)	0.0071 (5)	0.0053 (5)	0.0008 (5)
C11	0.0265 (7)	0.0172 (6)	0.0212 (7)	0.0094 (5)	0.0063 (5)	0.0072 (5)
C12	0.0288 (7)	0.0193 (6)	0.0265 (7)	0.0088 (5)	0.0067 (6)	0.0085 (5)
C13	0.0268 (7)	0.0240 (7)	0.0373 (8)	0.0098 (6)	0.0119 (6)	0.0138 (6)
C14	0.0370 (8)	0.0264 (7)	0.0356 (8)	0.0171 (6)	0.0198 (6)	0.0138 (6)
C15	0.0399 (8)	0.0242 (7)	0.0241 (7)	0.0153 (6)	0.0117 (6)	0.0063 (6)
C16	0.0290 (7)	0.0189 (6)	0.0217 (7)	0.0110 (5)	0.0054 (5)	0.0067 (5)
C17	0.0362 (8)	0.0254 (7)	0.0206 (7)	0.0137 (6)	-0.0020 (6)	0.0000 (6)
C18	0.0289 (7)	0.0281 (7)	0.0200 (7)	0.0096 (6)	0.0026 (5)	0.0040 (6)
C19	0.0524 (9)	0.0208 (7)	0.0330 (9)	0.0128 (7)	0.0078 (7)	0.0094 (6)
C20	0.0516 (9)	0.0201 (7)	0.0318 (8)	0.0105 (7)	0.0062 (7)	0.0056 (6)

### *Geometric parameters ( $\text{\AA}$ , $^\circ$ )*

O1—C3	1.3374 (16)	C8—C9	1.381 (2)
O1—C2	1.4618 (18)	C8—H8A	0.9500
O2—C3	1.2044 (16)	C9—C10	1.388 (2)
O3—C5	1.3864 (16)	C9—H9A	0.9500
O3—C4	1.4181 (16)	C10—H10A	0.9500
O4—C16	1.3816 (17)	C11—C12	1.399 (2)
O4—C17	1.4167 (17)	C11—C16	1.402 (2)
O5—C18	1.2017 (18)	C12—C13	1.3900 (19)
O6—C18	1.3362 (17)	C12—H12A	0.9500
O6—C19	1.4634 (17)	C13—C14	1.385 (2)
C1—C2	1.492 (2)	C13—H13A	0.9500
C1—H1A	0.9800	C14—C15	1.389 (2)
C1—H1B	0.9800	C14—H14A	0.9500

C1—H1C	0.9800	C15—C16	1.3913 (19)
C2—H2A	0.9900	C15—H15A	0.9500
C2—H2B	0.9900	C17—C18	1.514 (2)
C3—C4	1.5175 (19)	C17—H17A	0.9900
C4—H4A	0.9900	C17—H17B	0.9900
C4—H4B	0.9900	C19—C20	1.495 (2)
C5—C10	1.3916 (18)	C19—H19A	0.9900
C5—C6	1.3971 (18)	C19—H19B	0.9900
C6—C7	1.3966 (19)	C20—H20A	0.9800
C6—C11	1.4897 (18)	C20—H20B	0.9800
C7—C8	1.386 (2)	C20—H20C	0.9800
C7—H7A	0.9500		
C3—O1—C2	116.34 (11)	C9—C10—H10A	120.2
C5—O3—C4	116.66 (10)	C5—C10—H10A	120.2
C16—O4—C17	117.43 (11)	C12—C11—C16	118.38 (12)
C18—O6—C19	115.91 (11)	C12—C11—C6	119.93 (12)
C2—C1—H1A	109.5	C16—C11—C6	121.54 (12)
C2—C1—H1B	109.5	C13—C12—C11	121.36 (13)
H1A—C1—H1B	109.5	C13—C12—H12A	119.3
C2—C1—H1C	109.5	C11—C12—H12A	119.3
H1A—C1—H1C	109.5	C14—C13—C12	119.08 (13)
H1B—C1—H1C	109.5	C14—C13—H13A	120.5
O1—C2—C1	107.38 (12)	C12—C13—H13A	120.5
O1—C2—H2A	110.2	C13—C14—C15	120.95 (13)
C1—C2—H2A	110.2	C13—C14—H14A	119.5
O1—C2—H2B	110.2	C15—C14—H14A	119.5
C1—C2—H2B	110.2	C14—C15—C16	119.63 (14)
H2A—C2—H2B	108.5	C14—C15—H15A	120.2
O2—C3—O1	124.57 (13)	C16—C15—H15A	120.2
O2—C3—C4	125.14 (12)	O4—C16—C15	123.64 (12)
O1—C3—C4	110.27 (11)	O4—C16—C11	115.79 (12)
O3—C4—C3	111.80 (11)	C15—C16—C11	120.56 (13)
O3—C4—H4A	109.3	O4—C17—C18	111.40 (12)
C3—C4—H4A	109.3	O4—C17—H17A	109.3
O3—C4—H4B	109.3	C18—C17—H17A	109.3
C3—C4—H4B	109.3	O4—C17—H17B	109.3
H4A—C4—H4B	107.9	C18—C17—H17B	109.3
O3—C5—C10	123.59 (12)	H17A—C17—H17B	108.0
O3—C5—C6	115.63 (11)	O5—C18—O6	124.81 (13)
C10—C5—C6	120.77 (12)	O5—C18—C17	124.61 (13)
C7—C6—C5	118.18 (12)	O6—C18—C17	110.57 (12)
C7—C6—C11	119.86 (12)	O6—C19—C20	108.07 (12)
C5—C6—C11	121.75 (12)	O6—C19—H19A	110.1
C8—C7—C6	121.36 (13)	C20—C19—H19A	110.1
C8—C7—H7A	119.3	O6—C19—H19B	110.1
C6—C7—H7A	119.3	C20—C19—H19B	110.1
C9—C8—C7	119.49 (13)	H19A—C19—H19B	108.4
C9—C8—H8A	120.3	C19—C20—H20A	109.5
C7—C8—H8A	120.3	C19—C20—H20B	109.5

## supplementary materials

---

C8—C9—C10	120.55 (13)	H20A—C20—H20B	109.5
C8—C9—H9A	119.7	C19—C20—H20C	109.5
C10—C9—H9A	119.7	H20A—C20—H20C	109.5
C9—C10—C5	119.64 (13)	H20B—C20—H20C	109.5
C3—O1—C2—C1	171.24 (12)	C7—C6—C11—C16	-58.95 (18)
C2—O1—C3—O2	-1.3 (2)	C5—C6—C11—C16	126.46 (14)
C2—O1—C3—C4	176.98 (12)	C16—C11—C12—C13	2.17 (19)
C5—O3—C4—C3	-67.26 (14)	C6—C11—C12—C13	-173.39 (12)
O2—C3—C4—O3	-33.27 (18)	C11—C12—C13—C14	-0.6 (2)
O1—C3—C4—O3	148.41 (11)	C12—C13—C14—C15	-1.0 (2)
C4—O3—C5—C10	-15.26 (18)	C13—C14—C15—C16	0.9 (2)
C4—O3—C5—C6	166.12 (11)	C17—O4—C16—C15	-9.89 (19)
O3—C5—C6—C7	178.06 (11)	C17—O4—C16—C11	171.55 (11)
C10—C5—C6—C7	-0.60 (19)	C14—C15—C16—O4	-177.75 (12)
O3—C5—C6—C11	-7.26 (18)	C14—C15—C16—C11	0.8 (2)
C10—C5—C6—C11	174.08 (12)	C12—C11—C16—O4	176.37 (11)
C5—C6—C7—C8	1.4 (2)	C6—C11—C16—O4	-8.15 (18)
C11—C6—C7—C8	-173.42 (12)	C12—C11—C16—C15	-2.24 (19)
C6—C7—C8—C9	-0.9 (2)	C6—C11—C16—C15	173.24 (12)
C7—C8—C9—C10	-0.4 (2)	C16—O4—C17—C18	-75.23 (15)
C8—C9—C10—C5	1.1 (2)	C19—O6—C18—O5	-3.1 (2)
O3—C5—C10—C9	-179.17 (12)	C19—O6—C18—C17	175.61 (12)
C6—C5—C10—C9	-0.6 (2)	O4—C17—C18—O5	-17.8 (2)
C7—C6—C11—C12	116.47 (14)	O4—C17—C18—O6	163.46 (11)
C5—C6—C11—C12	-58.13 (18)	C18—O6—C19—C20	168.59 (13)

*Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )*

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C4—H4b…O1 <sup>i</sup>	0.99	2.50	3.472 (2)	167.7 (1)

Symmetry codes: (i)  $-x+1, -y+1, -z$ .

Fig. 1

