

N-[4-(Benzylsulfamoyl)phenyl]acetamide

Peter John,^a Waqar Ahmad,^a Islam Ullah Khan,^{a†} Shahzad Sharif^a and Edward R. T. Tiekink^{b*}

^aMaterials Chemistry Laboratory, Department of Chemistry, Government College University, Lahore 54000, Pakistan, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: edward.tiekink@gmail.com


Received 6 July 2010; accepted 13 July 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.041; wR factor = 0.124; data-to-parameter ratio = 18.2.

A folded conformation is found for the title compound, $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_3\text{S}$, whereby the benzene rings come into close proximity [centroid-centroid distance = 4.0357 (12) Å and the dihedral angle between them = 24.37 (10)°]. The amide group is coplanar with the benzene ring to which it is bound [C—C—N—C torsion angle = 11.1 (3)°]. In the crystal packing, two-dimensional arrays in the (101) plane are formed via N—H···O hydrogen bonding.

Related literature

For background to the pharmacological uses of sulfonamides, see: Beate *et al.* (1998); Kazmierski *et al.* (2004). For related structures, see: Khan *et al.* (2010); Sharif *et al.* (2010).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_3\text{S}$
 $M_r = 304.37$
 Monoclinic, $P2_1/n$
 $a = 9.0646$ (9) Å

$b = 13.6888$ (14) Å
 $c = 12.1651$ (12) Å
 $\beta = 98.635$ (5)°
 $V = 1492.4$ (3) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.23$ mm⁻¹

$T = 293$ K
 $0.19 \times 0.09 \times 0.07$ mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.868$, $T_{\max} = 0.948$

13777 measured reflections
 3577 independent reflections
 2689 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.124$
 $S = 1.02$
 3577 reflections
 197 parameters
 2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.25$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1n}\cdots\text{O3}^i$	0.89 (2)	2.00 (2)	2.877 (2)	168 (2)
$\text{N2}-\text{H2n}\cdots\text{O2}^{ii}$	0.90 (2)	2.03 (2)	2.921 (2)	172 (2)

Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and DIAMOND (Brandenburg, 2006); software used to prepare material for publication: publCIF (Westrip, 2010).

We are grateful to Mr Munawar Hussain, Engineering Cell GC University, Lahore, for providing support services to the Materials Chemistry Laboratory.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2043).

References

- Beate, G., Nadenik, P. & Wagner, H. (1998). WO Patent No. 9855481.
 Brandenburg, K. (2006). DIAMOND. Crystal Impact GbR, Bonn, Germany.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Kazmierski, W. M., Aquino, C. J., Bifulco, N., Boros, E. E., Chauder, B. A., Chong, P. Y., Duan, M., Deanda, F. Jr, Koble, C. S., Mclean, E. W., Peckham, J. P., Perkins, A. C., Thompson, J. B. & Vanderwall, D. (2004). WO Patent No. 2004054974.
 Khan, I. U., Mariam, I., Zia-ur-Rehman, M., Arif Sajjad, M. & Sharif, S. (2010). *Acta Cryst.* **E66**, o1088.
 Sharif, S., Iqbal, H., Khan, I. U., John, P. & Tiekink, E. R. T. (2010). *Acta Cryst.* **E66**, o1288.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

† Additional correspondence author, e-mail: iuklodhi@yahoo.com.

supplementary materials

Acta Cryst. (2010). E66, o2048 [doi:10.1107/S1600536810027698]

N-[4-(Benzylsulfamoyl)phenyl]acetamide

P. John, W. Ahmad, I. U. Khan, S. Sharif and E. R. T. Tiekink

Comment

Sulfonamide drugs are used, for example, as inhibitors of HIV infection (Kazmierski *et al.*, 2004) and as anti-hypertensive drugs (Beate *et al.*, 1998). In connection with on-going structural studies of sulfonamides (Khan *et al.*, 2010; Sharif *et al.*, 2010), the crystal and molecular structure of the title compound, C₁₅H₁₆N₂O₃S, was investigated.

The molecule of C₁₅H₁₆N₂O₃S has a folded conformation with the benzene ring of the benzyl group somewhat orientated over the S-bound benzene ring. The rings are approximately parallel, forming a dihedral angle of 24.37 (10) °; the distance between the ring centroids is 4.0357 (12) Å. The amide group is essentially co-planar with the ring to which it is bound as seen in the C10–C11–N2–C14 torsion angle of 11.1 (3) °.

The crystal packing is dominated by N–H···O hydrogen bonds whereby the N1–H atom forms a hydrogen bond to the amide-carbonyl, and the amide N2–H forms a contact with the S-bound O2 atom, Table 1. The former leads to centrosymmetric aggregates and these are connected by the latter into a 2-D array in the (1 0 1) plane, Fig. 2.

Experimental

To 4-acetamidobenzenesulfonyl chloride (498 mg, 2.14 mmol) in distilled water (10 ml) was added benzylamine (234 ml, 2.14 mmol), the reaction mixture was stirred at room temperature while maintaining the pH of the reaction mixture at 8 using 3% sodium carbonate. The progress of the reaction was monitored by TLC. After consumption of all the reactants, the precipitates were filtered, dried and crystallized from methanol to yield colourless crystals.

Refinement

The C-bound H atoms were geometrically placed (C–H = 0.93–0.97 Å) and refined as riding with $U_{iso}(H) = 1.2-1.5U_{eq}(C)$. The N-bound H atom was refined with the distance restraint N–H = 0.88±0.01 Å, and with $U_{iso}(H) = 1.2U_{eq}(N)$.

Figures




Fig. 1. The molecular structure of C₁₅H₁₆N₂O₃S showing the atom-labelling scheme and displacement ellipsoids at the 35% probability level.




Fig. 2. A view of the supramolecular 2-D array in the (1 0 1) plane mediated by N–H···O hydrogen bonding (orange dashed lines) in $C_{15}H_{16}N_2O_3S$. Colour code: S, yellow; O, red; N, blue; C, grey; and H, green.

N-[4-(Benzyulfamoyl)phenyl]acetamide

Crystal data

$C_{15}H_{16}N_2O_3S$

$M_r = 304.37$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 9.0646$ (9) Å

$b = 13.6888$ (14) Å

$c = 12.1651$ (12) Å

$\beta = 98.635$ (5)°

$V = 1492.4$ (3) Å³

$Z = 4$

$F(000) = 640$

$D_x = 1.355$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4154 reflections

$\theta = 2.6$ – 27.9 °

$\mu = 0.23$ mm⁻¹

$T = 293$ K

Prism, colourless

$0.19 \times 0.09 \times 0.07$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.868$, $T_{\max} = 0.948$

13777 measured reflections

3577 independent reflections

2689 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 28.0$ °, $\theta_{\min} = 2.3$ °

$h = -11 \rightarrow 11$

$k = -18 \rightarrow 17$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.124$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0643P)^2 + 0.3076P]$
3577 reflections	where $P = (F_o^2 + 2F_c^2)/3$
197 parameters	$(\Delta/\sigma)_{\max} = 0.001$
2 restraints	$\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.11090 (5)	0.33181 (3)	0.82483 (3)	0.03999 (15)
O1	0.21873 (16)	0.27094 (10)	0.88809 (10)	0.0536 (4)
O2	-0.04454 (15)	0.31532 (11)	0.82667 (12)	0.0594 (4)
O3	0.02478 (16)	0.40834 (11)	0.26180 (11)	0.0596 (4)
N1	0.14186 (17)	0.44201 (11)	0.86867 (12)	0.0448 (4)
H1N	0.0786 (18)	0.4846 (12)	0.8311 (15)	0.054*
N2	0.20419 (16)	0.31175 (10)	0.35277 (11)	0.0399 (3)
H2N	0.2876 (15)	0.2778 (12)	0.3473 (16)	0.048*
C1	0.2944 (2)	0.47687 (16)	0.90443 (16)	0.0559 (5)
H1A	0.3409	0.4347	0.9638	0.067*
H1B	0.2888	0.5418	0.9355	0.067*
C2	0.3944 (2)	0.48133 (13)	0.81674 (15)	0.0464 (4)
C3	0.3606 (2)	0.54229 (14)	0.72574 (17)	0.0532 (5)
H3	0.2748	0.5805	0.7184	0.064*
C4	0.4533 (3)	0.54678 (16)	0.6461 (2)	0.0661 (6)
H4	0.4295	0.5880	0.5852	0.079*
C5	0.5799 (3)	0.49141 (18)	0.6555 (2)	0.0732 (7)
H5	0.6416	0.4945	0.6011	0.088*
C6	0.6154 (3)	0.43158 (18)	0.7452 (2)	0.0745 (7)
H6	0.7021	0.3943	0.7523	0.089*
C7	0.5230 (2)	0.42612 (15)	0.8254 (2)	0.0611 (6)
H7	0.5477	0.3848	0.8860	0.073*
C8	0.13636 (18)	0.32476 (11)	0.68406 (13)	0.0354 (3)
C9	0.0308 (2)	0.36444 (16)	0.60359 (15)	0.0499 (5)
H9	-0.0539	0.3934	0.6238	0.060*
C10	0.0494 (2)	0.36167 (15)	0.49303 (15)	0.0496 (5)
H10	-0.0223	0.3888	0.4390	0.059*

supplementary materials

C11	0.17580 (18)	0.31823 (11)	0.46307 (13)	0.0343 (3)
C12	0.2803 (2)	0.27757 (14)	0.54465 (14)	0.0436 (4)
H12	0.3647	0.2477	0.5250	0.052*
C13	0.26102 (19)	0.28078 (13)	0.65483 (14)	0.0426 (4)
H13	0.3321	0.2533	0.7091	0.051*
C14	0.1335 (2)	0.35627 (13)	0.26110 (14)	0.0418 (4)
C15	0.1978 (3)	0.33645 (17)	0.15660 (16)	0.0605 (6)
H15A	0.1635	0.2741	0.1270	0.091*
H15B	0.3048	0.3360	0.1730	0.091*
H15C	0.1664	0.3865	0.1029	0.091*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0444 (3)	0.0476 (3)	0.0295 (2)	-0.00397 (18)	0.01067 (17)	0.00348 (17)
O1	0.0718 (10)	0.0527 (7)	0.0358 (6)	0.0085 (6)	0.0066 (6)	0.0080 (6)
O2	0.0492 (8)	0.0830 (10)	0.0502 (8)	-0.0168 (7)	0.0209 (6)	0.0030 (7)
O3	0.0599 (9)	0.0774 (10)	0.0420 (7)	0.0218 (7)	0.0096 (6)	0.0121 (7)
N1	0.0510 (9)	0.0497 (9)	0.0341 (7)	0.0033 (7)	0.0079 (6)	-0.0005 (6)
N2	0.0407 (8)	0.0491 (8)	0.0312 (7)	0.0066 (6)	0.0094 (6)	-0.0011 (6)
C1	0.0667 (13)	0.0620 (12)	0.0363 (9)	-0.0135 (10)	-0.0015 (9)	-0.0055 (9)
C2	0.0470 (10)	0.0429 (9)	0.0467 (10)	-0.0085 (8)	-0.0014 (8)	-0.0028 (8)
C3	0.0547 (12)	0.0492 (10)	0.0555 (11)	0.0010 (8)	0.0081 (9)	0.0034 (9)
C4	0.0772 (16)	0.0596 (13)	0.0634 (14)	-0.0104 (11)	0.0167 (12)	0.0071 (11)
C5	0.0672 (15)	0.0670 (14)	0.0917 (18)	-0.0175 (12)	0.0319 (14)	-0.0142 (13)
C6	0.0467 (13)	0.0666 (14)	0.109 (2)	-0.0009 (10)	0.0096 (13)	-0.0133 (14)
C7	0.0554 (13)	0.0516 (11)	0.0706 (14)	-0.0016 (9)	-0.0091 (11)	0.0039 (10)
C8	0.0362 (9)	0.0418 (8)	0.0290 (7)	-0.0043 (6)	0.0079 (6)	0.0004 (6)
C9	0.0395 (10)	0.0759 (13)	0.0357 (9)	0.0162 (9)	0.0105 (7)	0.0016 (9)
C10	0.0420 (10)	0.0736 (12)	0.0329 (9)	0.0161 (9)	0.0050 (7)	0.0031 (9)
C11	0.0351 (8)	0.0379 (8)	0.0305 (7)	-0.0024 (6)	0.0074 (6)	-0.0015 (6)
C12	0.0397 (9)	0.0541 (10)	0.0383 (9)	0.0122 (8)	0.0098 (7)	0.0001 (8)
C13	0.0408 (10)	0.0523 (10)	0.0341 (8)	0.0080 (8)	0.0038 (7)	0.0044 (7)
C14	0.0461 (10)	0.0468 (9)	0.0327 (8)	-0.0030 (8)	0.0063 (7)	-0.0006 (7)
C15	0.0711 (14)	0.0797 (15)	0.0328 (9)	0.0057 (11)	0.0145 (9)	0.0024 (9)

Geometric parameters (\AA , $^\circ$)

S1—O1	1.4199 (13)	C5—C6	1.363 (4)
S1—O2	1.4304 (14)	C5—H5	0.9300
S1—N1	1.6105 (16)	C6—C7	1.380 (3)
S1—C8	1.7647 (16)	C6—H6	0.9300
O3—C14	1.217 (2)	C7—H7	0.9300
N1—C1	1.466 (2)	C8—C9	1.374 (2)
N1—H1N	0.893 (9)	C8—C13	1.373 (2)
N2—C14	1.346 (2)	C9—C10	1.381 (2)
N2—C11	1.4064 (19)	C9—H9	0.9300
N2—H2N	0.899 (9)	C10—C11	1.387 (2)
C1—C2	1.501 (3)	C10—H10	0.9300

C1—H1A	0.9700	C11—C12	1.382 (2)
C1—H1B	0.9700	C12—C13	1.378 (2)
C2—C7	1.380 (3)	C12—H12	0.9300
C2—C3	1.383 (3)	C13—H13	0.9300
C3—C4	1.376 (3)	C14—C15	1.501 (2)
C3—H3	0.9300	C15—H15A	0.9600
C4—C5	1.366 (3)	C15—H15B	0.9600
C4—H4	0.9300	C15—H15C	0.9600
O1—S1—O2	119.90 (9)	C7—C6—H6	119.9
O1—S1—N1	107.38 (8)	C2—C7—C6	120.8 (2)
O2—S1—N1	105.41 (9)	C2—C7—H7	119.6
O1—S1—C8	108.31 (8)	C6—C7—H7	119.6
O2—S1—C8	106.21 (8)	C9—C8—C13	120.05 (15)
N1—S1—C8	109.33 (8)	C9—C8—S1	119.43 (13)
C1—N1—S1	120.86 (14)	C13—C8—S1	120.52 (13)
C1—N1—H1N	116.3 (13)	C8—C9—C10	120.62 (16)
S1—N1—H1N	112.0 (13)	C8—C9—H9	119.7
C14—N2—C11	129.01 (14)	C10—C9—H9	119.7
C14—N2—H2N	118.3 (13)	C9—C10—C11	119.62 (16)
C11—N2—H2N	112.4 (13)	C9—C10—H10	120.2
N1—C1—C2	116.46 (15)	C11—C10—H10	120.2
N1—C1—H1A	108.2	C12—C11—C10	119.21 (15)
C2—C1—H1A	108.2	C12—C11—N2	117.12 (14)
N1—C1—H1B	108.2	C10—C11—N2	123.66 (15)
C2—C1—H1B	108.2	C13—C12—C11	120.82 (15)
H1A—C1—H1B	107.3	C13—C12—H12	119.6
C7—C2—C3	118.3 (2)	C11—C12—H12	119.6
C7—C2—C1	121.10 (19)	C8—C13—C12	119.68 (16)
C3—C2—C1	120.57 (18)	C8—C13—H13	120.2
C4—C3—C2	120.4 (2)	C12—C13—H13	120.2
C4—C3—H3	119.8	O3—C14—N2	123.00 (16)
C2—C3—H3	119.8	O3—C14—C15	121.99 (17)
C5—C4—C3	120.7 (2)	N2—C14—C15	115.01 (16)
C5—C4—H4	119.7	C14—C15—H15A	109.5
C3—C4—H4	119.7	C14—C15—H15B	109.5
C6—C5—C4	119.7 (2)	H15A—C15—H15B	109.5
C6—C5—H5	120.2	C14—C15—H15C	109.5
C4—C5—H5	120.2	H15A—C15—H15C	109.5
C5—C6—C7	120.2 (2)	H15B—C15—H15C	109.5
C5—C6—H6	119.9		
O1—S1—N1—C1	35.74 (15)	O1—S1—C8—C13	-11.08 (17)
O2—S1—N1—C1	164.65 (14)	O2—S1—C8—C13	-141.10 (15)
C8—S1—N1—C1	-81.56 (15)	N1—S1—C8—C13	105.63 (15)
S1—N1—C1—C2	64.4 (2)	C13—C8—C9—C10	-0.8 (3)
N1—C1—C2—C7	-118.2 (2)	S1—C8—C9—C10	178.69 (16)
N1—C1—C2—C3	62.7 (2)	C8—C9—C10—C11	0.1 (3)
C7—C2—C3—C4	0.4 (3)	C9—C10—C11—C12	0.6 (3)
C1—C2—C3—C4	179.41 (19)	C9—C10—C11—N2	179.85 (18)

supplementary materials

C2—C3—C4—C5	-0.1 (3)	C14—N2—C11—C12	-169.65 (17)
C3—C4—C5—C6	-0.5 (4)	C14—N2—C11—C10	11.1 (3)
C4—C5—C6—C7	0.7 (4)	C10—C11—C12—C13	-0.8 (3)
C3—C2—C7—C6	-0.1 (3)	N2—C11—C12—C13	179.98 (16)
C1—C2—C7—C6	-179.14 (19)	C9—C8—C13—C12	0.6 (3)
C5—C6—C7—C2	-0.5 (3)	S1—C8—C13—C12	-178.81 (14)
O1—S1—C8—C9	169.47 (15)	C11—C12—C13—C8	0.1 (3)
O2—S1—C8—C9	39.45 (17)	C11—N2—C14—O3	-3.2 (3)
N1—S1—C8—C9	-73.82 (16)	C11—N2—C14—C15	177.27 (17)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1n \cdots O3 ⁱ	0.894 (17)	1.996 (17)	2.877 (2)	168.2 (16)
N2—H2n \cdots O2 ⁱⁱ	0.898 (15)	2.029 (15)	2.921 (2)	171.5 (16)

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $x+1/2, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

