

5,8-Dibromo-2,11-dithia[3,3](2,6)-pyridinoparacyclophane

Si-Si Li,* Bei Zhang and Hong-Lin Zhang

Key Laboratory of Pesticides and Chemical Biology of the Ministry of Education, College of Chemistry, Central China Normal University, Wuhan 430079, People's Republic of China

Correspondence e-mail: 78766912@163.com


Received 1 July 2010; accepted 20 July 2010

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.031; wR factor = 0.078; data-to-parameter ratio = 20.9.

The title compound, $\text{C}_{15}\text{H}_{13}\text{Br}_2\text{NS}_2$ [systematic name: 1²,1⁵-dibromo-2,7-dithia-1(1,4)-benzena-5(2,6)-pyridinaoctaphane], contains a dibromo-substituted benzene ring and a pyridine ring that are linked by a pair of bridging $-\text{CH}_2\text{SCH}_2-$ groups. There is a weak $\pi-\pi$ interaction between the rings, the distance between the ring centroids being 3.572 (4) Å. The rings are not parallel, but form a dihedral angle of 18.29 (4)°.

Related literature

For the preparation of the title compound, see: Kay & Baek (1997); Scheytza *et al.* (1999); Xu *et al.* (2008). For further information on paracyclophane and its derivatives, see: Wang *et al.* (2006); Yamamoto *et al.* (1997).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{13}\text{Br}_2\text{NS}_2$	$V = 1525.2$ (4) Å ³
$M_r = 431.20$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 8.9275$ (15) Å	$\mu = 5.58$ mm ⁻¹
$b = 18.879$ (3) Å	$T = 298$ K
$c = 9.3213$ (15) Å	$0.16 \times 0.12 \times 0.10$ mm
$\beta = 103.878$ (3)°	

Data collection

Bruker SMART CCD area-detector diffractometer	11428 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	3775 independent reflections
$T_{\min} = 0.452$, $T_{\max} = 0.573$	3018 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$	181 parameters
$wR(F^2) = 0.078$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.56$ e Å ⁻³
3775 reflections	$\Delta\rho_{\text{min}} = -0.46$ e Å ⁻³

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Professor Sheng-Hua Liu for technical assistance and Dr Jian-Long Xia for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2253).

References

- Bruker (1997). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (1999). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Kay, K. Y. & Baek, Y. G. (1997). *Chem. Ber. Recl.* **130**, 581–584.
 Scheytza, H., Rademacher, O. & Reibig, H.-U. (1999). *Eur. J. Org. Chem.* **9**, 2373–2381.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wang, W., Xu, J., Zhang, X. & Lai, Y. H. (2006). *Macromolecules*, **39**, 7277–7285.
 Xu, J. W., Wang, W. L., Lin, T. T., Sun, Z. & Lai, Y. H. (2008). *Supramol. Chem.* **20**, 723–730.
 Yamamoto, M., Wu, L. P., Kuroda-Sowa, T., Maekawa, M., Suenaga, Y. & Munakata, M. (1997). *Inorg. Chim. Acta*, **258**, 89–91.

supplementary materials

Acta Cryst. (2010). E66, o2125 [doi:10.1107/S1600536810028850]

5,8-Dibromo-2,11-dithia[3,3](2,6)pyridinoparacyclophane

S.-S. Li, B. Zhang and H.-L. Zhang

Comment

In our research, we have synthesized the title compound 5,8-dibromo-2,11-dithia[3,3](2,6)pyridinoparacyclophane. In the crystal structure, there are no classical hydrogen bonds, but is a weak π - π interaction. The distance between the centroid of the pyridine ring and the centroid of the benzene ring is 3.574 (4) Å, in addition, the angle between pyridine ring and the benzene ring is 18.29 (4)°.

Experimental

The title compound was synthesized according to a modified literature procedure (Scheytza *et al.*, 1999).

A solution with equimolar amounts of the dithiol(3.26g, 10mmol) and 2,6-bis(bromomethyl) pyridine in degassed THF (500 mL) was added dropwise under N₂ over 12 h to a refluxing solution of K₂CO₃(6.9g,50mmol) in EtOH (1.5 L). After an additional 2h at the reflux temperature, the mixture was cooled and the solvent. The resulting residue was treated with CH₂Cl₂ (500 mL) and water (500mL). The organic phase was separated, and the aqueous layer extracted with CH₂Cl₂ three times. The combined organic layers were dried over MgSO₄, then the solvent was removed, and the resulting solid was chromatographed on silica gel using CH₂Cl₂ as eluent. Colourless single crystals of the title compound suitable for X-ray diffraction were obtained by slow evaporation of a dichloromethane-n-hexane (1:30) solution over a period of 6 hours.

Refinement

All H atoms were initially located in a difference map, but were constrained to an idealized geometry. Constrained bond lengths and isotropic displacement parameters: (C—H =0.93 Å) and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic H atoms, and (C—H =0.97 Å) and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for methylene.

Figures




Fig. 1. A view of the title compound, showing the atom-labelling scheme, with displacement ellipsoids drawn at the 50% probability level.

1²,1⁵-dibromo-2,7-dithia-1(1,4)-benzena-5(2,6)-pyridinaoctaphane

Crystal data

$C_{15}H_{13}Br_2NS_2$	$F(000) = 848$
$M_r = 431.20$	$D_x = 1.878 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 4039 reflections
$a = 8.9275 (15) \text{ \AA}$	$\theta = 2.5\text{--}27.8^\circ$
$b = 18.879 (3) \text{ \AA}$	$\mu = 5.58 \text{ mm}^{-1}$
$c = 9.3213 (15) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 103.878 (3)^\circ$	Block, colorless
$V = 1525.2 (4) \text{ \AA}^3$	$0.16 \times 0.12 \times 0.10 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART CCD area-detector diffractometer	3775 independent reflections
Radiation source: fine-focus sealed tube graphite	3018 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.026$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 2.2^\circ$
$T_{\text{min}} = 0.452$, $T_{\text{max}} = 0.573$	$h = -11 \rightarrow 11$
11428 measured reflections	$k = -25 \rightarrow 25$
	$l = -10 \rightarrow 12$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.031$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.078$	H-atom parameters constrained
$S = 1.03$	$w = 1/[\sigma^2(F_o^2) + (0.0414P)^2 + 0.2049P]$
3775 reflections	where $P = (F_o^2 + 2F_c^2)/3$
181 parameters	$(\Delta/\sigma)_{\text{max}} = 0.002$
0 restraints	$\Delta\rho_{\text{max}} = 0.56 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.46 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.63099 (3)	0.048688 (14)	0.87832 (3)	0.04897 (10)
Br2	0.22924 (4)	0.139645 (16)	0.22281 (3)	0.05124 (10)
C1	0.3799 (3)	0.03373 (12)	0.6206 (3)	0.0338 (5)
C2	0.5079 (3)	0.07344 (12)	0.6886 (3)	0.0334 (5)
C3	0.5520 (3)	0.13348 (12)	0.6246 (3)	0.0331 (5)
H3	0.6384	0.1587	0.6742	0.040*
C4	0.4686 (3)	0.15680 (13)	0.4867 (3)	0.0324 (5)
C5	0.3444 (3)	0.11522 (13)	0.4148 (3)	0.0338 (5)
C6	0.3002 (3)	0.05574 (12)	0.4802 (3)	0.0372 (5)
H6	0.2154	0.0298	0.4295	0.045*
C7	0.5137 (3)	0.22521 (13)	0.4279 (3)	0.0383 (6)
H7A	0.5180	0.2181	0.3259	0.046*
H7B	0.6168	0.2376	0.4835	0.046*
C8	0.3410 (3)	0.28873 (14)	0.6143 (3)	0.0371 (5)
H8A	0.4332	0.2727	0.6849	0.045*
H8B	0.3134	0.3346	0.6471	0.045*
C9	0.2123 (3)	0.23745 (12)	0.6156 (2)	0.0301 (5)
C10	0.0700 (3)	0.24389 (14)	0.5179 (3)	0.0382 (5)
H10	0.0501	0.2810	0.4505	0.046*
C11	-0.0417 (3)	0.19455 (15)	0.5220 (3)	0.0414 (6)
H11	-0.1389	0.1982	0.4580	0.050*
C12	-0.0085 (3)	0.13961 (13)	0.6216 (3)	0.0387 (6)
H12	-0.0819	0.1050	0.6243	0.046*
C13	0.1362 (3)	0.13660 (11)	0.7180 (3)	0.0308 (5)
C14	0.1781 (3)	0.07834 (13)	0.8309 (3)	0.0384 (6)
H14A	0.1171	0.0840	0.9034	0.046*
H14B	0.2858	0.0835	0.8820	0.046*
C15	0.3230 (3)	-0.02833 (13)	0.6927 (3)	0.0426 (6)
H15A	0.4036	-0.0430	0.7770	0.051*
H15B	0.3036	-0.0674	0.6232	0.051*
N1	0.2443 (2)	0.18540 (10)	0.7172 (2)	0.0302 (4)
S1	0.14839 (8)	-0.01069 (3)	0.75438 (8)	0.04531 (17)
S2	0.38563 (7)	0.29931 (3)	0.43559 (7)	0.03722 (15)

Atomic displacement parameters (\AA^2)

U^{11} U^{22} U^{33} U^{12} U^{13} U^{23}

supplementary materials

Br1	0.05493 (19)	0.04386 (16)	0.03946 (16)	0.00297 (12)	-0.00566 (13)	0.01018 (11)
Br2	0.0613 (2)	0.05608 (19)	0.02904 (15)	0.00966 (13)	-0.00356 (12)	-0.00142 (11)
C1	0.0373 (13)	0.0296 (11)	0.0345 (13)	0.0066 (9)	0.0086 (10)	-0.0027 (10)
C2	0.0351 (13)	0.0354 (12)	0.0282 (12)	0.0097 (10)	0.0047 (10)	0.0013 (9)
C3	0.0287 (12)	0.0383 (12)	0.0322 (12)	0.0040 (9)	0.0072 (10)	0.0009 (10)
C4	0.0294 (12)	0.0381 (12)	0.0320 (12)	0.0091 (9)	0.0122 (10)	0.0014 (10)
C5	0.0348 (13)	0.0402 (12)	0.0257 (11)	0.0097 (10)	0.0059 (9)	-0.0032 (10)
C6	0.0368 (13)	0.0375 (12)	0.0351 (13)	0.0034 (10)	0.0045 (11)	-0.0088 (10)
C7	0.0337 (13)	0.0467 (14)	0.0379 (13)	0.0047 (10)	0.0150 (10)	0.0061 (11)
C8	0.0401 (14)	0.0384 (13)	0.0332 (13)	-0.0020 (10)	0.0095 (11)	-0.0027 (10)
C9	0.0348 (12)	0.0311 (11)	0.0268 (11)	0.0025 (9)	0.0123 (9)	-0.0041 (9)
C10	0.0388 (13)	0.0422 (13)	0.0330 (13)	0.0072 (11)	0.0075 (10)	0.0056 (10)
C11	0.0263 (12)	0.0571 (16)	0.0394 (14)	0.0082 (11)	0.0049 (10)	0.0043 (12)
C12	0.0312 (13)	0.0447 (14)	0.0404 (14)	-0.0040 (10)	0.0092 (11)	-0.0022 (11)
C13	0.0314 (12)	0.0345 (12)	0.0283 (12)	0.0030 (9)	0.0105 (9)	-0.0032 (9)
C14	0.0462 (15)	0.0374 (13)	0.0332 (13)	0.0031 (11)	0.0122 (11)	0.0017 (10)
C15	0.0517 (16)	0.0303 (12)	0.0441 (15)	0.0008 (11)	0.0082 (12)	-0.0017 (11)
N1	0.0309 (10)	0.0332 (10)	0.0277 (10)	0.0038 (7)	0.0093 (8)	-0.0038 (8)
S1	0.0464 (4)	0.0343 (3)	0.0559 (4)	-0.0053 (3)	0.0138 (3)	0.0052 (3)
S2	0.0394 (3)	0.0376 (3)	0.0363 (3)	0.0021 (2)	0.0123 (3)	0.0090 (3)

Geometric parameters (\AA , $^\circ$)

Br1—C2	1.903 (2)	C8—H8B	0.9700
Br2—C5	1.895 (2)	C9—N1	1.347 (3)
C1—C2	1.386 (3)	C9—C10	1.379 (3)
C1—C6	1.395 (3)	C10—C11	1.371 (4)
C1—C15	1.498 (4)	C10—H10	0.9300
C2—C3	1.381 (3)	C11—C12	1.377 (4)
C3—C4	1.393 (3)	C11—H11	0.9300
C3—H3	0.9300	C12—C13	1.386 (3)
C4—C5	1.392 (3)	C12—H12	0.9300
C4—C7	1.496 (3)	C13—N1	1.335 (3)
C5—C6	1.380 (3)	C13—C14	1.506 (3)
C6—H6	0.9300	C14—S1	1.820 (3)
C7—S2	1.819 (3)	C14—H14A	0.9700
C7—H7A	0.9700	C14—H14B	0.9700
C7—H7B	0.9700	C15—S1	1.817 (3)
C8—C9	1.505 (3)	C15—H15A	0.9700
C8—S2	1.814 (2)	C15—H15B	0.9700
C8—H8A	0.9700		
C2—C1—C6	116.5 (2)	N1—C9—C10	122.3 (2)
C2—C1—C15	123.3 (2)	N1—C9—C8	116.2 (2)
C6—C1—C15	120.2 (2)	C10—C9—C8	121.5 (2)
C3—C2—C1	122.4 (2)	C11—C10—C9	118.9 (2)
C3—C2—Br1	116.22 (18)	C11—C10—H10	120.6
C1—C2—Br1	121.40 (18)	C9—C10—H10	120.6
C2—C3—C4	120.9 (2)	C10—C11—C12	119.3 (2)
C2—C3—H3	119.6	C10—C11—H11	120.3

C4—C3—H3	119.6	C12—C11—H11	120.3
C5—C4—C3	117.0 (2)	C11—C12—C13	119.0 (2)
C5—C4—C7	124.3 (2)	C11—C12—H12	120.5
C3—C4—C7	118.6 (2)	C13—C12—H12	120.5
C6—C5—C4	121.7 (2)	N1—C13—C12	122.0 (2)
C6—C5—Br2	117.85 (18)	N1—C13—C14	116.7 (2)
C4—C5—Br2	120.44 (19)	C12—C13—C14	121.3 (2)
C5—C6—C1	121.4 (2)	C13—C14—S1	114.34 (18)
C5—C6—H6	119.3	C13—C14—H14A	108.7
C1—C6—H6	119.3	S1—C14—H14A	108.7
C4—C7—S2	115.00 (17)	C13—C14—H14B	108.7
C4—C7—H7A	108.5	S1—C14—H14B	108.7
S2—C7—H7A	108.5	H14A—C14—H14B	107.6
C4—C7—H7B	108.5	C1—C15—S1	114.02 (17)
S2—C7—H7B	108.5	C1—C15—H15A	108.7
H7A—C7—H7B	107.5	S1—C15—H15A	108.7
C9—C8—S2	114.50 (16)	C1—C15—H15B	108.7
C9—C8—H8A	108.6	S1—C15—H15B	108.7
S2—C8—H8A	108.6	H15A—C15—H15B	107.6
C9—C8—H8B	108.6	C13—N1—C9	118.4 (2)
S2—C8—H8B	108.6	C15—S1—C14	103.75 (12)
H8A—C8—H8B	107.6	C8—S2—C7	103.29 (12)
C6—C1—C2—C3	2.6 (3)	S2—C8—C9—C10	-53.5 (3)
C15—C1—C2—C3	-175.3 (2)	N1—C9—C10—C11	-1.6 (4)
C6—C1—C2—Br1	-178.55 (17)	C8—C9—C10—C11	178.4 (2)
C15—C1—C2—Br1	3.6 (3)	C9—C10—C11—C12	-0.8 (4)
C1—C2—C3—C4	-0.2 (4)	C10—C11—C12—C13	1.5 (4)
Br1—C2—C3—C4	-179.17 (17)	C11—C12—C13—N1	0.1 (4)
C2—C3—C4—C5	-2.8 (3)	C11—C12—C13—C14	179.1 (2)
C2—C3—C4—C7	175.2 (2)	N1—C13—C14—S1	-126.8 (2)
C3—C4—C5—C6	3.5 (3)	C12—C13—C14—S1	54.2 (3)
C7—C4—C5—C6	-174.3 (2)	C2—C1—C15—S1	105.8 (2)
C3—C4—C5—Br2	-177.02 (16)	C6—C1—C15—S1	-71.9 (3)
C7—C4—C5—Br2	5.1 (3)	C12—C13—N1—C9	-2.4 (3)
C4—C5—C6—C1	-1.2 (4)	C14—C13—N1—C9	178.6 (2)
Br2—C5—C6—C1	179.31 (18)	C10—C9—N1—C13	3.2 (3)
C2—C1—C6—C5	-1.8 (3)	C8—C9—N1—C13	-176.80 (19)
C15—C1—C6—C5	176.1 (2)	C1—C15—S1—C14	-40.3 (2)
C5—C4—C7—S2	72.8 (3)	C13—C14—S1—C15	83.7 (2)
C3—C4—C7—S2	-105.0 (2)	C9—C8—S2—C7	-83.4 (2)
S2—C8—C9—N1	126.56 (19)	C4—C7—S2—C8	40.9 (2)

Fig. 1

