

(E)-1-(2,2-Dimethoxyethyl)-2-(nitromethylene)imidazolidine

Dongmei Li, Zhongzhen Tian,* Gaolei Wang, Peifeng Wei and Yanming Zhang

Shandong Provincial Key Laboratory of Fluorine Chemistry and Chemical Materials, School of Chemistry and Chemical Engineering, University of Jinan, People's Republic of China

Correspondence e-mail: chm_tianzz@ujn.edu.cn


Received 21 July 2010; accepted 29 July 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.037; wR factor = 0.110; data-to-parameter ratio = 16.8.

In the title compound, $C_8H_{15}N_3O_4$, the 2-(nitromethylene)-imidazolidine fragment is close to being planar (r.m.s. deviation = 0.027 Å), which may be correlated with delocalization of the electrons and the effect of the strongly electron-withdrawing NO_2 group. An intramolecular $N-H \cdots O$ link generates an $S(6)$ ring. The same H atom also forms a weak intermolecular $N-H \cdots O$ hydrogen bond, which results in $C(7)$ chains propagating in $[010]$.

Related literature

For background to neonicotinoid insecticides, see Moriya *et al.* (1992). For the synthesis, see: Tian *et al.* (2007).



Experimental

Crystal data

$C_8H_{15}N_3O_4$

$M_r = 217.23$

Monoclinic, $P2_1/c$

$a = 10.444$ (2) Å

$b = 6.8676$ (17) Å

$c = 14.441$ (3) Å

$\beta = 99.953$ (14)°

$V = 1020.2$ (4) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.11$ mm⁻¹
 $T = 296$ K

$0.32 \times 0.26 \times 0.15$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{min} = 0.965$, $T_{max} = 0.983$

8629 measured reflections
2330 independent reflections
1849 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.018$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.110$
 $S = 1.06$
2330 reflections

139 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 0.25$ e Å⁻³
 $\Delta\rho_{min} = -0.16$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$N2-H2 \cdots O2$	0.86	2.09	2.6394 (17)	121
$N2-H2 \cdots O3^i$	0.86	2.64	3.3554 (16)	141

Symmetry code: (i) $-x, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: WinGX (Farrugia, 1999).

The authors thank the National Natural Science Foundation of China (grant No. 20902037), the Opening Fund of Shanghai Key Laboratory of Chemical Biology (grant No. SKLCB-2008-08) and the Doctoral Foundation of the University of Jinan (B0542) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5567).

References

Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Burla, M. C., Polidori, G., Camalli, M. & Spagna, R. (1999). *SIR97*. Universities of Bari, Perugia and Rome, Italy.
Bruker (2005). *APEX2*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Moriya, K., Shibuya, K., Hattori, Y., Tsuboi, S., Shiokawa, K. & Kagabu, S. (1992). *Biosci. Biotech. Biochem.* **56**, 364–365.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Tian, Z. Z., Shao, X. S., Li, Z., Qian, X. H. & Huang, Q. C. (2007). *J. Agric. Food Chem.* **55**, 2288–2292.

supplementary materials

Acta Cryst. (2010). E66, o2216 [doi:10.1107/S1600536810030266]

(*E*)-1-(2,2-Dimethoxyethyl)-2-(nitromethylidene)imidazolidine

D. Li, Z. Tian, G. Wang, P. Wei and Y. Zhang

Comment

Since the debut of Imidacloprid in 1990s (Moriya *et al.*, 1992), neonicotinoid insecticides have become rapidly an important chemical class of insecticides. Our interest was introducing oxygen atom into the lead struture and synthesizing a series of new compounds, in which the title compound (I) exhibited good insecticidal activities against pea aphids.

The structure of (I) is shown in Fig. 1 with the atom-numbering scheme. The delocalization of the electrons as far as the strong electron-withdrawing group, NO₂, lead to a coplanar olefin-amine π -electron network. Intermolecular hydrogen bonds (N2—H2 \cdots O3) are found, and link the molecules into chains.

Experimental

The title compound was synthesized according to the literature (Tian *et al.*, 2007). Colourless prisms of (I) were obtained by slow evaporation of the solution of dichloromethane and ethyl acetate of the title compound.

Refinement

H atoms bonded to N and O atoms were located in a difference map and refined with distance restraints of N—H = 0.87 (2) Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. Other H atoms were positioned geometrically and refined using a riding model (including free rotation about the ethanol C—C bond), with C—H = 0.95–0.99 Å and with $U_{\text{iso}}(\text{H}) = 1.2$ (1.5 for methyl groups) times $U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of (I) with displacement ellipsoids drawn at the 50% probability level. The H atoms are shown as circles of arbitrary size.




Fig. 2. Intermolecular hydrogen bonding in the crystal structure of (I).

(E)-1-(2,2-Dimethoxyethyl)-2-(nitromethylidene)imidazolidine

Crystal data

$C_8H_{15}N_3O_4$	$F(000) = 464$
$M_r = 217.23$	$D_x = 1.414 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 3493 reflections
$a = 10.444 (2) \text{ \AA}$	$\theta = 2.9\text{--}27.3^\circ$
$b = 6.8676 (17) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$c = 14.441 (3) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 99.953 (14)^\circ$	Prism, colourless
$V = 1020.2 (4) \text{ \AA}^3$	$0.32 \times 0.26 \times 0.15 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD diffractometer	2330 independent reflections
Radiation source: fine-focus sealed tube graphite	1849 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.018$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$\theta_{\text{max}} = 27.6^\circ$, $\theta_{\text{min}} = 2.0^\circ$
$T_{\text{min}} = 0.965$, $T_{\text{max}} = 0.983$	$h = -13 \rightarrow 13$
8629 measured reflections	$k = -8 \rightarrow 8$
	$l = -18 \rightarrow 18$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.037$	H-atom parameters constrained
$wR(F^2) = 0.110$	$w = 1/[\sigma^2(F_o^2) + (0.0508P)^2 + 0.1844P]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
2330 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
139 parameters	$\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.16 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
	Extinction coefficient: 0.014 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.00359 (12)	0.08330 (19)	0.63950 (8)	0.0427 (3)
H1A	0.0562	0.0860	0.5937	0.051*
C2	0.06233 (12)	0.08637 (17)	0.73529 (8)	0.0406 (3)
C3	0.09094 (16)	0.0794 (2)	0.89827 (9)	0.0563 (4)
H3A	0.0816	-0.0387	0.9333	0.068*
H3B	0.0774	0.1913	0.9364	0.068*
C4	0.22254 (15)	0.0880 (2)	0.86762 (9)	0.0597 (4)
H4A	0.2701	0.2036	0.8921	0.072*
H4B	0.2741	-0.0262	0.8888	0.072*
C5	0.29097 (12)	0.08628 (19)	0.70666 (9)	0.0461 (3)
H5A	0.2533	0.0386	0.6447	0.055*
H5B	0.3573	-0.0057	0.7342	0.055*
C6	0.35394 (11)	0.28135 (19)	0.69663 (8)	0.0414 (3)
H6	0.3643	0.3483	0.7574	0.050*
C7	0.31179 (18)	0.5845 (2)	0.62367 (12)	0.0641 (4)
H7A	0.3323	0.6410	0.6853	0.096*
H7B	0.2446	0.6594	0.5860	0.096*
H7C	0.3880	0.5846	0.5948	0.096*
C8	0.48007 (15)	0.1666 (3)	0.58408 (11)	0.0699 (5)
H8A	0.4495	0.0354	0.5873	0.105*
H8B	0.5676	0.1650	0.5719	0.105*
H8C	0.4252	0.2349	0.5343	0.105*
N1	-0.12619 (11)	0.07657 (17)	0.61267 (8)	0.0481 (3)
N2	0.00157 (12)	0.0812 (2)	0.80881 (8)	0.0569 (3)
H2	-0.0816	0.0792	0.8041	0.068*
N3	0.19059 (11)	0.09392 (17)	0.76437 (7)	0.0473 (3)
O1	-0.17315 (10)	0.07294 (17)	0.52580 (7)	0.0654 (3)
O2	-0.20198 (10)	0.0741 (2)	0.67215 (8)	0.0725 (4)
O3	0.26831 (8)	0.39078 (13)	0.63099 (6)	0.0476 (3)
O4	0.47709 (8)	0.26264 (15)	0.67127 (6)	0.0524 (3)

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0413 (6)	0.0489 (7)	0.0367 (6)	-0.0019 (5)	0.0030 (5)	0.0045 (5)
C2	0.0448 (6)	0.0371 (6)	0.0384 (6)	-0.0040 (5)	0.0029 (5)	0.0046 (5)
C3	0.0749 (10)	0.0563 (9)	0.0359 (6)	-0.0084 (7)	0.0049 (6)	0.0040 (6)
C4	0.0647 (9)	0.0703 (10)	0.0380 (7)	-0.0080 (7)	-0.0082 (6)	0.0071 (6)
C5	0.0418 (6)	0.0450 (7)	0.0487 (7)	-0.0006 (5)	0.0001 (5)	-0.0001 (5)
C6	0.0349 (6)	0.0491 (7)	0.0379 (6)	-0.0021 (5)	-0.0001 (4)	-0.0028 (5)
C7	0.0774 (11)	0.0510 (9)	0.0627 (9)	-0.0060 (7)	0.0084 (8)	0.0084 (7)
C8	0.0467 (8)	0.1033 (13)	0.0602 (9)	0.0017 (8)	0.0108 (7)	-0.0211 (9)
N1	0.0461 (6)	0.0522 (7)	0.0437 (6)	-0.0003 (5)	0.0008 (5)	0.0042 (5)
N2	0.0519 (6)	0.0807 (9)	0.0379 (6)	-0.0050 (6)	0.0073 (5)	0.0037 (6)
N3	0.0450 (6)	0.0570 (7)	0.0368 (5)	-0.0083 (5)	-0.0013 (4)	0.0060 (5)
O1	0.0562 (6)	0.0879 (8)	0.0445 (5)	0.0038 (5)	-0.0124 (4)	0.0015 (5)
O2	0.0446 (6)	0.1143 (11)	0.0590 (6)	-0.0025 (6)	0.0096 (5)	0.0061 (6)
O3	0.0438 (5)	0.0477 (5)	0.0479 (5)	-0.0010 (4)	-0.0014 (4)	0.0026 (4)
O4	0.0342 (5)	0.0713 (7)	0.0495 (5)	-0.0050 (4)	0.0010 (4)	-0.0086 (5)

Geometric parameters (\AA , $^\circ$)

C1—N1	1.3447 (17)	C5—H5B	0.9700
C1—C2	1.4131 (17)	C6—O4	1.4028 (15)
C1—H1A	0.9300	C6—O3	1.4036 (15)
C2—N2	1.3282 (17)	C6—H6	0.9800
C2—N3	1.3339 (16)	C7—O3	1.4159 (18)
C3—N2	1.4573 (17)	C7—H7A	0.9600
C3—C4	1.516 (2)	C7—H7B	0.9600
C3—H3A	0.9700	C7—H7C	0.9600
C3—H3B	0.9700	C8—O4	1.4265 (18)
C4—N3	1.4711 (17)	C8—H8A	0.9600
C4—H4A	0.9700	C8—H8B	0.9600
C4—H4B	0.9700	C8—H8C	0.9600
C5—N3	1.4485 (18)	N1—O2	1.2647 (15)
C5—C6	1.5104 (18)	N1—O1	1.2658 (14)
C5—H5A	0.9700	N2—H2	0.8600
N1—C1—C2	121.87 (12)	O4—C6—H6	108.2
N1—C1—H1A	119.1	O3—C6—H6	108.2
C2—C1—H1A	119.1	C5—C6—H6	108.2
N2—C2—N3	110.00 (11)	O3—C7—H7A	109.5
N2—C2—C1	126.55 (12)	O3—C7—H7B	109.5
N3—C2—C1	123.45 (12)	H7A—C7—H7B	109.5
N2—C3—C4	102.41 (11)	O3—C7—H7C	109.5
N2—C3—H3A	111.3	H7A—C7—H7C	109.5
C4—C3—H3A	111.3	H7B—C7—H7C	109.5
N2—C3—H3B	111.3	O4—C8—H8A	109.5
C4—C3—H3B	111.3	O4—C8—H8B	109.5

H3A—C3—H3B	109.2	H8A—C8—H8B	109.5
N3—C4—C3	103.81 (11)	O4—C8—H8C	109.5
N3—C4—H4A	111.0	H8A—C8—H8C	109.5
C3—C4—H4A	111.0	H8B—C8—H8C	109.5
N3—C4—H4B	111.0	O2—N1—O1	119.47 (11)
C3—C4—H4B	111.0	O2—N1—C1	121.53 (11)
H4A—C4—H4B	109.0	O1—N1—C1	119.00 (12)
N3—C5—C6	113.17 (11)	C2—N2—C3	112.80 (12)
N3—C5—H5A	108.9	C2—N2—H2	123.6
C6—C5—H5A	108.9	C3—N2—H2	123.6
N3—C5—H5B	108.9	C2—N3—C5	127.22 (11)
C6—C5—H5B	108.9	C2—N3—C4	110.95 (11)
H5A—C5—H5B	107.8	C5—N3—C4	121.47 (11)
O4—C6—O3	112.25 (10)	C6—O3—C7	112.24 (10)
O4—C6—C5	112.21 (11)	C6—O4—C8	115.70 (10)
O3—C6—C5	107.61 (9)		
N1—C1—C2—N2	0.7 (2)	C1—C2—N3—C5	-4.8 (2)
N1—C1—C2—N3	-179.68 (12)	N2—C2—N3—C4	1.74 (15)
N2—C3—C4—N3	0.09 (15)	C1—C2—N3—C4	-177.89 (12)
N3—C5—C6—O4	158.56 (10)	C6—C5—N3—C2	105.58 (14)
N3—C5—C6—O3	-77.48 (13)	C6—C5—N3—C4	-81.94 (15)
C2—C1—N1—O2	0.51 (19)	C3—C4—N3—C2	-1.09 (15)
C2—C1—N1—O1	-179.59 (12)	C3—C4—N3—C5	-174.68 (12)
N3—C2—N2—C3	-1.71 (16)	O4—C6—O3—C7	-62.38 (14)
C1—C2—N2—C3	177.90 (12)	C5—C6—O3—C7	173.69 (12)
C4—C3—N2—C2	0.95 (16)	O3—C6—O4—C8	-61.01 (16)
N2—C2—N3—C5	174.88 (12)	C5—C6—O4—C8	60.33 (16)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H2 \cdots O2	0.86	2.09	2.6394 (17)	121
N2—H2 \cdots O3 ⁱ	0.86	2.64	3.3554 (16)	141

Symmetry codes: (i) $-x, y-1/2, -z+3/2$.

Fig. 1




Fig. 2

