

## (4*S*,5*S*,6*S*)-4-Hydroxy-3-methoxy-5-methyl-5,6-epoxycyclohex-2-en-1-one

Srinuan Tansuwan, Porntana Chanaprat, Thapong Teerawatananond, Nongnuj Muangsin and Surachai Pornpakakul\*

Research Centre of Bioorganic Chemistry, Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

Correspondence e-mail: nongnuj.j@chula.ac.th


Received 11 July 2010; accepted 2 August 2010

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.041;  $wR$  factor = 0.116; data-to-parameter ratio = 15.8.

The title compound,  $\text{C}_8\text{H}_{10}\text{O}_4$ , was isolated from culture extracts of the endophytic fungus *Xylaria sp.* (PB-30). The cyclohexenone ring exhibits a flattened boat conformation. In the crystal structure, molecules related by translation along the  $b$  axis are linked into chains through  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds. Weak non-classical  $\text{C}-\text{H}\cdots\text{O}$  contacts are also observed in the structure.

### Related literature

For background to the structures of bioactive secondary metabolites from endophytic fungus and their activities, see: Tansuwan *et al.* (2007); Shiono *et al.* (2005); Mitsui *et al.* (2004). For related structures and the assignment of the absolute configuration, see: Mitsui *et al.* (2004); Shiono *et al.* (2005). For puckering parameters, see: Cremer & Pople (1975).



### Experimental

#### Crystal data

$\text{C}_8\text{H}_{10}\text{O}_4$	$V = 798.80(4)\text{ \AA}^3$
$M_r = 170.16$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 4.2208(1)\text{ \AA}$	$\mu = 0.11\text{ mm}^{-1}$
$b = 7.5459(3)\text{ \AA}$	$T = 293\text{ K}$
$c = 25.0802(8)\text{ \AA}$	$0.42 \times 0.40 \times 0.30\text{ mm}$

#### Data collection

Bruker SMART APEXII CCD area-detector diffractometer  
6038 measured reflections

1768 independent reflections  
1551 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.116$   
 $S = 1.07$   
1768 reflections

112 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.32\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.24\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2A $\cdots$ O1 <sup>i</sup>	0.82	1.99	2.8148 (17)	180
C4—H4 $\cdots$ O2 <sup>ii</sup>	0.98	2.54	3.521 (2)	176
C6—H6 $\cdots$ O4 <sup>iii</sup>	0.98	2.56	3.5208 (17)	167

Symmetry codes: (i)  $x, y + 1, z$ ; (ii)  $x - 1, y, z$ ; (iii)  $x - \frac{1}{2}, -y + \frac{1}{2}, -z + 2$ .

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Financial support from the Department of Chemistry, Faculty of Science, Chulalongkorn University, the Rachadapiseksomponj Endowment, Chulalongkorn University, the Thailand research fund (grant No. MRG5280213), the National Center of Excellence for Petroleum, Petrochemicals and Advanced Materials, and the A1–B1 project, Faculty of Science, Chulalongkorn University, is gratefully acknowledged. The Thai Government Stimulus Package 2 (TKK2555), under the Project for Establishment of a Comprehensive Center for Innovative Food, Health Products and Agriculture, is acknowledged for support of the X-ray Crystallographic analysis.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2746).

### References

- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Mitsui, T., Miyake, Y., Kakeya, H., Osada, H. & Kataoka, T. (2004). *J. Immunol.* **172**, 3428–3436.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shiono, Y., Murayama, T., Takahashi, K., Okada, K., Katohda, S. & Ikeda, M. (2005). *Biosci. Biotechnol. Biochem.* **69**, 287–292.
- Tansuwan, S., Pornpakakul, S., Roengsumran, S., Petsom, A., Muangsin, N., Sihanonta, P. & Chaichit, N. (2007). *J. Nat. Prod.* **70**, 1620–1623.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## **supplementary materials**

*Acta Cryst.* (2010). E66, o2263 [doi:10.1107/S1600536810030850]

### (4*S*,5*S*,6*S*)-4-Hydroxy-3-methoxy-5-methyl-5,6-epoxycyclohex-2-en-1-one

**S. Tansuwan, P. Chanaprat, T. Teerawatananond, N. Muangsin and S. Pornpakakul**

#### Comment

Endophytic fungi have been proven to be a rich source of novel structural compounds with interesting biological activities and a high level of biodiversity. In the previous investigations of bioactive compounds produced by an endophytic fungus, *Xylaria* sp (strain PB-30), two antimalarial benzoquinones were isolated (Tansuwan *et al.*, 2007). As a part of our continuing search for anticancer metabolites of this fungus, we found that the title compound is one of major metabolite showing cytotoxicity against various human cell lines, for example breast ductal carcinoma (BT474), human undifferentiated lung carcinoma (CHAGO), human liver hepatoblastoma (HEP-G2), human gastric carcinoma (KATO-3), and human colon adenocarcinoma (SW620) at IC<sub>50</sub> values of 10.51, 11.11, 6.25, 5.61 and 5.31 µg/ml, respectively.

The title compound was previously isolated from the organic extracts of the fungus xylariaceous endophytic fungus (strain YUA-026), and elucidated on the basis of spectroscopic analysis (Shiono *et al.*, 2005). Herein we present the crystal structure of the title compound, which was isolated from the fermentation culture of the endophytic fungus *Xylaria* sp (strain PB-30) at room temperature under a static condition.

The cyclohexenone ring exhibits a flattened boat conformation with puckering amplitudes  $Q = 0.232$  (1),  $\phi = 176.5$  (3)° and  $\theta = 81.8$ °. The 3-methoxy substituent is in the plane of the cyclohexenone ring to which it is attached [dihedral angle = 7.4 (2)°]. Epoxide ring makes dihedral angles of 87.23 (6)° with the cyclohexenone rings C1—C6. The hydroxyl group locates on the same side of the epoxide ring. In the crystal, the molecules are linked to each other through O—H···O hydrogen bonds, building a polymeric chain parallel to the *b*-axis. Weak non-classical C—H···O contacts are also observed in the structure. The absolute configuration could not be determined therefore it was assigned on the basis of literature data (Shiono *et al.*, 2005).

#### Experimental

The endophytic fungus *Xylaria* sp. PB-30 was cultivated in 100 ml of malt extract broth (MEB) in a 250 ml flask (x 300) under static condition at room temperature for 35 days. The culture was filtered through filter paper (Whatman No.1). The culture broth (23 L) was concentrated by rotary evaporator *in vacuo* to give the concentrated broth (1.5 L) and then extracted with EtOAc (x5), CH<sub>2</sub>Cl<sub>2</sub>: MeOH (1:1) (x5) and MeOH (x5), respectively. The solvents were evaporated under reduced pressure at 30°C to give EtOAc crude as yellow viscous liquid (29.61 g), CH<sub>2</sub>Cl<sub>2</sub>: MeOH crude as brown viscous liquid (21.45 g) and MeOH crude as brown viscous liquid (4.74 g).

The EtOAc crude extract (29.61 g) was subjected to a column chromatography [Sephadex™ LH-20 (400 g), column diameter 3.6 cm] using 5% dichloromethane in MeOH as eluent. 10 ml of each fraction was collected. The similar fractions were combined on the basis of TLC profile and monitored by UV, iodine vapor and vanillin/H<sub>2</sub>SO<sub>4</sub> reagent to give 14 combined fractions.

## supplementary materials

The fractions EB-5 and EB-6 were combined and purified by crystallization from a 1:1 mixture of CH<sub>2</sub>Cl<sub>2</sub> and acetone to obtain the title compound as colorless crystals (25.8 mg).

m.p. 153–155 °C; [α]<sub>D</sub><sup>20</sup> -100 ° (c = 0.1, MeOH); λ<sub>max</sub> (MeOH) (log ε) 260 nm (3.75); HRESIMS *m/z* 363.0579 [2M+Na]<sup>+</sup> calc. for (C<sub>8</sub>H<sub>10</sub>O<sub>4</sub>)<sub>2</sub> Na 363.1055 FT—IR (KBr) ν<sub>max</sub> (cm<sup>-1</sup>): 3445, 3070, 3033, 2987, 2940, 2917, 2857, 1642, 1605, 1386, 1300, 1253, 1213, 1070, and 1014; <sup>1</sup>H-NMR δ (CDCl<sub>3</sub>, 400 MHz) 1.64 (3H, s, Me), 2.58 (1H, d, *J*=6.0 Hz, 4-OH), 3.32 (1H, d, *J*=2.0 Hz, 6-H), 3.76 (3H, s, OMe), 4.48 (1H, d, *J*=6.0 Hz, 4-H) and 5.25 (1H, d, *J*= 2.0 Hz, 2-H) p.p.m.; <sup>13</sup>C-NMR δ (CDCl<sub>3</sub>, 100 MHz) 18.9 (Me), 56.6 (OMe), 59.4 (C-5), 60.5 (C-6), 69.1 (C-4), 98.2 (C-2), 171.3 (C-3) and 193.4 (C-1) p.p.m.

### Refinement

All H atoms were positioned geometrically and treated as riding, with C—H bonding lengths constrained to 0.93 Å (aromatic CH), 0.96 Å (methyl CH<sub>3</sub>), 0.97 Å (methylene CH<sub>2</sub>), and O—H = 0.84 Å, and with *U*<sub>iso</sub>(H) = 1.2*U*<sub>eq</sub>(CH) or *U*<sub>iso</sub>(H) = 1.5*U*<sub>eq</sub>(CH<sub>3</sub>, methylene C or OH). 1159 Friedel pair were merged before the final refinement. The absolute configuration was assigned on the basis of the known configuration of the starting material as *R*, *S* and *S* for C4, C5 and C6-positions, respectively.

### Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atom numbering scheme. Hydrogen atoms are shown as spheres of arbitrary radius.

### (4*S*,5*S*,6*S*)-4-Hydroxy-3-methoxy-5-methyl-5,6- epoxycyclohex-2-en-1-one

#### Crystal data

C <sub>8</sub> H <sub>10</sub> O <sub>4</sub>	<i>F</i> (000) = 360
<i>M<sub>r</sub></i> = 170.16	<i>D<sub>x</sub></i> = 1.415 Mg m <sup>-3</sup>
Orthorhombic, <i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Mo <i>K</i> α radiation, $\lambda$ = 0.71073 Å
Hall symbol: P 2ac 2ab	Cell parameters from 2855 reflections
<i>a</i> = 4.2208 (1) Å	$\theta$ = 2.8–32.8°
<i>b</i> = 7.5459 (3) Å	$\mu$ = 0.11 mm <sup>-1</sup>
<i>c</i> = 25.0802 (8) Å	<i>T</i> = 293 K
<i>V</i> = 798.80 (4) Å <sup>3</sup>	Prism, colourless
<i>Z</i> = 4	0.42 × 0.40 × 0.30 mm

#### Data collection

Bruker SMART APEXII CCD area-detector  
diffractometer 1551 reflections with *I* > 2σ(*I*)

Radiation source: Mo	$R_{\text{int}} = 0.021$
graphite	$\theta_{\text{max}} = 33.1^\circ, \theta_{\text{min}} = 2.8^\circ$
$\varphi$ and $\omega$ scans	$h = -6 \rightarrow 6$
6038 measured reflections	$k = -11 \rightarrow 7$
1768 independent reflections	$l = -37 \rightarrow 32$

*Refinement*

Refinement on $F^2$	0 restraints
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.041$	$w = 1/[\sigma^2(F_o^2) + (0.0798P)^2 + 0.0232P]$
	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.116$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.07$	$\Delta\rho_{\text{max}} = 0.32 \text{ e \AA}^{-3}$
1768 reflections	$\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$
112 parameters	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6030 (4)	0.12154 (17)	0.88766 (6)	0.0349 (3)
C2	0.6543 (4)	0.19049 (16)	0.83465 (5)	0.0337 (3)
H2	0.6778	0.1121	0.8063	0.04*
C3	0.6687 (3)	0.36664 (16)	0.82548 (5)	0.0268 (2)
C4	0.6073 (3)	0.50632 (15)	0.86741 (5)	0.0248 (2)
H4	0.3894	0.5489	0.8632	0.03*
C5	0.6459 (3)	0.43752 (15)	0.92384 (5)	0.0242 (2)
C6	0.6377 (4)	0.24623 (17)	0.93348 (5)	0.0293 (3)
H6	0.5586	0.2064	0.9682	0.035*
C7	0.5554 (5)	0.56563 (19)	0.96737 (5)	0.0372 (3)
H7A	0.678	0.6721	0.9638	0.056*
H7B	0.5964	0.5128	1.0015	0.056*
H7C	0.3342	0.5936	0.9645	0.056*
C8	0.7980 (6)	0.3240 (2)	0.73390 (6)	0.0452 (4)
H8A	0.6217	0.2477	0.7264	0.068*
H8B	0.9791	0.2537	0.7433	0.068*
H8C	0.8461	0.3938	0.7029	0.068*
O1	0.5405 (5)	-0.03468 (14)	0.89674 (5)	0.0594 (5)
O2	0.8156 (3)	0.65036 (13)	0.85837 (4)	0.0392 (3)
H2A	0.735	0.7419	0.8696	0.059*

## supplementary materials

---

O3	0.7187 (4)	0.43952 (13)	0.77758 (4)	0.0379 (3)
O4	0.9367 (3)	0.34098 (15)	0.93283 (4)	0.0350 (2)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0486 (8)	0.0182 (5)	0.0379 (6)	0.0011 (5)	0.0071 (7)	-0.0006 (4)
C2	0.0484 (8)	0.0210 (5)	0.0318 (6)	-0.0004 (5)	0.0055 (6)	-0.0066 (4)
C3	0.0325 (6)	0.0226 (5)	0.0253 (5)	0.0013 (5)	0.0029 (5)	-0.0029 (4)
C4	0.0297 (5)	0.0174 (4)	0.0272 (5)	0.0020 (4)	0.0042 (4)	-0.0011 (4)
C5	0.0254 (5)	0.0211 (5)	0.0261 (5)	-0.0009 (4)	0.0018 (4)	-0.0029 (4)
C6	0.0360 (6)	0.0231 (5)	0.0289 (5)	0.0008 (5)	0.0029 (5)	0.0024 (4)
C7	0.0521 (9)	0.0286 (6)	0.0310 (6)	-0.0041 (6)	0.0082 (6)	-0.0093 (5)
C8	0.0645 (11)	0.0422 (8)	0.0290 (6)	0.0011 (9)	0.0103 (7)	-0.0080 (6)
O1	0.1057 (14)	0.0182 (4)	0.0545 (7)	-0.0065 (6)	0.0200 (9)	0.0002 (4)
O2	0.0545 (7)	0.0199 (4)	0.0431 (5)	-0.0081 (4)	0.0158 (5)	-0.0024 (4)
O3	0.0593 (7)	0.0294 (4)	0.0250 (4)	0.0029 (5)	0.0074 (5)	-0.0019 (3)
O4	0.0280 (5)	0.0368 (5)	0.0400 (5)	0.0022 (4)	-0.0047 (4)	0.0019 (4)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C1—O1	1.2292 (17)	C5—C7	1.5074 (17)
C1—C2	1.444 (2)	C6—O4	1.4506 (18)
C1—C6	1.4924 (19)	C6—H6	0.98
C2—C3	1.3503 (17)	C7—H7A	0.96
C2—H2	0.93	C7—H7B	0.96
C3—O3	1.3379 (15)	C7—H7C	0.96
C3—C4	1.5113 (16)	C8—O3	1.4393 (17)
C4—O2	1.4162 (17)	C8—H8A	0.96
C4—C5	1.5162 (17)	C8—H8B	0.96
C4—H4	0.98	C8—H8C	0.96
C5—O4	1.4448 (16)	O2—H2A	0.82
C5—C6	1.4640 (18)		
O1—C1—C2	123.24 (14)	O4—C6—C5	59.43 (9)
O1—C1—C6	118.88 (14)	O4—C6—C1	112.79 (12)
C2—C1—C6	117.85 (12)	C5—C6—C1	119.79 (11)
C3—C2—C1	121.21 (11)	O4—C6—H6	117.2
C3—C2—H2	119.4	C5—C6—H6	117.2
C1—C2—H2	119.4	C1—C6—H6	117.2
O3—C3—C2	124.38 (11)	C5—C7—H7A	109.5
O3—C3—C4	111.41 (10)	C5—C7—H7B	109.5
C2—C3—C4	124.08 (11)	H7A—C7—H7B	109.5
O2—C4—C3	108.50 (10)	C5—C7—H7C	109.5
O2—C4—C5	110.21 (11)	H7A—C7—H7C	109.5
C3—C4—C5	113.09 (10)	H7B—C7—H7C	109.5
O2—C4—H4	108.3	O3—C8—H8A	109.5
C3—C4—H4	108.3	O3—C8—H8B	109.5
C5—C4—H4	108.3	H8A—C8—H8B	109.5

O4—C5—C6	59.82 (9)	O3—C8—H8C	109.5
O4—C5—C7	115.19 (12)	H8A—C8—H8C	109.5
C6—C5—C7	120.44 (11)	H8B—C8—H8C	109.5
O4—C5—C4	114.18 (10)	C4—O2—H2A	109.5
C6—C5—C4	119.29 (10)	C3—O3—C8	118.11 (11)
C7—C5—C4	115.41 (11)	C5—O4—C6	60.75 (8)
O1—C1—C2—C3	168.6 (2)	C7—C5—C6—O4	-103.26 (15)
C6—C1—C2—C3	-13.3 (3)	C4—C5—C6—O4	102.54 (13)
C1—C2—C3—O3	179.30 (16)	O4—C5—C6—C1	-100.39 (15)
C1—C2—C3—C4	-5.2 (3)	C7—C5—C6—C1	156.36 (15)
O3—C3—C4—O2	-40.34 (16)	C4—C5—C6—C1	2.2 (2)
C2—C3—C4—O2	143.68 (16)	O1—C1—C6—O4	125.98 (19)
O3—C3—C4—C5	-162.94 (13)	C2—C1—C6—O4	-52.22 (19)
C2—C3—C4—C5	21.1 (2)	O1—C1—C6—C5	-167.29 (19)
O2—C4—C5—O4	-72.54 (13)	C2—C1—C6—C5	14.5 (2)
C3—C4—C5—O4	49.10 (15)	C2—C3—O3—C8	-7.4 (3)
O2—C4—C5—C6	-140.21 (13)	C4—C3—O3—C8	176.65 (16)
C3—C4—C5—C6	-18.57 (18)	C7—C5—O4—C6	111.98 (13)
O2—C4—C5—C7	64.33 (16)	C4—C5—O4—C6	-111.05 (12)
C3—C4—C5—C7	-174.03 (13)	C1—C6—O4—C5	112.19 (13)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H2A···O1 <sup>i</sup>	0.82	1.99	2.8148 (17)	180
C4—H4···O2 <sup>ii</sup>	0.98	2.54	3.521 (2)	176
C6—H6···O4 <sup>iii</sup>	0.98	2.56	3.5208 (17)	167

Symmetry codes: (i)  $x, y+1, z$ ; (ii)  $x-1, y, z$ ; (iii)  $x-1/2, -y+1/2, -z+2$ .

## supplementary materials

Fig. 1

