

5-Chloro-3-ethylsulfinyl-2-(4-iodophenyl)-7-methyl-1-benzofuran

Hong Dae Choi,^a Pil Ja Seo,^a Byeng Wha Son^b and Uk Lee^{b*}

^aDepartment of Chemistry, Dongeui University, San 24 Kaya-dong Busanjin-gu, Busan 614-714, Republic of Korea, and ^bDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong Nam-gu, Busan 608-737, Republic of Korea
Correspondence e-mail: uklee@pknu.ac.kr


Received 2 August 2010; accepted 5 August 2010

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; disorder in main residue; R factor = 0.024; wR factor = 0.065; data-to-parameter ratio = 17.1.

In the title compound, $C_{17}H_{14}ClO_2S$, the 4-iodophenyl ring makes a dihedral angle of $1.61(9)^\circ$ with the benzofuran ring system. In the crystal, molecules are linked through a weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond and an $\text{I}\cdots\text{O}$ contact [$3.416(2)\text{ \AA}$]. The ethyl group is disordered over two orientations with site-occupancy factors of 0.402(7) and 0.598(7).

Related literature

For the pharmacological activity of benzofuran compounds, see: Aslam *et al.* (2006); Galal *et al.* (2009); Khan *et al.* (2005). For natural products with benzofuran rings, see: Akgul & Anil (2003); Soekamto *et al.* (2003). For the structures of related 3-ethylsulfinyl-2-(4-iodophenyl)-1-benzofuran derivatives, see: Choi *et al.* (2010a,b). For a review of halogen bonding, see: Politzer *et al.* (2007).



Experimental

Crystal data

$C_{17}H_{14}ClO_2S$

$M_r = 444.69$

Triclinic, $P\bar{1}$
 $a = 7.8013(2)\text{ \AA}$
 $b = 10.4240(3)\text{ \AA}$
 $c = 11.6003(3)\text{ \AA}$
 $\alpha = 115.962(1)^\circ$
 $\beta = 98.040(1)^\circ$
 $\gamma = 96.661(1)^\circ$
 $V = 823.00(4)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 2.24\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.48 \times 0.34 \times 0.16\text{ mm}$

Data collection

Bruker SMART APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2009)
 $R_{\text{int}} = 0.024$
 $T_{\text{min}} = 0.473$, $T_{\text{max}} = 0.746$
14721 measured reflections
3773 independent reflections
3408 reflections with $I > 2\sigma(I)$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.024$
 $wR(F^2) = 0.065$
 $S = 1.06$
3773 reflections
221 parameters
28 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.28\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.60\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|--|--------------|--------------------|-------------|----------------------|
| $\text{C}16\text{A}-\text{H}16\text{B}\cdots\text{O}2^i$ | 0.97 | 2.40 | 3.297 (6) | 154 |

Symmetry code: (i) $-x, -y, -z$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 1998); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2588).

References

- Akgul, Y. Y. & Anil, H. (2003). *Phytochemistry*, **63**, 939–943.
- Aslam, S. N., Stevenson, P. C., Phythian, S. J., Veitch, N. C. & Hall, D. R. (2006). *Tetrahedron*, **62**, 4214–4226.
- Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2009). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2010a). *Acta Cryst. E66*, o1862.
- Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2010b). *Acta Cryst. E66*, o1985.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Galal, S. A., Abd El-All, A. S., Abdallah, M. M. & El-Diwani, H. I. (2009). *Bioorg. Med. Chem. Lett.* **19**, 2420–2428.
- Khan, M. W., Alam, M. J., Rashid, M. A. & Chowdhury, R. (2005). *Bioorg. Med. Chem.* **13**, 4796–4805.
- Politzer, P., Lane, P., Concha, M. C., Ma, Y. & Murray, J. S. (2007). *J. Mol. Model.* **13**, 305–311.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Soekamto, N. H., Achmad, S. A., Ghisalberti, E. L., Hakim, E. H. & Syah, Y. M. (2003). *Phytochemistry*, **64**, 831–834.

supplementary materials

Acta Cryst. (2010). E66, o2277 [doi:10.1107/S160053681003134X]

5-Chloro-3-ethylsulfinyl-2-(4-iodophenyl)-7-methyl-1-benzofuran

H. D. Choi, P. J. Seo, B. W. Son and U. Lee

Comment

Many compounds containing a benzofuran ring have received considerable attention in view of their pharmacological properties such as antifungal, antitumor and antiviral, antimicrobial activities (Aslam *et al.*, 2006, Galal *et al.*, 2009, Khan *et al.*, 2005). These compounds widely occur in nature (Akgul & Anil, 2003; Soekamto *et al.*, 2003). As a part of our continuing studies of the substituent effect on the solid state structures of 3-ethylsulfinyl-2-(4-iodophenyl)-1-benzofuran analogues (Choi *et al.*, 2010*a, b*), we report the crystal structure of the title compound (Fig. 1).

The benzofuran unit is essentially planar, with a mean deviation of 0.007 (2) Å from the least-squares plane defined by the nine constituent atoms. The ethyl group is disordered over two positions with site-occupancy factors of 0.402 (7) (for atom labelled A) and 0.598 (7) (for atom labelled B) in Fig. 1. The dihedral angle formed by the benzofuran plane and the 4-iodophenyl ring is 1.61 (9)°. The molecular packing (Fig. 2) is stabilized by a weak intermolecular C—H···O hydrogen bond between the methylene H atom of the ethyl group and the oxygen of the S=O unit, with a C16A—H16B···O2ⁱ (Table 1). The crystal packing (Fig. 2) is further stabilized by an I···O halogen-bonding between the iodine and the oxygen of the S=O unit [I···O2ⁱⁱ = 3.416 (2) Å; C12—I···O2ⁱⁱ = 161.46 (7)°] (Politzer *et al.*, 2007).

Experimental

77% 3-chloroperoxybenzoic acid (157 mg, 0.7 mmol) was added in small portions to a stirred solution of 5-chloro-3-ethylsulfinyl-2-(4-iodophenyl)-7-methyl-1-benzofuran (300 mg, 0.7 mmol) in dichloromethane (40 mL) at 273 K. After being stirred at room temperature for 5 h, the mixture was washed with saturated sodium bicarbonate solution and the organic layer was separated, dried over magnesium sulfate, filtered and concentrated at reduced pressure. The residue was purified by column chromatography (hexane–ethyl acetate, 1:1 v/v) to afford the title compound as a colorless solid [yield 76%, m.p. 457–458 K; R_f = 0.64 (hexane–ethyl acetate, 1:1 v/v)]. Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in acetone at room temperature.

Refinement

All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93 Å for aryl, 0.97 Å for methylene, and 0.96 Å for methyl H atoms, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aryl and methylene H atoms, and $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms. The ethyl group was found to be disordered over two positions and modelled with site-occupancy factors, from refinement of 0.402 (7) (part A) and 0.598 (7) (part B), respectively. The displacement ellipsoids of part B were restrained using command ISOR (0.01), sets of C atoms were restrained using command DELU and the distances of C—C were restrained to 1.480 (2) Å using command DFIX. The distances of C—S and C—C were restrained to 0.001 Å using command SADI.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as a small spheres of arbitrary radius. The ethyl group is disordered over two positions with site-occupancy factors, from refinement of 0.402 (7) and 0.598 (7).




Fig. 2. C—H···O and I···O interactions (dotted lines) in the crystal structure of the title compound. The disordered component of the ethyl group, part B, has been omitted for clarity as have H atoms not involved in intermolecular contacts. [Symmetry codes: (i) -x, -y, -z; (ii) x + 1, y + 1, z + 1; (iii) x - 1, y - 1, z - 1.]

5-Chloro-3-ethylsulfinyl-2-(4-iodophenyl)-7-methyl-1-benzofuran

Crystal data

| | |
|---|--|
| C ₁₇ H ₁₄ ClIO ₂ S | Z = 2 |
| M _r = 444.69 | F(000) = 436 |
| Triclinic, P $\bar{1}$ | D _x = 1.794 Mg m ⁻³ |
| Hall symbol: -P 1 | Mo K α radiation, λ = 0.71073 Å |
| a = 7.8013 (2) Å | Cell parameters from 9231 reflections |
| b = 10.4240 (3) Å | θ = 2.2–27.5° |
| c = 11.6003 (3) Å | μ = 2.24 mm ⁻¹ |
| α = 115.962 (1)° | T = 173 K |
| β = 98.040 (1)° | Block, colourless |
| γ = 96.661 (1)° | 0.48 × 0.34 × 0.16 mm |
| V = 823.00 (4) Å ³ | |

Data collection

| | |
|---|--|
| Bruker SMART APEXII CCD diffractometer | 3773 independent reflections |
| Radiation source: rotating anode graphite multilayer | 3408 reflections with $I > 2\sigma(I)$ |
| Detector resolution: 10.0 pixels mm ⁻¹ | $R_{\text{int}} = 0.024$ |
| φ and ω scans | $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 2.0^\circ$ |
| Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009) | $h = -9 \rightarrow 10$ |
| $T_{\text{min}} = 0.473$, $T_{\text{max}} = 0.746$ | $k = -13 \rightarrow 13$ |
| 14721 measured reflections | $l = -15 \rightarrow 15$ |

Refinement

| | |
|----------------------------|--|
| Refinement on F^2 | Primary atom site location: structure-invariant direct methods |
| Least-squares matrix: full | Secondary atom site location: difference Fourier map |

| | |
|---------------------------------|--|
| $R[F^2 > 2\sigma(F^2)] = 0.024$ | Hydrogen site location: difference Fourier map |
| $wR(F^2) = 0.065$ | H-atom parameters constrained |
| $S = 1.06$ | $w = 1/[\sigma^2(F_o^2) + (0.034P)^2 + 0.2748P]$ where $P = (F_o^2 + 2F_c^2)/3$ |
| 3773 reflections | $(\Delta/\sigma)_{\max} = 0.001$ |
| 221 parameters | $\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$ |
| 28 restraints | $\Delta\rho_{\min} = -0.60 \text{ e } \text{\AA}^{-3}$ |

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ | Occ. (<1) |
|-----|--------------|---------------|---------------|----------------------------------|-----------|
| I | 0.64468 (2) | 0.843070 (18) | 0.633828 (13) | 0.05745 (7) | |
| Cl | -0.13606 (8) | 0.12905 (7) | -0.54903 (5) | 0.06029 (16) | |
| S | 0.05956 (8) | 0.20738 (7) | 0.00343 (6) | 0.05340 (14) | |
| O1 | 0.26320 (19) | 0.55334 (15) | -0.03279 (13) | 0.0397 (3) | |
| O2 | -0.1151 (3) | 0.1273 (2) | -0.0800 (2) | 0.0830 (7) | |
| C3 | -0.0084 (3) | 0.2242 (2) | -0.2921 (2) | 0.0432 (4) | |
| H3 | -0.0663 | 0.1352 | -0.3031 | 0.052* | |
| C1 | 0.1344 (3) | 0.3441 (2) | -0.04020 (19) | 0.0398 (4) | |
| C2 | 0.0905 (3) | 0.3316 (2) | -0.17052 (19) | 0.0383 (4) | |
| C4 | -0.0153 (3) | 0.2577 (2) | -0.3946 (2) | 0.0440 (4) | |
| C5 | 0.0668 (3) | 0.3905 (2) | -0.3815 (2) | 0.0450 (5) | |
| H5 | 0.0567 | 0.4065 | -0.4547 | 0.054* | |
| C6 | 0.1637 (3) | 0.4998 (2) | -0.2612 (2) | 0.0413 (4) | |
| C7 | 0.1723 (3) | 0.4632 (2) | -0.15913 (18) | 0.0377 (4) | |
| C8 | 0.2378 (3) | 0.4790 (2) | 0.03875 (18) | 0.0384 (4) | |
| C9 | 0.3273 (3) | 0.5601 (2) | 0.17695 (19) | 0.0396 (4) | |
| C10 | 0.3167 (3) | 0.5045 (3) | 0.2654 (2) | 0.0498 (5) | |
| H10 | 0.2503 | 0.4120 | 0.2370 | 0.060* | |
| C11 | 0.4045 (3) | 0.5857 (3) | 0.3958 (2) | 0.0512 (5) | |
| H11 | 0.3943 | 0.5481 | 0.4545 | 0.061* | |
| C12 | 0.5061 (3) | 0.7208 (2) | 0.4386 (2) | 0.0446 (5) | |
| C13 | 0.5151 (4) | 0.7794 (3) | 0.3532 (2) | 0.0611 (7) | |
| H13 | 0.5815 | 0.8721 | 0.3825 | 0.073* | |
| C14 | 0.4253 (4) | 0.7001 (3) | 0.2240 (2) | 0.0590 (6) | |

supplementary materials

| | | | | | |
|------|-------------|------------|-------------|-------------|-----------|
| H14 | 0.4303 | 0.7410 | 0.1673 | 0.071* | |
| C15 | 0.2513 (4) | 0.6454 (3) | -0.2418 (3) | 0.0580 (6) | |
| H15A | 0.3745 | 0.6468 | -0.2421 | 0.087* | |
| H15B | 0.2385 | 0.7197 | -0.1593 | 0.087* | |
| H15C | 0.1971 | 0.6631 | -0.3116 | 0.087* | |
| C16A | 0.2618 (6) | 0.1332 (6) | 0.0089 (5) | 0.0561 (19) | 0.402 (7) |
| H16A | 0.3649 | 0.2108 | 0.0570 | 0.067* | 0.402 (7) |
| H16B | 0.2543 | 0.0697 | 0.0498 | 0.067* | 0.402 (7) |
| C17A | 0.2695 (15) | 0.0509 (8) | -0.1310 (5) | 0.074 (2) | 0.402 (7) |
| H17A | 0.1592 | -0.0159 | -0.1789 | 0.112* | 0.402 (7) |
| H17B | 0.3633 | -0.0024 | -0.1382 | 0.112* | 0.402 (7) |
| H17C | 0.2909 | 0.1174 | -0.1665 | 0.112* | 0.402 (7) |
| C16B | 0.1842 (6) | 0.0735 (5) | -0.0927 (4) | 0.0586 (13) | 0.598 (7) |
| H16C | 0.1382 | -0.0208 | -0.0998 | 0.070* | 0.598 (7) |
| H16D | 0.1686 | 0.0643 | -0.1806 | 0.070* | 0.598 (7) |
| C17B | 0.3749 (6) | 0.1176 (7) | -0.0305 (6) | 0.0832 (18) | 0.598 (7) |
| H17D | 0.4174 | 0.2156 | -0.0140 | 0.125* | 0.598 (7) |
| H17E | 0.4386 | 0.0533 | -0.0882 | 0.125* | 0.598 (7) |
| H17F | 0.3920 | 0.1126 | 0.0508 | 0.125* | 0.598 (7) |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|------|--------------|--------------|-------------|--------------|--------------|-------------|
| I | 0.05606 (11) | 0.06525 (12) | 0.03984 (9) | -0.00242 (7) | -0.00008 (6) | 0.02082 (7) |
| Cl | 0.0624 (3) | 0.0565 (3) | 0.0396 (3) | 0.0026 (3) | 0.0005 (2) | 0.0076 (2) |
| S | 0.0596 (3) | 0.0483 (3) | 0.0604 (3) | 0.0012 (3) | 0.0147 (3) | 0.0339 (3) |
| O1 | 0.0480 (7) | 0.0332 (7) | 0.0359 (7) | 0.0025 (6) | 0.0051 (5) | 0.0165 (5) |
| O2 | 0.0949 (15) | 0.0656 (12) | 0.0693 (12) | -0.0320 (11) | -0.0043 (11) | 0.0322 (10) |
| C3 | 0.0463 (10) | 0.0333 (10) | 0.0431 (10) | 0.0038 (8) | 0.0085 (8) | 0.0130 (8) |
| C1 | 0.0467 (10) | 0.0359 (10) | 0.0403 (10) | 0.0077 (8) | 0.0110 (8) | 0.0202 (8) |
| C2 | 0.0414 (10) | 0.0341 (9) | 0.0392 (9) | 0.0078 (8) | 0.0097 (8) | 0.0163 (8) |
| C4 | 0.0418 (10) | 0.0425 (11) | 0.0356 (9) | 0.0073 (8) | 0.0036 (8) | 0.0090 (8) |
| C5 | 0.0496 (11) | 0.0497 (12) | 0.0378 (10) | 0.0118 (9) | 0.0089 (8) | 0.0218 (9) |
| C6 | 0.0459 (10) | 0.0390 (10) | 0.0428 (10) | 0.0083 (8) | 0.0113 (8) | 0.0217 (8) |
| C7 | 0.0404 (9) | 0.0343 (9) | 0.0358 (9) | 0.0059 (7) | 0.0067 (7) | 0.0146 (7) |
| C8 | 0.0446 (10) | 0.0359 (10) | 0.0387 (9) | 0.0088 (8) | 0.0107 (8) | 0.0199 (8) |
| C9 | 0.0414 (10) | 0.0376 (10) | 0.0379 (9) | 0.0068 (8) | 0.0071 (7) | 0.0163 (8) |
| C10 | 0.0586 (13) | 0.0419 (11) | 0.0452 (11) | -0.0001 (10) | 0.0009 (9) | 0.0222 (9) |
| C11 | 0.0588 (13) | 0.0525 (13) | 0.0437 (11) | 0.0043 (10) | 0.0028 (9) | 0.0273 (10) |
| C12 | 0.0428 (10) | 0.0503 (12) | 0.0363 (9) | 0.0061 (9) | 0.0046 (8) | 0.0179 (9) |
| C13 | 0.0743 (16) | 0.0502 (13) | 0.0463 (12) | -0.0156 (12) | -0.0007 (11) | 0.0216 (10) |
| C14 | 0.0800 (16) | 0.0503 (13) | 0.0409 (11) | -0.0101 (12) | 0.0022 (11) | 0.0246 (10) |
| C15 | 0.0709 (15) | 0.0502 (13) | 0.0551 (13) | -0.0005 (11) | 0.0093 (11) | 0.0306 (11) |
| C16A | 0.076 (4) | 0.041 (3) | 0.050 (3) | 0.005 (3) | 0.002 (3) | 0.025 (3) |
| C17A | 0.112 (6) | 0.057 (4) | 0.062 (4) | 0.028 (4) | 0.015 (4) | 0.033 (3) |
| C16B | 0.084 (3) | 0.044 (2) | 0.051 (3) | 0.017 (2) | 0.019 (2) | 0.0216 (19) |
| C17B | 0.081 (4) | 0.094 (4) | 0.099 (4) | 0.037 (3) | 0.035 (3) | 0.057 (3) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|-----------------------|-------------|----------------|-------------|
| I—C12 | 2.095 (2) | C10—C11 | 1.389 (3) |
| I—O2 ⁱ | 3.416 (2) | C10—H10 | 0.9300 |
| Cl—C4 | 1.739 (2) | C11—C12 | 1.370 (3) |
| S—O2 | 1.466 (2) | C11—H11 | 0.9300 |
| S—C1 | 1.774 (2) | C12—C13 | 1.378 (3) |
| S—C16A | 1.841 (4) | C13—C14 | 1.381 (3) |
| S—C16B | 1.842 (4) | C13—H13 | 0.9300 |
| O1—C7 | 1.372 (2) | C14—H14 | 0.9300 |
| O1—C8 | 1.379 (2) | C15—H15A | 0.9600 |
| C3—C4 | 1.374 (3) | C15—H15B | 0.9600 |
| C3—C2 | 1.397 (3) | C15—H15C | 0.9600 |
| C3—H3 | 0.9300 | C16A—C17A | 1.4807 (15) |
| C1—C8 | 1.368 (3) | C16A—H16A | 0.9700 |
| C1—C2 | 1.447 (3) | C16A—H16B | 0.9700 |
| C2—C7 | 1.386 (3) | C17A—H17A | 0.9600 |
| C4—C5 | 1.390 (3) | C17A—H17B | 0.9600 |
| C5—C6 | 1.389 (3) | C17A—H17C | 0.9600 |
| C5—H5 | 0.9300 | C16B—C17B | 1.4812 (15) |
| C6—C7 | 1.391 (3) | C16B—H16C | 0.9700 |
| C6—C15 | 1.495 (3) | C16B—H16D | 0.9700 |
| C8—C9 | 1.461 (3) | C17B—H17D | 0.9600 |
| C9—C10 | 1.389 (3) | C17B—H17E | 0.9600 |
| C9—C14 | 1.392 (3) | C17B—H17F | 0.9600 |
| C12—I—O2 ⁱ | 161.46 (7) | C12—C11—C10 | 120.4 (2) |
| O2—S—C1 | 108.18 (11) | C12—C11—H11 | 119.8 |
| O2—S—C16A | 126.7 (2) | C10—C11—H11 | 119.8 |
| C1—S—C16A | 99.2 (2) | C11—C12—C13 | 119.9 (2) |
| O2—S—C16B | 96.09 (17) | C11—C12—I | 121.10 (16) |
| C1—S—C16B | 96.62 (17) | C13—C12—I | 119.02 (16) |
| C7—O1—C8 | 106.81 (15) | C12—C13—C14 | 119.8 (2) |
| C4—C3—C2 | 116.20 (19) | C12—C13—H13 | 120.1 |
| C4—C3—H3 | 121.9 | C14—C13—H13 | 120.1 |
| C2—C3—H3 | 121.9 | C13—C14—C9 | 121.3 (2) |
| C8—C1—C2 | 106.99 (17) | C13—C14—H14 | 119.3 |
| C8—C1—S | 128.36 (16) | C9—C14—H14 | 119.3 |
| C2—C1—S | 124.63 (15) | C6—C15—H15A | 109.5 |
| C7—C2—C3 | 119.59 (18) | C6—C15—H15B | 109.5 |
| C7—C2—C1 | 105.15 (16) | H15A—C15—H15B | 109.5 |
| C3—C2—C1 | 135.26 (19) | C6—C15—H15C | 109.5 |
| C3—C4—C5 | 123.51 (19) | H15A—C15—H15C | 109.5 |
| C3—C4—C1 | 118.70 (17) | H15B—C15—H15C | 109.5 |
| C5—C4—C1 | 117.78 (17) | C17A—C16A—S | 103.8 (4) |
| C6—C5—C4 | 121.43 (19) | C17A—C16A—H16A | 111.0 |
| C6—C5—H5 | 119.3 | S—C16A—H16A | 111.0 |
| C4—C5—H5 | 119.3 | C17A—C16A—H16B | 111.0 |

supplementary materials

| | | | |
|--------------|--------------|----------------------------|--------------|
| C5—C6—C7 | 114.33 (19) | S—C16A—H16B | 111.0 |
| C5—C6—C15 | 123.3 (2) | H16A—C16A—H16B | 109.0 |
| C7—C6—C15 | 122.3 (2) | C17B—C16B—S | 111.2 (3) |
| O1—C7—C2 | 110.77 (17) | C17B—C16B—H16C | 109.4 |
| O1—C7—C6 | 124.31 (18) | S—C16B—H16C | 109.4 |
| C2—C7—C6 | 124.93 (18) | C17B—C16B—H16D | 109.4 |
| C1—C8—O1 | 110.28 (16) | S—C16B—H16D | 109.4 |
| C1—C8—C9 | 135.71 (19) | H16C—C16B—H16D | 108.0 |
| O1—C8—C9 | 114.00 (16) | C16B—C17B—H17D | 109.5 |
| C10—C9—C14 | 117.86 (19) | C16B—C17B—H17E | 109.5 |
| C10—C9—C8 | 122.76 (19) | H17D—C17B—H17E | 109.5 |
| C14—C9—C8 | 119.37 (19) | C16B—C17B—H17F | 109.5 |
| C9—C10—C11 | 120.6 (2) | H17D—C17B—H17F | 109.5 |
| C9—C10—H10 | 119.7 | H17E—C17B—H17F | 109.5 |
| C11—C10—H10 | 119.7 | | |
| O2—S—C1—C8 | 147.2 (2) | C2—C1—C8—O1 | -0.2 (2) |
| C16A—S—C1—C8 | -79.1 (3) | S—C1—C8—O1 | -178.82 (15) |
| C16B—S—C1—C8 | -114.1 (2) | C2—C1—C8—C9 | -179.1 (2) |
| O2—S—C1—C2 | -31.1 (2) | S—C1—C8—C9 | 2.3 (4) |
| C16A—S—C1—C2 | 102.6 (2) | C7—O1—C8—C1 | 0.6 (2) |
| C16B—S—C1—C2 | 67.5 (2) | C7—O1—C8—C9 | 179.74 (16) |
| C4—C3—C2—C7 | 0.8 (3) | C1—C8—C9—C10 | -1.6 (4) |
| C4—C3—C2—C1 | -179.5 (2) | O1—C8—C9—C10 | 179.5 (2) |
| C8—C1—C2—C7 | -0.2 (2) | C1—C8—C9—C14 | 179.6 (3) |
| S—C1—C2—C7 | 178.46 (15) | O1—C8—C9—C14 | 0.7 (3) |
| C8—C1—C2—C3 | -179.9 (2) | C14—C9—C10—C11 | -1.3 (4) |
| S—C1—C2—C3 | -1.2 (4) | C8—C9—C10—C11 | 179.9 (2) |
| C2—C3—C4—C5 | -1.2 (3) | C9—C10—C11—C12 | -1.4 (4) |
| C2—C3—C4—Cl | 180.00 (15) | C10—C11—C12—C13 | 2.9 (4) |
| C3—C4—C5—C6 | 0.3 (3) | C10—C11—C12—I | -178.48 (19) |
| Cl—C4—C5—C6 | 179.08 (17) | O2 ⁱ —I—C12—C11 | -167.99 (19) |
| C4—C5—C6—C7 | 1.0 (3) | O2 ⁱ —I—C12—C13 | 10.6 (4) |
| C4—C5—C6—C15 | -178.8 (2) | C11—C12—C13—C14 | -1.6 (4) |
| C8—O1—C7—C2 | -0.7 (2) | I—C12—C13—C14 | 179.7 (2) |
| C8—O1—C7—C6 | 179.17 (19) | C12—C13—C14—C9 | -1.1 (4) |
| C3—C2—C7—O1 | -179.68 (18) | C10—C9—C14—C13 | 2.5 (4) |
| C1—C2—C7—O1 | 0.6 (2) | C8—C9—C14—C13 | -178.6 (3) |
| C3—C2—C7—C6 | 0.4 (3) | O2—S—C16A—C17A | 48.1 (6) |
| C1—C2—C7—C6 | -179.3 (2) | C1—S—C16A—C17A | -72.9 (5) |
| C5—C6—C7—O1 | 178.80 (18) | C16B—S—C16A—C17A | 15.4 (5) |
| C15—C6—C7—O1 | -1.5 (3) | O2—S—C16B—C17B | -177.3 (4) |
| C5—C6—C7—C2 | -1.3 (3) | C1—S—C16B—C17B | 73.5 (4) |
| C15—C6—C7—C2 | 178.4 (2) | C16A—S—C16B—C17B | -23.2 (4) |

Symmetry codes: (i) $x+1, y+1, z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

| D—H···A | D—H | H···A | D···A |
|---------|-----|-------|-------|
|---------|-----|-------|-------|


C16A—H16B···O2ⁱⁱ

0.97

2.40

3.297 (6)

154

Symmetry codes: (ii) $-x, -y, -z$.**Fig. 1**

supplementary materials

Fig. 2

