

2-Hydroxyethanaminium 2,4-dinitrophenolate hemihydrate

Yong Yan and Guoxia Han*

School of Chemistry and Chemical Engineering, Nanjing University, Nanjing 210093, People's Republic of China
Correspondence e-mail: gfwref@sina.cn


Received 2 August 2010; accepted 9 August 2010

Key indicators: single-crystal X-ray study; $T = 153\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.024; wR factor = 0.066; data-to-parameter ratio = 7.5.

In the title salt, $\text{C}_2\text{H}_8\text{NO}^+\cdot\text{C}_6\text{H}_3\text{N}_2\text{O}_5^-\cdot0.5\text{H}_2\text{O}$, the anions, cations and water molecules are linked via $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a three-dimensional network.

Related literature

For comparable structures, see: Goddard *et al.* (2002); Iwasaki & Kawano (1977); Kunnert *et al.* (1995); Rais & Bergman (2004); Sieler *et al.* (1994); Yuan *et al.* (2005).



Experimental

Crystal data


 $M_r = 254.20$

 Orthorhombic, $Pca2_1$
 $a = 24.5688(4)\text{ \AA}$
 $b = 10.5945(2)\text{ \AA}$
 $c = 8.4238(2)\text{ \AA}$
 $V = 2192.67(8)\text{ \AA}^3$
 $Z = 8$

 Mo $K\alpha$ radiation

 $\mu = 0.14\text{ mm}^{-1}$
 $T = 153\text{ K}$
 $0.56 \times 0.45 \times 0.33\text{ mm}$

Data collection

Rigaku R-Axis RAPID diffractometer

20722 measured reflections

2686 independent reflections

 2646 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.018$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.024$
 $wR(F^2) = 0.066$
 $S = 1.00$

2686 reflections

357 parameters

1 restraint

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.25\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.17\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O11—H11O \cdots O1	0.84 (3)	1.76 (3)	2.5914 (14)	168 (3)
O12—H12O \cdots O6	0.77 (3)	1.94 (3)	2.6963 (16)	166 (3)
O13—H0A \cdots O1	0.84 (2)	1.94 (2)	2.7315 (13)	156 (2)
O13—H0A \cdots O5	0.84 (2)	2.31 (2)	2.8938 (15)	127 (2)
O13—H0B \cdots O12 ⁱ	0.85 (2)	1.97 (2)	2.8214 (15)	171 (2)
N5—H5B \cdots O9 ⁱⁱ	0.92 (2)	2.13 (2)	2.9620 (15)	150 (2)
N5—H5C \cdots O11 ⁱⁱⁱ	0.92 (2)	1.85 (2)	2.7620 (16)	171 (2)
N5—H5A \cdots O13	0.82 (2)	2.11 (2)	2.9038 (14)	161 (2)
N6—H6C \cdots O5 ⁱⁱ	0.91 (2)	2.07 (2)	2.9127 (16)	155 (2)
N6—H6B \cdots O6 ⁱⁱ	0.87 (3)	1.93 (3)	2.7453 (17)	155 (2)
N6—H6B \cdots O10 ⁱⁱ	0.87 (3)	2.34 (2)	2.9517 (16)	127 (2)
N6—H6A \cdots O13	0.93 (2)	1.87 (2)	2.7937 (17)	175 (2)

Symmetry codes: (i) $-x + 1, -y + 1, z + \frac{1}{2}$; (ii) $-x + 1, -y + 1, z - \frac{1}{2}$; (iii) $-x + 1, -y, z + \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku/MSC, 2004); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5319).

References

- Goddard, R., Herzog, H. M. & Reetz, M. T. (2002). *Tetrahedron*, **58**, 7847–7853.
- Iwasaki, F. & Kawano, Y. (1977). *Acta Cryst. B* **33**, 2455–2459.
- Kunnert, M., Zahn, G. & Sieler, J. (1995). *Z. Anorg. Allg. Chem.* **621**, 1597–1582.
- Rais, D. & Bergman, R. G. (2004). *Chem. Eur. J.* **10**, 3970–3974.
- Rigaku/MSC (2004). *RAPID-AUTO*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sieler, J., Pink, M. & Zahn, G. (1994). *Z. Anorg. Allg. Chem.* **620**, 743–746.
- Yuan, J.-X., Shen, Z.-L., Li, L., Song, X.-Y. & Jin, Z.-M. (2005). *Acta Cryst. E* **61**, o3712–o3714.

supplementary materials

Acta Cryst. (2010). E66, o2342 [doi:10.1107/S1600536810031983]

2-Hydroxyethanaminium 2,4-dinitrophenolate hemihydrate

Y. Yan and G. Han

Comment

As shown in Fig. 1, compound (I) consists of two crystallographically independent ionic $O^+—H\cdots O^-$ hydrogen bonding (Table 2) pairs of ethanolammonium DNPL (A and B for containing O1 and O6, respectively), and one water molecule. The ion pairs of A and B are associated *via* the water molecule by a $N—H\cdots O$ hydrogen bonds (Table 2).

Following analysis shows the feature of quinoic-phenolic resonance of DNPL in ionic pairs A. The C1—C6 and C1—C2 bond lengths are extremely longer than the normal length (1.38 Å), and the C3—C4 bond length is obviously longer than the normal length. The C2—C3 bond length is the shortest one in DNPL, and is slightly shorter than the normal one (Table 1). The N1—C6 [1.4416 (16) Å] and N2—C4 [1.4478 (17) Å] bond lengths are obviously shorter than that observed in 2,4-dinitrophenol [1.484 (5) Å, Iwasaki & Kawano, 1977]. The C1—O1 bond length [1.2671 (16) Å] is shorter than that of phenolates, 1.317 (4) Å for a naphthalate (Yuan *et al.*, 2005), 1.283 (2)—1.331 (6) Å for phenolates in different compounds (Sieler *et al.*, 1994; Kunnert *et al.*, 1995; Goddard *et al.*, 2002; Rais *et al.*, 2004).

The nitro groups of the DNPL at C2 and C4 are twisted relatively to the aromatic ring with dihedral angles lesser than about 4°, which are indicated by torsion angles in Table 2. This twist of the nitro group relative to the benzene ring is a result of the participation of nitro O atoms in $O—H\cdots O$ hydrogen bonds (Table 2).

The $N—H\cdots O$ and $O—H\cdots O$ hydrogen bonding in the structure leads to a three-dimensional hydrogen-bonded network (Table 2). As shown in Fig. 2, the water molecule acts as well a hydrogen-bond donor for DNPL anion and ethanolammonium cation. Furthermore, the DNPL anions are linked to ethanolammonium cations *via* various hydrogen bonds of $N—H\cdots O$ (Table 2 & Fig. 2).

Experimental

Admixture of ethanolamine, DNP and water in the molar ratio of 2:1:1 were heated to a temperature where clear solutions resulted. The crystals of (I) were formed by making the resulting solutions standing overnight at 293K.

Refinement

The H atoms bonded to N and O were taken from a difference fourier map, and were refined. Others were placed in calculated positions and allowed to ride on their parent atoms at distances of 0.95 (phenyl), 0.99 (methylene), with $U_{iso}(H)$ values 1.2 times U_{eq} of the parent atoms.

supplementary materials

Figures




Fig. 1. The cell unit of (I) with atom labels, showing 40% probability displacement ellipsoids. Hydrogen bonds are illustrated as thin lines.




Fig. 2. A diagram of crystal packing viewed down along the b axis. Hydrogen bonds are drawn as dashed lines.

2-Hydroxyethanaminium 2,4-dinitrophenolate hemihydrate

Crystal data

$\text{C}_2\text{H}_8\text{NO}^+\cdot\text{C}_6\text{H}_3\text{N}_2\text{O}_5^- \cdot 0.5\text{H}_2\text{O}$	$F(000) = 1064$
$M_r = 254.20$	$D_x = 1.540 \text{ Mg m}^{-3}$
Orthorhombic, $Pca2_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2c -2ac	Cell parameters from 20146 reflections
$a = 24.5688 (4) \text{ \AA}$	$\theta = 3.1\text{--}27.5^\circ$
$b = 10.5945 (2) \text{ \AA}$	$\mu = 0.14 \text{ mm}^{-1}$
$c = 8.4238 (2) \text{ \AA}$	$T = 153 \text{ K}$
$V = 2192.67 (8) \text{ \AA}^3$	Prism, yellow
$Z = 8$	$0.56 \times 0.45 \times 0.33 \text{ mm}$

Data collection

Rigaku R-AXIS RAPID diffractometer	2646 reflections with $I > 2\sigma(I)$
Radiation source: Rotating Anode graphite	$R_{\text{int}} = 0.018$
ω scans	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.1^\circ$
20722 measured reflections	$h = -31 \rightarrow 31$
2686 independent reflections	$k = -13 \rightarrow 13$
	$l = -10 \rightarrow 10$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.024$	H atoms treated by a mixture of independent and constrained refinement

$wR(F^2) = 0.066$	$w = 1/[\sigma^2(F_o^2) + (0.0506P)^2 + 0.256P]$
	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.00$	$(\Delta/\sigma)_{\max} = 0.001$
2686 reflections	$\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$
357 parameters	$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$
1 restraint	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0213 (13)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.40970 (4)	0.16257 (9)	0.70708 (14)	0.0246 (2)
O2	0.15648 (4)	0.16118 (13)	0.7433 (2)	0.0436 (4)
O3	0.17398 (5)	0.31833 (12)	0.89817 (19)	0.0390 (3)
O4	0.35318 (5)	0.43069 (14)	1.0234 (2)	0.0489 (4)
O5	0.42268 (4)	0.35022 (10)	0.91162 (16)	0.0300 (3)
O6	0.38672 (4)	0.63686 (10)	0.71654 (14)	0.0262 (2)
O7	0.13452 (4)	0.69048 (12)	0.70476 (19)	0.0384 (3)
O8	0.15347 (5)	0.83799 (13)	0.87468 (18)	0.0407 (3)
O9	0.33436 (4)	0.93189 (10)	1.00421 (14)	0.0292 (2)
O10	0.40336 (4)	0.83871 (10)	0.89816 (14)	0.0265 (2)
O11	0.46306 (4)	0.03331 (10)	0.49735 (13)	0.0236 (2)
O12	0.45349 (4)	0.58299 (10)	0.47171 (13)	0.0220 (2)
N1	0.37306 (4)	0.35618 (10)	0.92905 (16)	0.0211 (2)
N2	0.18870 (5)	0.23265 (12)	0.81025 (19)	0.0276 (3)
N3	0.35365 (5)	0.85094 (10)	0.91370 (14)	0.0194 (2)
N4	0.16743 (5)	0.75211 (13)	0.78413 (18)	0.0270 (3)
N5	0.56485 (5)	0.01705 (10)	0.68611 (15)	0.0185 (2)
N6	0.53325 (5)	0.39558 (11)	0.44028 (16)	0.0218 (2)
C1	0.35954 (5)	0.17880 (12)	0.73496 (16)	0.0175 (2)
C2	0.31902 (6)	0.09935 (12)	0.66134 (18)	0.0224 (3)
H2	0.3308	0.0330	0.5938	0.027*
C3	0.26441 (6)	0.11540 (13)	0.68459 (19)	0.0236 (3)
H3	0.2390	0.0615	0.6335	0.028*

supplementary materials

C4	0.24637 (6)	0.21292 (11)	0.78519 (17)	0.0210 (3)
C5	0.28234 (5)	0.29052 (12)	0.86297 (18)	0.0202 (3)
H5	0.2694	0.3552	0.9314	0.024*
C6	0.33790 (5)	0.27319 (12)	0.84031 (16)	0.0176 (3)
C7	0.33707 (5)	0.66684 (12)	0.73307 (15)	0.0186 (3)
C8	0.29498 (6)	0.59359 (13)	0.65700 (19)	0.0243 (3)
H8	0.3054	0.5241	0.5926	0.029*
C9	0.24074 (6)	0.61971 (12)	0.67346 (19)	0.0249 (3)
H9	0.2143	0.5687	0.6220	0.030*
C10	0.22461 (5)	0.72307 (13)	0.76756 (18)	0.0217 (3)
C11	0.26181 (5)	0.79730 (12)	0.84427 (16)	0.0194 (3)
H11	0.2502	0.8663	0.9078	0.023*
C12	0.31712 (5)	0.77025 (12)	0.82799 (16)	0.0180 (3)
C13	0.54204 (6)	-0.08272 (12)	0.58103 (18)	0.0209 (3)
H13A	0.5540	-0.1666	0.6191	0.025*
H13B	0.5561	-0.0711	0.4718	0.025*
C14	0.48060 (6)	-0.07767 (13)	0.57902 (19)	0.0233 (3)
H14A	0.4660	-0.1535	0.5251	0.028*
H14B	0.4666	-0.0767	0.6892	0.028*
C15	0.48229 (6)	0.39280 (14)	0.34614 (19)	0.0273 (3)
H15A	0.4720	0.3040	0.3251	0.033*
H15B	0.4886	0.4346	0.2427	0.033*
C16	0.43622 (5)	0.45829 (13)	0.4314 (2)	0.0246 (3)
H16A	0.4037	0.4621	0.3621	0.029*
H16B	0.4265	0.4110	0.5288	0.029*
O13	0.51234 (4)	0.26120 (9)	0.71903 (13)	0.0197 (2)
H0A	0.4788 (9)	0.249 (2)	0.732 (3)	0.036 (5)*
H0B	0.5255 (8)	0.3027 (19)	0.797 (3)	0.029 (5)*
H5A	0.5523 (9)	0.088 (2)	0.673 (3)	0.037 (6)*
H5B	0.6013 (9)	0.0271 (17)	0.667 (3)	0.031 (5)*
H5C	0.5571 (8)	-0.0081 (19)	0.788 (3)	0.028 (5)*
H6A	0.5286 (8)	0.3495 (18)	0.533 (3)	0.024 (4)*
H6B	0.5606 (10)	0.366 (2)	0.386 (3)	0.041 (6)*
H6C	0.5420 (8)	0.4771 (19)	0.461 (3)	0.028 (5)*
H11O	0.4426 (10)	0.078 (2)	0.555 (4)	0.042 (6)*
H12O	0.4350 (10)	0.610 (2)	0.537 (4)	0.044 (7)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0164 (4)	0.0285 (5)	0.0289 (5)	0.0015 (3)	0.0016 (4)	-0.0118 (4)
O2	0.0166 (5)	0.0491 (7)	0.0652 (10)	-0.0100 (5)	-0.0046 (6)	-0.0124 (7)
O3	0.0200 (5)	0.0422 (6)	0.0548 (8)	0.0079 (4)	0.0030 (5)	-0.0084 (6)
O4	0.0238 (5)	0.0577 (8)	0.0653 (9)	0.0005 (5)	0.0029 (6)	-0.0452 (8)
O5	0.0154 (4)	0.0274 (5)	0.0472 (7)	-0.0029 (4)	0.0007 (5)	-0.0153 (5)
O6	0.0180 (4)	0.0364 (5)	0.0242 (5)	0.0073 (4)	-0.0034 (4)	-0.0102 (5)
O7	0.0173 (5)	0.0440 (6)	0.0539 (8)	-0.0048 (4)	-0.0075 (5)	-0.0008 (7)
O8	0.0223 (5)	0.0569 (7)	0.0429 (7)	0.0128 (5)	0.0000 (5)	-0.0094 (7)

O9	0.0240 (5)	0.0303 (5)	0.0333 (6)	0.0067 (4)	-0.0053 (5)	-0.0156 (5)
O10	0.0158 (4)	0.0313 (5)	0.0326 (6)	-0.0033 (4)	0.0014 (4)	-0.0080 (5)
O11	0.0221 (4)	0.0288 (5)	0.0199 (5)	0.0050 (4)	0.0004 (4)	-0.0066 (4)
O12	0.0188 (4)	0.0221 (5)	0.0250 (5)	-0.0002 (4)	0.0048 (4)	-0.0013 (4)
N1	0.0164 (5)	0.0197 (5)	0.0274 (6)	0.0004 (4)	0.0000 (5)	-0.0073 (5)
N2	0.0151 (5)	0.0309 (6)	0.0368 (7)	-0.0006 (4)	-0.0012 (5)	-0.0004 (6)
N3	0.0184 (5)	0.0197 (5)	0.0199 (6)	0.0016 (4)	-0.0014 (5)	-0.0019 (4)
N4	0.0160 (5)	0.0330 (6)	0.0319 (7)	0.0020 (5)	-0.0023 (5)	0.0046 (5)
N5	0.0168 (5)	0.0181 (5)	0.0206 (6)	0.0016 (4)	0.0013 (4)	0.0027 (4)
N6	0.0210 (5)	0.0191 (5)	0.0253 (6)	0.0023 (4)	0.0065 (5)	0.0015 (5)
C1	0.0167 (5)	0.0178 (5)	0.0178 (6)	0.0003 (4)	0.0000 (5)	-0.0013 (5)
C2	0.0221 (6)	0.0198 (6)	0.0253 (7)	-0.0014 (5)	-0.0010 (5)	-0.0060 (5)
C3	0.0206 (6)	0.0226 (6)	0.0275 (7)	-0.0052 (5)	-0.0036 (6)	-0.0027 (6)
C4	0.0129 (5)	0.0231 (5)	0.0269 (7)	-0.0009 (5)	0.0001 (5)	0.0014 (5)
C5	0.0159 (6)	0.0204 (5)	0.0242 (7)	0.0016 (4)	0.0007 (5)	-0.0024 (5)
C6	0.0155 (5)	0.0173 (5)	0.0201 (6)	-0.0015 (4)	0.0000 (5)	-0.0028 (5)
C7	0.0182 (6)	0.0218 (6)	0.0160 (6)	0.0028 (5)	-0.0025 (5)	-0.0009 (5)
C8	0.0242 (7)	0.0217 (6)	0.0269 (7)	0.0024 (5)	-0.0064 (6)	-0.0065 (5)
C9	0.0223 (7)	0.0219 (5)	0.0304 (8)	-0.0016 (5)	-0.0080 (6)	-0.0019 (6)
C10	0.0155 (6)	0.0250 (6)	0.0245 (7)	0.0014 (5)	-0.0021 (5)	0.0032 (5)
C11	0.0167 (6)	0.0221 (5)	0.0195 (6)	0.0037 (4)	-0.0009 (5)	0.0001 (5)
C12	0.0162 (6)	0.0197 (5)	0.0179 (6)	0.0008 (4)	-0.0018 (5)	-0.0012 (5)
C13	0.0242 (6)	0.0182 (5)	0.0203 (6)	0.0026 (5)	0.0045 (5)	-0.0009 (5)
C14	0.0230 (6)	0.0238 (6)	0.0231 (7)	-0.0047 (5)	0.0040 (5)	-0.0030 (6)
C15	0.0287 (7)	0.0279 (6)	0.0253 (7)	0.0012 (5)	0.0016 (6)	-0.0055 (6)
C16	0.0187 (6)	0.0240 (6)	0.0310 (8)	-0.0036 (5)	0.0015 (5)	-0.0012 (6)
O13	0.0156 (4)	0.0214 (4)	0.0220 (5)	-0.0015 (3)	0.0009 (4)	-0.0011 (4)

Geometric parameters (Å, °)

O1—C1	1.2661 (16)	C2—C3	1.3665 (19)
O2—N2	1.2322 (18)	C2—H2	0.9500
O3—N2	1.2261 (19)	C3—C4	1.408 (2)
O4—N1	1.2223 (17)	C3—H3	0.9500
O5—N1	1.2293 (15)	C4—C5	1.3734 (19)
O6—C7	1.2683 (16)	C5—C6	1.3905 (17)
O7—N4	1.2357 (19)	C5—H5	0.9500
O8—N4	1.236 (2)	C7—C12	1.4421 (18)
O9—N3	1.2417 (15)	C7—C8	1.4429 (18)
O10—N3	1.2351 (15)	C8—C9	1.3680 (19)
O11—C14	1.4288 (17)	C8—H8	0.9500
O11—H11O	0.84 (3)	C9—C10	1.409 (2)
O12—C16	1.4285 (17)	C9—H9	0.9500
O12—H12O	0.77 (3)	C10—C11	1.3681 (18)
N1—C6	1.4416 (16)	C11—C12	1.3955 (17)
N2—C4	1.4478 (17)	C11—H11	0.9500
N3—C12	1.4345 (17)	C13—C14	1.5105 (19)
N4—C10	1.4449 (16)	C13—H13A	0.9900
N5—C13	1.4882 (18)	C13—H13B	0.9900

supplementary materials

N5—H5A	0.82 (2)	C14—H14A	0.9900
N5—H5B	0.92 (2)	C14—H14B	0.9900
N5—H5C	0.92 (2)	C15—C16	1.510 (2)
N6—C15	1.4824 (19)	C15—H15A	0.9900
N6—H6A	0.93 (2)	C15—H15B	0.9900
N6—H6B	0.87 (3)	C16—H16A	0.9900
N6—H6C	0.91 (2)	C16—H16B	0.9900
C1—C6	1.4388 (18)	O13—H0A	0.84 (2)
C1—C2	1.4438 (18)	O13—H0B	0.85 (2)
C14—O11—H11O	111.2 (18)	O6—C7—C12	125.31 (12)
C16—O12—H12O	109.9 (18)	O6—C7—C8	120.39 (12)
O4—N1—O5	120.48 (12)	C12—C7—C8	114.28 (11)
O4—N1—C6	119.45 (11)	C9—C8—C7	122.97 (13)
O5—N1—C6	120.06 (11)	C9—C8—H8	118.5
O3—N2—O2	122.82 (13)	C7—C8—H8	118.5
O3—N2—C4	118.92 (12)	C8—C9—C10	119.23 (12)
O2—N2—C4	118.25 (13)	C8—C9—H9	120.4
O10—N3—O9	121.00 (11)	C10—C9—H9	120.4
O10—N3—C12	120.19 (11)	C11—C10—C9	121.64 (12)
O9—N3—C12	118.81 (11)	C11—C10—N4	118.79 (12)
O7—N4—O8	122.78 (13)	C9—C10—N4	119.57 (13)
O7—N4—C10	118.13 (13)	C10—C11—C12	119.09 (12)
O8—N4—C10	119.08 (13)	C10—C11—H11	120.5
C13—N5—H5A	115.5 (17)	C12—C11—H11	120.5
C13—N5—H5B	110.1 (13)	C11—C12—N3	115.94 (12)
H5A—N5—H5B	103.6 (19)	C11—C12—C7	122.78 (12)
C13—N5—H5C	105.7 (13)	N3—C12—C7	121.28 (11)
H5A—N5—H5C	108 (2)	N5—C13—C14	110.97 (11)
H5B—N5—H5C	113.8 (18)	N5—C13—H13A	109.4
C15—N6—H6A	109.6 (12)	C14—C13—H13A	109.4
C15—N6—H6B	111.3 (16)	N5—C13—H13B	109.4
H6A—N6—H6B	110 (2)	C14—C13—H13B	109.4
C15—N6—H6C	108.8 (13)	H13A—C13—H13B	108.0
H6A—N6—H6C	111.6 (19)	O11—C14—C13	109.63 (11)
H6B—N6—H6C	105 (2)	O11—C14—H14A	109.7
O1—C1—C6	124.65 (12)	C13—C14—H14A	109.7
O1—C1—C2	120.82 (12)	O11—C14—H14B	109.7
C6—C1—C2	114.53 (11)	C13—C14—H14B	109.7
C3—C2—C1	122.89 (12)	H14A—C14—H14B	108.2
C3—C2—H2	118.6	N6—C15—C16	111.68 (13)
C1—C2—H2	118.6	N6—C15—H15A	109.3
C2—C3—C4	119.12 (12)	C16—C15—H15A	109.3
C2—C3—H3	120.4	N6—C15—H15B	109.3
C4—C3—H3	120.4	C16—C15—H15B	109.3
C5—C4—C3	121.60 (12)	H15A—C15—H15B	107.9
C5—C4—N2	118.28 (12)	O12—C16—C15	108.38 (11)
C3—C4—N2	120.12 (12)	O12—C16—H16A	110.0
C4—C5—C6	119.15 (12)	C15—C16—H16A	110.0
C4—C5—H5	120.4	O12—C16—H16B	110.0

C6—C5—H5	120.4	C15—C16—H16B	110.0
C5—C6—C1	122.63 (12)	H16A—C16—H16B	108.4
C5—C6—N1	115.89 (11)	H0A—O13—H0B	111 (2)
C1—C6—N1	121.48 (11)		
O1—C1—C2—C3	−178.24 (14)	C12—C7—C8—C9	−0.1 (2)
C6—C1—C2—C3	2.5 (2)	C7—C8—C9—C10	0.5 (2)
C1—C2—C3—C4	−0.4 (2)	C8—C9—C10—C11	−0.7 (2)
C2—C3—C4—C5	−1.4 (2)	C8—C9—C10—N4	179.33 (14)
C2—C3—C4—N2	179.31 (13)	O7—N4—C10—C11	174.57 (14)
O3—N2—C4—C5	0.8 (2)	O8—N4—C10—C11	−4.5 (2)
O2—N2—C4—C5	−178.37 (15)	O7—N4—C10—C9	−5.4 (2)
O3—N2—C4—C3	−179.87 (15)	O8—N4—C10—C9	175.54 (14)
O2—N2—C4—C3	1.0 (2)	C9—C10—C11—C12	0.4 (2)
C3—C4—C5—C6	0.7 (2)	N4—C10—C11—C12	−179.61 (12)
N2—C4—C5—C6	−179.92 (13)	C10—C11—C12—N3	−179.11 (12)
C4—C5—C6—C1	1.6 (2)	C10—C11—C12—C7	0.1 (2)
C4—C5—C6—N1	−178.49 (12)	O10—N3—C12—C11	−175.46 (12)
O1—C1—C6—C5	177.64 (13)	O9—N3—C12—C11	5.19 (18)
C2—C1—C6—C5	−3.1 (2)	O10—N3—C12—C7	5.35 (19)
O1—C1—C6—N1	−2.2 (2)	O9—N3—C12—C7	−174.00 (13)
C2—C1—C6—N1	177.00 (12)	O6—C7—C12—C11	−178.70 (13)
O4—N1—C6—C5	3.9 (2)	C8—C7—C12—C11	−0.2 (2)
O5—N1—C6—C5	−177.39 (13)	O6—C7—C12—N3	0.4 (2)
O4—N1—C6—C1	−176.23 (15)	C8—C7—C12—N3	178.91 (12)
O5—N1—C6—C1	2.5 (2)	N5—C13—C14—O11	69.73 (14)
O6—C7—C8—C9	178.49 (14)	N6—C15—C16—O12	54.43 (16)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O11—H11O···O1	0.84 (3)	1.76 (3)	2.5914 (14)	168 (3)
O12—H12O···O6	0.77 (3)	1.94 (3)	2.6963 (16)	166 (3)
O13—H0A···O1	0.84 (2)	1.94 (2)	2.7315 (13)	156 (2)
O13—H0A···O5	0.84 (2)	2.31 (2)	2.8938 (15)	127 (2)
O13—H0B···O12 ⁱ	0.85 (2)	1.97 (2)	2.8214 (15)	171 (2)
N5—H5B···O9 ⁱⁱ	0.92 (2)	2.13 (2)	2.9620 (15)	150 (2)
N5—H5C···O11 ⁱⁱⁱ	0.92 (2)	1.85 (2)	2.7620 (16)	171 (2)
N5—H5A···O13	0.82 (2)	2.11 (2)	2.9038 (14)	161 (2)
N6—H6C···O5 ⁱⁱ	0.91 (2)	2.07 (2)	2.9127 (16)	155 (2)
N6—H6B···O6 ⁱⁱ	0.87 (3)	1.93 (3)	2.7453 (17)	155 (2)
N6—H6B···O10 ⁱⁱ	0.87 (3)	2.34 (2)	2.9517 (16)	127 (2)
N6—H6A···O13	0.93 (2)	1.87 (2)	2.7937 (17)	175 (2)

Symmetry codes: (i) $-x+1, -y+1, z+1/2$; (ii) $-x+1, -y+1, z-1/2$; (iii) $-x+1, -y, z+1/2$.

supplementary materials

Fig. 1




Fig. 2

