

4-Chloro-N-[*(E*)-(3,4-dimethoxyphenyl)-methylidene]aniline

M. Nawaz Tahir,^{a*} Muhammad Ilyas Tariq,^b Muhammad Sarfraz,^b Shahbaz Ahmad^b and Riaz H. Tariq^c

^aDepartment of Physics, University of Sargodha, Sargodha, Pakistan, ^bDepartment of Chemistry, University of Sargodha, Sargodha, Pakistan, and ^cInstitute of Chemical and Pharmaceutical Sciences, The University of Faisalabad, Faisalabad, Pakistan
Correspondence e-mail: dmntahir_uos@yahoo.com


Received 14 August 2010; accepted 15 August 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.040; wR factor = 0.120; data-to-parameter ratio = 14.4.

The asymmetric unit of the title compound, $\text{C}_{15}\text{H}_{14}\text{ClNO}_2$, contains two molecules with significantly different conformations: the dihedral angles between the 4-chloroaniline and 3,4-dimethoxyphenyl (excluding C atoms) moieties are $19.68(7)$ and $45.54(4)^\circ$. In the crystal, the molecules are linked by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds and weak $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For related structures, see: Dehno Khalaji *et al.* (2009); Shang & Tan (2007).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{14}\text{ClNO}_2$
 $M_r = 275.72$
Monoclinic, $P2_1/c$
 $a = 12.4227(4)\text{ \AA}$
 $b = 7.3638(2)\text{ \AA}$
 $c = 30.4583(13)\text{ \AA}$
 $\beta = 96.080(2)^\circ$
 $V = 2770.60(17)\text{ \AA}^3$
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.27\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.35 \times 0.22 \times 0.20\text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.932$, $T_{\max} = 0.950$
20975 measured reflections
5007 independent reflections
3456 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.120$
 $S = 1.05$
5007 reflections
347 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.19\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.30\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$, $Cg2$ and $Cg3$ are the centroids of the C1–C6, C8–C13 and C16–C21 rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C14–H14B···O3 ⁱ	0.96	2.52	3.468 (3)	170
C6–H6···Cg3 ⁱⁱ	0.93	2.85	3.602 (2)	139
C18–H18···Cg1	0.93	2.89	3.588 (2)	133
C21–H21···Cg3 ⁱⁱⁱ	0.93	2.88	3.549 (2)	130
C29–H29C···Cg2	0.96	2.88	3.782 (2)	157
C30–H30C···Cg1 ^{iv}	0.96	2.76	3.613 (2)	148

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $x, y + 1, z$; (iii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $x + 1, y - 1, z$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the provision of funds for the purchase of diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5610).

References

- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dehno Khalaji, A., Asghari, J., Fejfarová, K. & Dušek, M. (2009). *Acta Cryst. E65*, o253.
- Farrugia, L. J. (1997). *J. Appl. Cryst. 30*, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst. 32*, 837–838.
- Shang, Z.-H. & Tan, S. (2007). *Acta Cryst. E63*, o2960–o2961.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o2355 [doi:10.1107/S1600536810032782]

4-Chloro-N-[*(E*)-(3,4-dimethoxyphenyl)methylidene]aniline

M. N. Tahir, M. I. Tariq, M. Sarfraz, S. Ahmad and R. H. Tariq

Comment

The crystal structures of (II) i.e., 4-chloro-*N*-(3,4,5-trimethoxybenzylidene)aniline (Dehno Khalaji *et al.*, 2009) and (III) i.e., 4-[(4-chlorophenyl)iminomethyl]-2-methoxyphenol (Shang & Tan, 2007) have been published which are related to the title compound (I, Fig. 1)

The title compound consists of two molecules in the crystallographic asymmetric unit which differ from each other geometrically. In one molecule, the 4-chloroanilinic group A (C1—C6/N1/CL1) and group B (C7—C13/O1/O2) of 3,4-dimethoxyphenyl are planar with r.m.s. deviation of 0.0081 and 0.0146 Å, respectively. The dihedral angle between A/B is 45.54 (4)°. The C-atoms C14 and C15 of 3,4-dimethoxyphenyl are at a distance of -0.0058 (35) and 0.1132 (34) Å, respectively from the parent group B. In the second molecule, the 4-chloroanilinic group C (C16—C21/N2/CL2) and group D (C22—C28/O3/O4) of 3,4-dimethoxyphenyl are planar with r.m.s. deviation of 0.0094 and 0.0063 Å, respectively. The dihedral angle between C/D is 19.68 (7)°. The C-atoms C29 and C30 of 3,4-dimethoxyphenyl are at a distance of -0.2256 (33) and -0.2205 (31) Å, respectively from the parent group D. This shows that both molecules differ at large from each other. The molecules are stabilized through C—H···O type of H-bonding and π ··· π interactions (Table 1).

Experimental

Equimolar quantities of 4-chloroaniline and 3,4-dimethoxybenzaldehyde were refluxed in methanol for 30 min. The solution was kept at room temperature which afforded orange light yellow prisms of (I) after 48 h.

Refinement

The H-atoms were positioned geometrically (C—H = 0.93–0.96 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl and $x = 1.2$ for other H-atoms.

Figures




Fig. 1. View of (I) with displacement ellipsoids drawn at the 50% probability level. H-atoms are shown by small circles of arbitrary radii.

supplementary materials

4-Chloro-N-[*(E*)-(3,4-dimethoxyphenyl)methylidene]aniline

Crystal data

C ₁₅ H ₁₄ ClNO ₂	<i>F</i> (000) = 1152
<i>M_r</i> = 275.72	<i>D_x</i> = 1.322 Mg m ⁻³
Monoclinic, <i>P2₁/c</i>	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: -P 2ybc	Cell parameters from 3456 reflections
<i>a</i> = 12.4227 (4) Å	θ = 1.7–25.3°
<i>b</i> = 7.3638 (2) Å	μ = 0.27 mm ⁻¹
<i>c</i> = 30.4583 (13) Å	<i>T</i> = 296 K
β = 96.080 (2)°	Prism, light yellow
<i>V</i> = 2770.60 (17) Å ³	0.35 × 0.22 × 0.20 mm
<i>Z</i> = 8	

Data collection

Bruker Kappa APEXII CCD diffractometer	5007 independent reflections
Radiation source: fine-focus sealed tube graphite	3456 reflections with $I > 2\sigma(I)$
Detector resolution: 8.10 pixels mm ⁻¹	$R_{\text{int}} = 0.026$
ω scans	$\theta_{\text{max}} = 25.3^\circ$, $\theta_{\text{min}} = 1.7^\circ$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	$h = -14 \rightarrow 14$
$T_{\text{min}} = 0.932$, $T_{\text{max}} = 0.950$	$k = -8 \rightarrow 8$
20975 measured reflections	$l = -36 \rightarrow 29$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.040$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.120$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0465P)^2 + 1.0036P]$
5007 reflections	where $P = (F_o^2 + 2F_c^2)/3$
347 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.19 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	-0.47381 (6)	0.59742 (12)	0.19551 (3)	0.1029 (3)
O1	0.34541 (11)	0.7300 (2)	0.00839 (5)	0.0666 (6)
O2	0.15540 (11)	0.7756 (2)	-0.03006 (4)	0.0622 (5)
N1	-0.10335 (14)	0.6680 (2)	0.09001 (5)	0.0531 (6)
C1	-0.18906 (15)	0.6522 (3)	0.11683 (6)	0.0480 (6)
C2	-0.27949 (16)	0.5526 (3)	0.10049 (7)	0.0534 (7)
C3	-0.36655 (17)	0.5352 (3)	0.12478 (7)	0.0582 (8)
C4	-0.36430 (18)	0.6199 (3)	0.16498 (8)	0.0597 (8)
C5	-0.27691 (19)	0.7213 (3)	0.18141 (7)	0.0641 (8)
C6	-0.18948 (17)	0.7378 (3)	0.15726 (7)	0.0572 (8)
C7	-0.00599 (17)	0.6629 (3)	0.10774 (7)	0.0547 (7)
C8	0.08663 (16)	0.6815 (3)	0.08263 (7)	0.0505 (7)
C9	0.07247 (15)	0.7174 (3)	0.03730 (6)	0.0481 (7)
C10	0.15987 (16)	0.7355 (3)	0.01380 (6)	0.0477 (6)
C11	0.26563 (16)	0.7137 (3)	0.03515 (7)	0.0505 (7)
C12	0.27930 (17)	0.6779 (3)	0.07967 (7)	0.0595 (8)
C13	0.19067 (17)	0.6630 (3)	0.10334 (7)	0.0606 (8)
C14	0.45431 (17)	0.7121 (4)	0.02793 (9)	0.0797 (10)
C15	0.05181 (18)	0.8106 (4)	-0.05280 (7)	0.0704 (9)
Cl2	-0.28854 (6)	0.13333 (12)	0.26687 (2)	0.0916 (3)
O3	0.34354 (11)	0.2151 (2)	0.04035 (5)	0.0685 (6)
O4	0.53216 (11)	0.1383 (2)	0.07674 (5)	0.0614 (5)
N2	0.08863 (14)	0.1523 (2)	0.16357 (6)	0.0538 (6)
C16	0.00252 (16)	0.1438 (3)	0.19048 (6)	0.0474 (7)
C17	-0.09141 (16)	0.2363 (3)	0.17535 (7)	0.0526 (7)
C18	-0.18002 (17)	0.2347 (3)	0.19879 (7)	0.0575 (7)
C19	-0.17623 (17)	0.1391 (3)	0.23760 (7)	0.0535 (7)
C20	-0.08492 (18)	0.0463 (3)	0.25324 (7)	0.0614 (8)
C21	0.00371 (17)	0.0481 (3)	0.22977 (7)	0.0572 (7)
C22	0.18473 (17)	0.1189 (3)	0.17922 (7)	0.0597 (8)
C23	0.27594 (16)	0.1226 (3)	0.15321 (7)	0.0547 (7)
C24	0.26207 (16)	0.1687 (3)	0.10854 (7)	0.0511 (7)
C25	0.34811 (16)	0.1725 (3)	0.08394 (7)	0.0502 (7)
C26	0.45269 (15)	0.1286 (3)	0.10403 (7)	0.0515 (7)

supplementary materials

C27	0.46640 (17)	0.0814 (3)	0.14774 (8)	0.0654 (8)
C28	0.37878 (18)	0.0795 (4)	0.17212 (8)	0.0700 (9)
C29	0.23950 (17)	0.2317 (3)	0.01634 (7)	0.0672 (8)
C30	0.63635 (16)	0.0663 (3)	0.09192 (8)	0.0675 (9)
H2	-0.28136	0.49722	0.07297	0.0640*
H3	-0.42638	0.46651	0.11397	0.0699*
H5	-0.27641	0.77864	0.20863	0.0769*
H6	-0.13016	0.80729	0.16828	0.0686*
H7	0.00653	0.64637	0.13813	0.0656*
H9	0.00291	0.72905	0.02296	0.0577*
H12	0.34870	0.66362	0.09400	0.0714*
H13	0.20112	0.64022	0.13353	0.0727*
H14A	0.46754	0.79754	0.05169	0.1193*
H14B	0.50294	0.73565	0.00611	0.1193*
H14C	0.46581	0.59102	0.03914	0.1193*
H15A	0.00931	0.70152	-0.05403	0.1056*
H15B	0.05986	0.85091	-0.08225	0.1056*
H15C	0.01616	0.90321	-0.03750	0.1056*
H17	-0.09447	0.30033	0.14893	0.0631*
H18	-0.24229	0.29817	0.18840	0.0690*
H20	-0.08271	-0.01779	0.27963	0.0737*
H21	0.06554	-0.01593	0.24041	0.0687*
H22	0.19776	0.09042	0.20906	0.0716*
H24	0.19328	0.19723	0.09526	0.0614*
H27	0.53481	0.05060	0.16104	0.0785*
H28	0.38922	0.04869	0.20187	0.0840*
H29A	0.19869	0.12289	0.01982	0.1007*
H29B	0.24729	0.24997	-0.01437	0.1007*
H29C	0.20224	0.33350	0.02736	0.1007*
H30A	0.66641	0.13300	0.11741	0.1012*
H30B	0.68330	0.07689	0.06896	0.1012*
H30C	0.62931	-0.05923	0.09960	0.1012*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0819 (5)	0.1266 (7)	0.1086 (6)	0.0040 (4)	0.0499 (4)	0.0131 (5)
O1	0.0419 (8)	0.0932 (12)	0.0652 (10)	-0.0014 (8)	0.0074 (7)	0.0011 (9)
O2	0.0502 (8)	0.0884 (11)	0.0482 (8)	0.0024 (8)	0.0058 (7)	0.0031 (8)
N1	0.0510 (10)	0.0576 (10)	0.0509 (10)	-0.0004 (8)	0.0067 (8)	0.0022 (8)
C1	0.0490 (11)	0.0469 (11)	0.0480 (11)	0.0033 (9)	0.0048 (9)	0.0062 (9)
C2	0.0576 (13)	0.0507 (12)	0.0512 (12)	0.0006 (10)	0.0028 (10)	0.0015 (10)
C3	0.0496 (12)	0.0575 (13)	0.0664 (14)	-0.0020 (10)	0.0005 (10)	0.0109 (11)
C4	0.0568 (13)	0.0607 (14)	0.0640 (14)	0.0086 (11)	0.0179 (11)	0.0117 (11)
C5	0.0749 (16)	0.0635 (14)	0.0554 (13)	0.0048 (12)	0.0139 (12)	-0.0054 (11)
C6	0.0574 (13)	0.0566 (13)	0.0572 (13)	-0.0035 (10)	0.0045 (10)	-0.0052 (10)
C7	0.0574 (13)	0.0604 (13)	0.0464 (11)	0.0015 (10)	0.0059 (10)	0.0051 (10)
C8	0.0492 (12)	0.0535 (12)	0.0487 (11)	0.0007 (9)	0.0043 (9)	0.0026 (9)

C9	0.0419 (11)	0.0496 (12)	0.0518 (12)	0.0011 (9)	0.0001 (9)	-0.0002 (9)
C10	0.0482 (11)	0.0469 (11)	0.0472 (11)	0.0005 (9)	0.0015 (9)	-0.0022 (9)
C11	0.0450 (11)	0.0514 (12)	0.0548 (12)	-0.0009 (9)	0.0043 (9)	-0.0033 (10)
C12	0.0438 (11)	0.0724 (15)	0.0601 (14)	0.0015 (10)	-0.0049 (10)	-0.0011 (11)
C13	0.0588 (13)	0.0740 (15)	0.0478 (12)	0.0017 (11)	0.0000 (10)	0.0064 (11)
C14	0.0445 (13)	0.113 (2)	0.0814 (17)	-0.0016 (13)	0.0054 (12)	-0.0137 (16)
C15	0.0656 (15)	0.0966 (18)	0.0477 (12)	0.0126 (13)	0.0000 (11)	-0.0030 (12)
Cl2	0.0734 (4)	0.1239 (6)	0.0828 (5)	0.0051 (4)	0.0329 (3)	-0.0075 (4)
O3	0.0441 (8)	0.1016 (12)	0.0596 (9)	0.0077 (8)	0.0045 (7)	0.0180 (9)
O4	0.0406 (8)	0.0748 (10)	0.0686 (10)	0.0043 (7)	0.0045 (7)	0.0094 (8)
N2	0.0529 (10)	0.0587 (11)	0.0503 (10)	0.0024 (8)	0.0072 (8)	0.0034 (8)
C16	0.0516 (12)	0.0447 (11)	0.0452 (11)	0.0001 (9)	0.0023 (9)	-0.0007 (9)
C17	0.0593 (13)	0.0538 (12)	0.0432 (11)	0.0071 (10)	-0.0019 (9)	0.0047 (9)
C18	0.0528 (12)	0.0617 (13)	0.0562 (13)	0.0120 (10)	-0.0030 (10)	-0.0051 (11)
C19	0.0538 (12)	0.0565 (12)	0.0509 (12)	0.0000 (10)	0.0094 (10)	-0.0085 (10)
C20	0.0689 (15)	0.0616 (14)	0.0543 (13)	0.0024 (11)	0.0092 (11)	0.0119 (11)
C21	0.0549 (12)	0.0573 (13)	0.0586 (13)	0.0091 (10)	0.0021 (10)	0.0126 (11)
C22	0.0588 (14)	0.0723 (15)	0.0474 (12)	0.0005 (11)	0.0035 (10)	0.0030 (11)
C23	0.0500 (12)	0.0606 (13)	0.0531 (12)	0.0010 (10)	0.0031 (10)	0.0013 (10)
C24	0.0425 (11)	0.0537 (12)	0.0564 (12)	0.0045 (9)	0.0016 (9)	0.0031 (10)
C25	0.0471 (11)	0.0496 (12)	0.0531 (12)	0.0015 (9)	0.0011 (9)	0.0050 (10)
C26	0.0423 (11)	0.0512 (12)	0.0600 (13)	0.0006 (9)	0.0005 (9)	0.0018 (10)
C27	0.0440 (12)	0.0853 (17)	0.0642 (14)	0.0044 (11)	-0.0073 (10)	0.0105 (12)
C28	0.0580 (14)	0.0963 (18)	0.0538 (13)	0.0041 (12)	-0.0028 (11)	0.0113 (13)
C29	0.0576 (13)	0.0810 (16)	0.0606 (14)	0.0047 (12)	-0.0046 (11)	0.0085 (12)
C30	0.0416 (12)	0.0733 (15)	0.0867 (17)	0.0067 (10)	0.0029 (11)	0.0070 (13)

Geometric parameters (Å, °)

Cl1—C4	1.735 (2)	C13—H13	0.9300
Cl2—C19	1.734 (2)	C14—H14A	0.9600
O1—C11	1.354 (2)	C14—H14B	0.9600
O1—C14	1.425 (3)	C14—H14C	0.9600
O2—C15	1.419 (3)	C15—H15A	0.9600
O2—C10	1.364 (2)	C15—H15B	0.9600
O3—C25	1.360 (3)	C15—H15C	0.9600
O3—C29	1.421 (3)	C16—C17	1.387 (3)
O4—C30	1.430 (3)	C16—C21	1.388 (3)
O4—C26	1.358 (2)	C17—C18	1.374 (3)
N1—C7	1.272 (3)	C18—C19	1.372 (3)
N1—C1	1.414 (2)	C19—C20	1.366 (3)
N2—C22	1.262 (3)	C20—C21	1.375 (3)
N2—C16	1.416 (3)	C22—C23	1.450 (3)
C1—C6	1.384 (3)	C23—C24	1.395 (3)
C1—C2	1.389 (3)	C23—C28	1.382 (3)
C2—C3	1.380 (3)	C24—C25	1.369 (3)
C3—C4	1.372 (3)	C25—C26	1.414 (3)
C4—C5	1.368 (3)	C26—C27	1.369 (3)
C5—C6	1.380 (3)	C27—C28	1.381 (3)

supplementary materials

C7—C8	1.454 (3)	C17—H17	0.9300
C8—C9	1.398 (3)	C18—H18	0.9300
C8—C13	1.384 (3)	C20—H20	0.9300
C9—C10	1.369 (3)	C21—H21	0.9300
C10—C11	1.412 (3)	C22—H22	0.9300
C11—C12	1.374 (3)	C24—H24	0.9300
C12—C13	1.383 (3)	C27—H27	0.9300
C2—H2	0.9300	C28—H28	0.9300
C3—H3	0.9300	C29—H29A	0.9600
C5—H5	0.9300	C29—H29B	0.9600
C6—H6	0.9300	C29—H29C	0.9600
C7—H7	0.9300	C30—H30A	0.9600
C9—H9	0.9300	C30—H30B	0.9600
C12—H12	0.9300	C30—H30C	0.9600
C11—O1—C14	117.62 (17)	H15A—C15—H15B	109.00
C10—O2—C15	117.28 (15)	O2—C15—H15A	109.00
C25—O3—C29	117.64 (16)	O2—C15—H15B	109.00
C26—O4—C30	118.33 (17)	N2—C16—C17	116.51 (17)
C1—N1—C7	119.52 (17)	N2—C16—C21	125.58 (19)
C16—N2—C22	121.01 (18)	C17—C16—C21	117.89 (19)
N1—C1—C6	123.42 (18)	C16—C17—C18	121.0 (2)
N1—C1—C2	117.81 (17)	C17—C18—C19	119.7 (2)
C2—C1—C6	118.70 (18)	C12—C19—C18	119.96 (17)
C1—C2—C3	120.48 (19)	C12—C19—C20	119.48 (17)
C2—C3—C4	119.6 (2)	C18—C19—C20	120.6 (2)
C3—C4—C5	121.0 (2)	C19—C20—C21	119.7 (2)
C11—C4—C3	119.47 (17)	C16—C21—C20	121.2 (2)
C11—C4—C5	119.53 (18)	N2—C22—C23	123.6 (2)
C4—C5—C6	119.5 (2)	C22—C23—C24	120.96 (18)
C1—C6—C5	120.8 (2)	C22—C23—C28	120.6 (2)
N1—C7—C8	122.94 (19)	C24—C23—C28	118.41 (19)
C7—C8—C9	120.89 (18)	C23—C24—C25	121.05 (19)
C7—C8—C13	120.26 (19)	O3—C25—C24	125.80 (19)
C9—C8—C13	118.85 (19)	O3—C25—C26	114.64 (17)
C8—C9—C10	120.72 (18)	C24—C25—C26	119.57 (19)
O2—C10—C11	114.53 (17)	O4—C26—C25	114.67 (18)
O2—C10—C9	125.53 (18)	O4—C26—C27	125.80 (18)
C9—C10—C11	119.93 (17)	C25—C26—C27	119.54 (19)
O1—C11—C10	114.68 (18)	C26—C27—C28	120.1 (2)
O1—C11—C12	126.11 (19)	C23—C28—C27	121.3 (2)
C10—C11—C12	119.21 (19)	C16—C17—H17	119.00
C11—C12—C13	120.5 (2)	C18—C17—H17	119.00
C8—C13—C12	120.8 (2)	C17—C18—H18	120.00
C1—C2—H2	120.00	C19—C18—H18	120.00
C3—C2—H2	120.00	C19—C20—H20	120.00
C2—C3—H3	120.00	C21—C20—H20	120.00
C4—C3—H3	120.00	C16—C21—H21	119.00
C4—C5—H5	120.00	C20—C21—H21	119.00
C6—C5—H5	120.00	N2—C22—H22	118.00

C1—C6—H6	120.00	C23—C22—H22	118.00
C5—C6—H6	120.00	C23—C24—H24	119.00
C8—C7—H7	119.00	C25—C24—H24	119.00
N1—C7—H7	119.00	C26—C27—H27	120.00
C8—C9—H9	120.00	C28—C27—H27	120.00
C10—C9—H9	120.00	C23—C28—H28	119.00
C11—C12—H12	120.00	C27—C28—H28	119.00
C13—C12—H12	120.00	O3—C29—H29A	109.00
C12—C13—H13	120.00	O3—C29—H29B	109.00
C8—C13—H13	120.00	O3—C29—H29C	109.00
O1—C14—H14B	109.00	H29A—C29—H29B	109.00
H14A—C14—H14C	109.00	H29A—C29—H29C	109.00
O1—C14—H14C	109.00	H29B—C29—H29C	109.00
H14A—C14—H14B	109.00	O4—C30—H30A	109.00
O1—C14—H14A	109.00	O4—C30—H30B	109.00
H14B—C14—H14C	109.00	O4—C30—H30C	109.00
O2—C15—H15C	109.00	H30A—C30—H30B	109.00
H15A—C15—H15C	109.00	H30A—C30—H30C	109.00
H15B—C15—H15C	109.00	H30B—C30—H30C	110.00
C14—O1—C11—C10	179.1 (2)	O2—C10—C11—O1	-2.1 (3)
C14—O1—C11—C12	-1.3 (3)	O2—C10—C11—C12	178.25 (19)
C15—O2—C10—C9	3.4 (3)	C9—C10—C11—O1	178.4 (2)
C15—O2—C10—C11	-176.1 (2)	C9—C10—C11—C12	-1.3 (3)
C29—O3—C25—C26	169.99 (18)	O1—C11—C12—C13	-179.5 (2)
C29—O3—C25—C24	-9.7 (3)	C10—C11—C12—C13	0.1 (3)
C30—O4—C26—C25	-169.27 (18)	C11—C12—C13—C8	0.8 (3)
C30—O4—C26—C27	10.8 (3)	N2—C16—C17—C18	179.24 (19)
C1—N1—C7—C8	-178.84 (19)	C21—C16—C17—C18	0.7 (3)
C7—N1—C1—C6	41.7 (3)	N2—C16—C21—C20	-179.1 (2)
C7—N1—C1—C2	-141.6 (2)	C17—C16—C21—C20	-0.6 (3)
C22—N2—C16—C21	-21.7 (3)	C16—C17—C18—C19	-0.5 (3)
C16—N2—C22—C23	179.0 (2)	C17—C18—C19—Cl2	-178.91 (17)
C22—N2—C16—C17	159.9 (2)	C17—C18—C19—C20	0.3 (3)
N1—C1—C6—C5	178.2 (2)	Cl2—C19—C20—C21	178.95 (17)
C6—C1—C2—C3	-1.9 (3)	C18—C19—C20—C21	-0.3 (3)
N1—C1—C2—C3	-178.78 (19)	C19—C20—C21—C16	0.5 (3)
C2—C1—C6—C5	1.5 (3)	N2—C22—C23—C24	1.6 (3)
C1—C2—C3—C4	1.2 (3)	N2—C22—C23—C28	-178.1 (2)
C2—C3—C4—Cl1	-179.88 (17)	C22—C23—C24—C25	179.9 (2)
C2—C3—C4—C5	-0.1 (3)	C28—C23—C24—C25	-0.4 (3)
Cl1—C4—C5—C6	179.46 (17)	C22—C23—C28—C27	179.6 (2)
C3—C4—C5—C6	-0.3 (3)	C24—C23—C28—C27	0.0 (4)
C4—C5—C6—C1	-0.4 (3)	C23—C24—C25—O3	179.9 (2)
N1—C7—C8—C9	3.9 (3)	C23—C24—C25—C26	0.2 (3)
N1—C7—C8—C13	-175.8 (2)	O3—C25—C26—O4	0.8 (3)
C7—C8—C9—C10	179.8 (2)	O3—C25—C26—C27	-179.21 (19)
C9—C8—C13—C12	-0.6 (3)	C24—C25—C26—O4	-179.42 (19)
C13—C8—C9—C10	-0.5 (3)	C24—C25—C26—C27	0.5 (3)
C7—C8—C13—C12	179.1 (2)	O4—C26—C27—C28	179.0 (2)

supplementary materials

C8—C9—C10—C11	1.5 (3)	C25—C26—C27—C28	-1.0 (3)
C8—C9—C10—O2	-178.0 (2)	C26—C27—C28—C23	0.7 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1, Cg2 and Cg3 are the centroids of the C1—C6, C8—C13 and C16—C21 rings, respectively.

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C14—H14B···O3 ⁱ	0.96	2.52	3.468 (3)	170
C6—H6···Cg3 ⁱⁱ	0.93	2.85	3.602 (2)	139
C18—H18···Cg1	0.93	2.89	3.588 (2)	133
C21—H21···Cg3 ⁱⁱⁱ	0.93	2.88	3.549 (2)	130
C29—H29C···Cg2	0.96	2.88	3.782 (2)	157
C30—H30C···Cg1 ^{iv}	0.96	2.76	3.613 (2)	148

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $x, y+1, z$; (iii) $-x, y-1/2, -z+1/2$; (iv) $x+1, y-1, z$.

Fig. 1

