

5-Carboxy-2,4-dihydroxyanilinium chloride

Syeda Sohaila Naz,^a Nazar Ul Islam^a and M. Nawaz Tahir^{b*}

^aInstitute of Chemical Sciences, University of Peshawar, Peshawar, Pakistan, and

^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 15 August 2010; accepted 18 August 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.044; wR factor = 0.098; data-to-parameter ratio = 12.8.

In the title salt, $\text{C}_7\text{H}_8\text{NO}_4^+\cdot\text{Cl}^-$, the organic group is planar with an r.m.s. deviation of 0.0265 \AA . An $S(6)$ ring motif is formed due to an intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond. The compound consists of dimers due to intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds with an $R_2^2(8)$ ring motif. The dimers are interlinked through strong $\text{N}-\text{H}\cdots\text{Cl}$ and $\text{O}-\text{H}\cdots\text{Cl}$ hydrogen bonds, resulting in a three-dimensional polymeric network.

Related literature

For related structures, see: Bendjedou *et al.* (2009); Dobson & Gerkin (1998). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_7\text{H}_8\text{NO}_4^+\cdot\text{Cl}^-$	$V = 912.47(8)\text{ \AA}^3$
$M_r = 205.59$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 5.0667(3)\text{ \AA}$	$\mu = 0.40\text{ mm}^{-1}$
$b = 28.4071(13)\text{ \AA}$	$T = 296\text{ K}$
$c = 6.3966(3)\text{ \AA}$	$0.28 \times 0.18 \times 0.16\text{ mm}$
$\beta = 97.649(3)^\circ$	

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.926$, $T_{\max} = 0.935$

7094 measured reflections
1635 independent reflections
1108 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.052$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.098$
 $S = 1.01$
1635 reflections
128 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.23\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.22\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots O2 ⁱ	0.85 (3)	1.81 (3)	2.653 (3)	177 (4)
N1—H1A \cdots Cl1 ⁱⁱ	0.89	2.24	3.124 (3)	176
N1—H1B \cdots Cl1 ⁱⁱⁱ	0.89	2.29	3.176 (3)	173
N1—H1C \cdots Cl1 ^{iv}	0.89	2.35	3.217 (2)	163
O3—H3 \cdots O2	0.85 (4)	1.84 (3)	2.632 (3)	154 (3)
O4—H4A \cdots Cl1 ^v	0.85 (3)	2.15 (3)	2.985 (2)	170 (3)

Symmetry codes: (i) $-x + 2, -y, -z$; (ii) $x - 1, y, z - 1$; (iii) $x, y, z - 1$; (iv) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$; (v) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the provision of funds for the purchase of diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SI2289).

References

- Bendjedou, L., Cherouana, A., Hadjadj, N., Dahaoui, S. & Lecomte, C. (2009). *Acta Cryst. E65*, o1770–o1771.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dobson, A. J. & Gerkin, R. E. (1998). *Acta Cryst. C54*, 1632–1634.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o2372 [doi:10.1107/S1600536810033337]

5-Carboxy-2,4-dihydroxyanilinium chloride

S. S. Naz, N. U. Islam and M. N. Tahir

Comment

The title compound (I, Fig. 1) has been prepared for derivatization and for the synthesis of metallic complexes.

The crystal structure of (II) *i.e.*, 5-ammoniosalicylic Acid chloride monohydrate (Dobson & Gerkin, 1998) and (III) *i.e.*, bis(3-carboxyanilinium) bis(perchlorate) monohydrate (Bendjeddou *et al.*, 2009) have been published which are related to the title compound.

In (I), the organic group (C1—C7/O1—O4/N1) is planar with r. m. s. deviation of 0.0265 Å. There exist a strong intramolecular H-bond of O—H···O type (Table 1, Fig. 1) completing an S(6) ring motif (Bernstein *et al.*, 1995) in the organic part. The title compound consists of dimers due to intermolecular H-bond of O—H···O type (Table 1, Fig. 2) completing $R_2^2(8)$ ring motif. The dimers are interlinked through strong H-bondings of N—H···Cl and O—H···Cl types (Table 1, Fig. 2) resulting in a three dimensional polymeric network.

Experimental

Concentrated nitric acid (2 mL, 67%) was added drop by drop to β-resorcylic acid (1 g, 97%, 6.3 mmol) in a round bottom flask. The mixture was protected from moisture by CaCl_2 (anhydrous) tube and was allowed to stand for 12 h at room temperature. Then the reaction mixture was diluted with water. The crude material was filtered and recrystallized from water to afford the 5-nitro-β-resorcylic acid.

Then a mixture of 5-nitro-β-resorcylic acid (1.5 g, 7.5 mmol), tin (3 g, 25 mmol), concentrated hydrochloric acid (8 ml, 37%) and absolute ethanol (5 ml) were taken in a 100 ml round bottom flask and heated under reflux with stirring for 40 min. The completion of the reaction was monitored by TLC. The reaction mixture was filtered to remove any unreacted tin. The filtrate was kept for seven days to afford light green prisms of (I).

Refinement

The coordinates of H-atoms of hydroxy groups were refined. The H-atoms were positioned geometrically ($\text{N}—\text{H} = 0.89$, $\text{C}—\text{H} = 0.93$ Å) and refined as riding with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N}, \text{O})$, where $x = 1.2$ for all H-atoms.

supplementary materials

Figures




Fig. 1. View of the title compound with the atom numbering scheme. The displacement ellipsoids are drawn at the 50% probability level. H-atoms are shown as small spheres of arbitrary radii. The dotted line represents the intramolecular H-bonding.




Fig. 2. The partial packing (*PLATON*; Spek, 2009) which shows that molecules form dimers which are interlinked through H-bondings to form a three-dimensional polymeric network.

5-Carboxy-2,4-dihydroxyanilinium chloride

Crystal data



$M_r = 205.59$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 5.0667(3)$ Å

$b = 28.4071(13)$ Å

$c = 6.3966(3)$ Å

$\beta = 97.649(3)^\circ$

$V = 912.47(8)$ Å³

$Z = 4$

$F(000) = 424$

$D_x = 1.497$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1108 reflections

$\theta = 2.9\text{--}25.2^\circ$

$\mu = 0.40$ mm⁻¹

$T = 296$ K

Prism, light green

0.28 × 0.18 × 0.16 mm

Data collection

Bruker Kappa APEXII CCD diffractometer

1635 independent reflections

Radiation source: fine-focus sealed tube graphite

1108 reflections with $I > 2\sigma(I)$

Detector resolution: 8.10 pixels mm⁻¹

$R_{\text{int}} = 0.052$

ω scans

$\theta_{\text{max}} = 25.2^\circ$, $\theta_{\text{min}} = 2.9^\circ$

Absorption correction: multi-scan (*SADABS*; Bruker, 2005)

$h = -5\text{--}6$

$T_{\text{min}} = 0.926$, $T_{\text{max}} = 0.935$

$k = -34\text{--}32$

7094 measured reflections

$l = -7\text{--}7$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.044$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.098$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.01$	$w = 1/[\sigma^2(F_o^2) + (0.036P)^2 + 0.4226P]$ where $P = (F_o^2 + 2F_c^2)/3$
1635 reflections	$(\Delta/\sigma)_{\max} < 0.001$
128 parameters	$\Delta\rho_{\max} = 0.23 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.22 \text{ e \AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.8567 (5)	0.05383 (8)	-0.1244 (4)	0.0527 (8)
O2	0.7828 (4)	0.01248 (8)	0.1583 (3)	0.0553 (8)
O3	0.4286 (5)	0.04378 (8)	0.3905 (4)	0.0620 (10)
O4	0.0113 (4)	0.18833 (7)	0.1726 (3)	0.0498 (8)
N1	0.2950 (5)	0.19898 (8)	-0.1420 (4)	0.0404 (8)
C1	0.7367 (6)	0.04694 (11)	0.0429 (5)	0.0420 (11)
C2	0.5471 (5)	0.08321 (10)	0.0823 (4)	0.0373 (10)
C3	0.4020 (6)	0.08012 (10)	0.2540 (5)	0.0404 (10)
C4	0.2197 (6)	0.11460 (10)	0.2882 (5)	0.0415 (11)
C5	0.1841 (6)	0.15295 (10)	0.1561 (4)	0.0366 (10)
C6	0.3332 (6)	0.15690 (10)	-0.0114 (4)	0.0343 (9)
C7	0.5084 (6)	0.12258 (10)	-0.0495 (4)	0.0376 (10)
Cl1	0.75184 (16)	0.19834 (3)	0.56278 (12)	0.0476 (3)
H1	0.974 (7)	0.0328 (12)	-0.131 (5)	0.0632*
H1A	0.13993	0.19710	-0.22519	0.0485*
H1B	0.42649	0.20130	-0.22080	0.0485*
H1C	0.29515	0.22425	-0.05970	0.0485*
H3	0.543 (7)	0.0264 (13)	0.342 (5)	0.0743*
H4	0.12136	0.11186	0.40041	0.0498*
H4A	-0.063 (6)	0.1874 (11)	0.284 (5)	0.0597*
H7	0.60291	0.12534	-0.16402	0.0451*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0541 (15)	0.0463 (15)	0.0613 (14)	0.0190 (11)	0.0212 (13)	0.0073 (11)
O2	0.0518 (14)	0.0443 (14)	0.0728 (15)	0.0160 (12)	0.0189 (12)	0.0144 (12)
O3	0.0706 (18)	0.0536 (16)	0.0664 (16)	0.0192 (13)	0.0267 (14)	0.0218 (13)
O4	0.0599 (15)	0.0434 (14)	0.0525 (13)	0.0153 (12)	0.0311 (12)	0.0053 (10)
N1	0.0393 (14)	0.0373 (15)	0.0483 (14)	0.0049 (12)	0.0195 (12)	0.0009 (12)
C1	0.0349 (18)	0.0375 (18)	0.0538 (19)	0.0028 (15)	0.0067 (16)	0.0026 (16)
C2	0.0321 (16)	0.0322 (17)	0.0480 (18)	0.0030 (14)	0.0073 (14)	0.0010 (14)
C3	0.0410 (18)	0.0357 (18)	0.0453 (18)	0.0028 (15)	0.0085 (15)	0.0084 (14)
C4	0.0448 (19)	0.0414 (19)	0.0415 (17)	-0.0006 (16)	0.0179 (15)	0.0034 (15)
C5	0.0361 (18)	0.0348 (18)	0.0406 (16)	-0.0005 (14)	0.0116 (14)	-0.0047 (14)
C6	0.0343 (16)	0.0297 (17)	0.0409 (16)	0.0009 (13)	0.0125 (14)	0.0028 (13)
C7	0.0346 (17)	0.0381 (18)	0.0417 (16)	0.0014 (14)	0.0115 (14)	-0.0018 (14)
Cl1	0.0500 (5)	0.0494 (5)	0.0480 (4)	0.0043 (4)	0.0240 (4)	0.0048 (4)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.314 (4)	N1—H1C	0.8900
O2—C1	1.230 (4)	C1—C2	1.453 (4)
O3—C3	1.347 (4)	C2—C7	1.399 (4)
O4—C5	1.346 (4)	C2—C3	1.403 (4)
O1—H1	0.85 (3)	C3—C4	1.384 (4)
O3—H3	0.85 (4)	C4—C5	1.376 (4)
O4—H4A	0.85 (3)	C5—C6	1.395 (4)
N1—C6	1.456 (4)	C6—C7	1.362 (4)
N1—H1A	0.8900	C4—H4	0.9300
N1—H1B	0.8900	C7—H7	0.9300
Cl1…N1 ⁱ	3.176 (3)	C1…C1 ^{vi}	3.580 (4)
Cl1…N1 ⁱⁱ	3.124 (3)	C1…O2 ^{vi}	3.245 (4)
Cl1…C7 ⁱ	3.622 (3)	C1…O2 ^v	3.357 (4)
Cl1…O4 ⁱⁱⁱ	2.985 (2)	C1…C4 ⁱⁱⁱ	3.335 (4)
Cl1…N1 ^{iv}	3.217 (2)	C2…C4 ⁱⁱⁱ	3.598 (4)
Cl1…H4A ⁱⁱⁱ	2.15 (3)	C3…C1 ^{viii}	3.587 (4)
Cl1…H1A ⁱⁱ	2.2400	C4…C2 ^{viii}	3.598 (4)
Cl1…H1B ⁱ	2.2900	C4…C1 ^{viii}	3.335 (4)
Cl1…H1C ^{iv}	2.3500	C7…O4 ⁱⁱⁱ	3.322 (4)
Cl1…H7 ⁱ	2.8800	C7…Cl1 ^{ix}	3.622 (3)
O1…O2 ^v	2.653 (3)	C1…H1 ^v	2.72 (3)
O2…O3	2.632 (3)	C1…H3	2.34 (3)
O2…C1 ^{vi}	3.245 (4)	H1…O2 ^v	1.81 (3)
O2…O1 ^v	2.653 (3)	H1…C1 ^v	2.72 (3)
O2…C1 ^v	3.357 (4)	H1…H1 ^v	2.50 (5)
O3…O3 ^{vii}	2.899 (3)	H1A…Cl1 ^x	2.2400

O3···O2	2.632 (3)	H1A···O4	2.7200
O4···N1	2.642 (3)	H1B···H7	2.3500
O4···Cl1 ^{viii}	2.985 (2)	H1B···Cl1 ^{ix}	2.2900
O4···C7 ^{viii}	3.322 (4)	H1C···O4	2.4300
O1···H7	2.4000	H1C···Cl1 ^{xi}	2.3500
O2···H3	1.84 (3)	H3···C1	2.34 (3)
O2···H1 ^v	1.81 (3)	H3···O2	1.84 (3)
O3···H3 ^{vii}	2.62 (3)	H3···H3 ^{vii}	2.60 (5)
O4···H1C	2.4300	H3···O3 ^{vii}	2.62 (3)
O4···H1A	2.7200	H4···H4A	2.4200
N1···Cl1 ^{ix}	3.176 (3)	H4A···H4	2.4200
N1···Cl1 ^x	3.124 (3)	H4A···Cl1 ^{viii}	2.15 (3)
N1···O4	2.642 (3)	H7···Cl1 ^{ix}	2.8800
N1···Cl1 ^{xi}	3.217 (2)	H7···O1	2.4000
C1···C3 ⁱⁱⁱ	3.587 (4)	H7···H1B	2.3500
C1—O1—H1	110 (2)	O3—C3—C4	116.8 (3)
C3—O3—H3	103 (2)	C2—C3—C4	120.7 (3)
C5—O4—H4A	114 (2)	O3—C3—C2	122.5 (3)
C6—N1—H1C	109.00	C3—C4—C5	120.0 (3)
H1A—N1—H1B	109.00	C4—C5—C6	119.7 (3)
C6—N1—H1A	109.00	O4—C5—C6	115.1 (2)
C6—N1—H1B	109.00	O4—C5—C4	125.2 (3)
H1B—N1—H1C	109.00	N1—C6—C5	117.5 (3)
H1A—N1—H1C	109.00	N1—C6—C7	121.7 (2)
O1—C1—C2	115.1 (3)	C5—C6—C7	120.8 (3)
O1—C1—O2	122.4 (3)	C2—C7—C6	120.5 (3)
O2—C1—C2	122.5 (3)	C3—C4—H4	120.00
C1—C2—C7	120.4 (2)	C5—C4—H4	120.00
C3—C2—C7	118.4 (3)	C2—C7—H7	120.00
C1—C2—C3	121.2 (3)	C6—C7—H7	120.00
O1—C1—C2—C3	179.2 (3)	O3—C3—C4—C5	179.4 (3)
O1—C1—C2—C7	-1.8 (4)	C2—C3—C4—C5	-1.6 (5)
O2—C1—C2—C3	-1.2 (5)	C3—C4—C5—O4	179.3 (3)
O2—C1—C2—C7	177.8 (3)	C3—C4—C5—C6	-0.2 (4)
C1—C2—C3—O3	-0.2 (4)	O4—C5—C6—N1	2.6 (4)
C1—C2—C3—C4	-179.2 (3)	O4—C5—C6—C7	-177.7 (3)
C7—C2—C3—O3	-179.2 (3)	C4—C5—C6—N1	-177.8 (3)
C7—C2—C3—C4	1.8 (4)	C4—C5—C6—C7	1.9 (4)
C1—C2—C7—C6	-179.2 (3)	N1—C6—C7—C2	178.0 (3)
C3—C2—C7—C6	-0.2 (4)	C5—C6—C7—C2	-1.7 (4)

Symmetry codes: (i) $x, y, z+1$; (ii) $x+1, y, z+1$; (iii) $x+1, y, z$; (iv) $x+1/2, -y+1/2, z+1/2$; (v) $-x+2, -y, -z$; (vi) $-x+1, -y, -z$; (vii) $-x+1, -y, -z+1$; (viii) $x-1, y, z$; (ix) $x, y, z-1$; (x) $x-1, y, z-1$; (xi) $x-1/2, -y+1/2, z-1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)


$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
----------------------	--------------	-------------	-------------	----------------------

supplementary materials

O1—H1···O2 ^v	0.85 (3)	1.81 (3)	2.653 (3)	177 (4)
N1—H1A···Cl1 ^x	0.89	2.24	3.124 (3)	176
N1—H1B···Cl1 ^{ix}	0.89	2.29	3.176 (3)	173
N1—H1C···Cl1 ^{xi}	0.89	2.35	3.217 (2)	163
O3—H3···O2	0.85 (4)	1.84 (3)	2.632 (3)	154 (3)
O4—H4A···Cl1 ^{viii}	0.85 (3)	2.15 (3)	2.985 (2)	170 (3)

Symmetry codes: (v) $-x+2, -y, -z$; (x) $x-1, y, z-1$; (ix) $x, y, z-1$; (xi) $x-1/2, -y+1/2, z-1/2$; (viii) $x-1, y, z$.

Fig. 1



supplementary materials

Fig. 2

