

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(*E,E*)-1,2-Bis(2,4,6-trimethoxybenzylidene)hydrazineHoong-Kun Fun,^{a,*} Patcharaporn Jansrisewangwong^b and Suchada Chantrapromma^c§

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bDepartment of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand, and ^cCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand

Correspondence e-mail: hkfun@usm.my


Received 8 August 2010; accepted 20 August 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.040; wR factor = 0.115; data-to-parameter ratio = 20.8.

The title molecule, $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_6$, lies on an inversion centre. All non-H atoms are essentially coplanar, with an r.m.s. deviation of 0.0415 (1) Å and a maximum deviation of 0.1476 (1) Å for the methoxy C atom at the 4-position of the benzene ring. The crystal structure is stabilized by weak $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For standard bond-length data, see: Allen *et al.* (1987). For related structures, see: Jansrisewangwong *et al.* (2010); Zhao *et al.* (2006). For background and the biological activity of hydrozones, see: El-Tabl *et al.* (2008); Qin *et al.* (2009); Ramamohan *et al.* (1995); Rollas & Küçüküzgel (2007). For the stability of the temperature controller used in the data collection, see Cosier & Glazer, (1986).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_6$
 $M_r = 388.41$
 Triclinic, $P\bar{1}$
 $a = 7.3851$ (2) Å

$b = 7.4043$ (2) Å
 $c = 9.5440$ (2) Å
 $\alpha = 71.412$ (1)°
 $\beta = 78.095$ (1)°

* Thomson Reuters ResearcherID: A-3561-2009.

§ Additional correspondence author, e-mail: suchada.c@psu.ac.th. Thomson Reuters ResearcherID: A-5085-2009.

$\gamma = 79.449$ (1)°
 $V = 480.13$ (2) Å³
 $Z = 1$
 Mo $K\alpha$ radiation

$\mu = 0.10$ mm⁻¹
 $T = 100$ K
 $0.29 \times 0.14 \times 0.08$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.972$, $T_{\max} = 0.992$

11100 measured reflections
 2791 independent reflections
 2244 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.115$
 $S = 1.03$
 2791 reflections
 134 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.42$ e Å⁻³
 $\Delta\rho_{\min} = -0.23$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg is the centroid of the C1–C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C10}-\text{H10B}\cdots\text{N1}^i$	0.96	2.49	3.3876 (15)	155
$\text{C8}-\text{H8C}\cdots\text{Cg}^{ii}$	0.97	2.79	3.6678 (13)	152
$\text{C10}-\text{H10C}\cdots\text{Cg}^{iii}$	0.97	2.63	3.4385 (13)	142

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $-x + 1, -y + 2, -z$; (iii) $-x + 1, -y + 1, -z + 1$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

PJ thanks the Center of Excellence for Innovation in Chemistry (PERCH-CIC), the Commission on Higher Education, Ministry of Education and the Graduate School, Prince of Songkla University, for financial support. The authors thank the Prince of Songkla University for financial support through the Crystal Materials Research Unit (CMRU) and Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5117).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
 El-Tabl, A. S., El-Saied, F. A., Plass, W. & Al-Hakimi, A. N. (2008). *Spectrochim. Acta Part A*, **71**, 90–99.
 Jansrisewangwong, P., Chantrapromma, S. & Fun, H.-K. (2010). *Acta Cryst. E* **66**, o2170.
 Qin, D.-D., Yang, Z.-Y. & Qi, G.-F. (2009). *Spectrochim. Acta Part A*, **74**, 415–420.

Ramamohan, L., Shikkaragol, R. K., Angadi, S. D. & Kulkarni, V. H. (1995).
Asian J. Pure Appl. Chem. **1**, 86.
Rollas, S. & Küçükgülzel, Ş. G. (2007). *Molecules*, **12**, 1910–1939.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D36**, 148–155.
Zhao, B., Feng, F., Guo, G. & Wang, W. (2006). *Acta Cryst.* **E62**, o2413–o2414.

supplementary materials

Acta Cryst. (2010). E66, o2401-o2402 [doi:10.1107/S1600536810033684]

(*E,E*)-1,2-Bis(2,4,6-trimethoxybenzylidene)hydrazine

H.-K. Fun, P. Jansrisewangwong and S. Chantrapromma

Comment

Hydrazones and their complexes are interesting due to their fluorescence properties (Qin *et al.*, 2009) and various biological activities such as insecticidal, antitumor, antioxidant, antifungal, antibacterial and antiviral properties (El-Tabl *et al.*, 2008; Ramamohan *et al.*, 1995; Rollas & Küçükgülzel, 2007). These interesting properties led us to synthesize the title hydrazone derivative (I) in order to study its antibacterial activity and fluorescence property. Experiments show that (I) does not possess antibacterial activities, however it does exhibit fluorescence with the maximum emission at 410 nm when the compound is excited at 280 nm. Herein the crystal structure of (I) is reported.

The asymmetric unit of (I), (Fig. 1), $C_{20}H_{24}N_2O_6$, contains one half-molecule and the complete molecule is generated by an inversion centre (symmetry code $-x, 2-y, 1-z$). The mean plane through the C=N-N=C bridge forms a dihedral angle of $4.96(9)^\circ$ with the benzene rings. The methoxy groups attached to atoms C1 and C5 (positions 2 and 6) are approximately coplanar with the benzene ring whereas the one attached to atom C3 (position 4) is slightly twisted with respect to the benzene ring as described by the torsion angles of C8–O1–C1–C2 = $2.86(15)^\circ$, C10–O3–C5–C4 = $3.58(14)^\circ$ and C9–O2–C3–C4 = $8.39(15)^\circ$, respectively. The N–N bond length, $1.4117(18) \text{ \AA}$ is comparable with $1.419(3) \text{ \AA}$ and the C=N–N angle = $110.7(2)^\circ$, is almost similar to $112.2(2)^\circ$ observed in (*E,E*)-1,2-bis(3,4,5-trimethoxybenzylidene)hydrazine (Zhao *et al.*, 2006). The bond distances have normal values (Allen *et al.*, 1987) and are comparable with related structures (Jansrisewangwong *et al.*, 2010; Zhao *et al.*, 2006). The crystal structure is stabilized by weak C—H \cdots N and C—H \cdots π interactions (Fig. 2).

Experimental

The title compound was synthesized by mixing a solution (1:2 molar ratio) of hydrazine hydrate (0.097 ml, 2 mmol) and 2,4,6-trimethoxybenzaldehyde (0.785 mg, 4 mmol) in ethanol (20 ml). The resulting solution was refluxed for 5 h, yielding the yellow solid. The resultant solid was filtered off and washed with methanol. Yellow block-shaped single crystals of the title compound suitable for *x*-ray structure determination were recrystallized from acetone by slow evaporation of the solvent at room temperature over several days, mp. 484–486 K.

Refinement

The H atom attached to C7 was located in a difference map and refined isotropically. The remaining H atoms were positioned geometrically and allowed to ride on their parent atoms, with $d(C-H) = 0.93 \text{ \AA}$ for aromatic and 0.96 \AA for CH_3 atoms. The U_{iso} values were constrained to be $1.5U_{eq}$ of the carrier atom for methyl H atoms and $1.2U_{eq}$ for the remaining H atoms. A rotating group model was used for the methyl groups.

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atom-numbering scheme. Atoms with suffix A were generated by symmetry code $-x, 2 - y, 1 - z$.




Fig. 2. Part of the crystal structure showing weak hydrogen bonds as dashed lines.

(*E,E*)-1,2-Bis(2,4,6-trimethoxybenzylidene)hydrazine

Crystal data

$C_{20}H_{24}N_2O_6$

$M_r = 388.41$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.3851\ (2)\ \text{\AA}$

$b = 7.4043\ (2)\ \text{\AA}$

$c = 9.5440\ (2)\ \text{\AA}$

$\alpha = 71.412\ (1)^\circ$

$\beta = 78.095\ (1)^\circ$

$\gamma = 79.449\ (1)^\circ$

$V = 480.13\ (2)\ \text{\AA}^3$

$Z = 1$

$F(000) = 206$

$D_x = 1.343\ \text{Mg m}^{-3}$

Melting point = 484–486 K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2791 reflections

$\theta = 2.3\text{--}30.0^\circ$

$\mu = 0.10\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Block, yellow

$0.29 \times 0.14 \times 0.08\ \text{mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer

Radiation source: sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2005)

$T_{\min} = 0.972$, $T_{\max} = 0.992$

11100 measured reflections

2791 independent reflections

2244 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 30.0^\circ$, $\theta_{\min} = 2.3^\circ$

$h = -10 \rightarrow 10$

$k = -10 \rightarrow 10$

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.115$

$S = 1.03$

2791 reflections

134 parameters

0 restraints

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0603P)^2 + 0.1087P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.23 \text{ e } \text{\AA}^{-3}$$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 120.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.20858 (11)	0.87907 (11)	0.08848 (8)	0.01756 (18)
O2	0.82788 (11)	0.57545 (11)	0.12544 (8)	0.01906 (19)
O3	0.44058 (10)	0.78425 (11)	0.53298 (8)	0.01517 (17)
N1	0.09376 (12)	0.95751 (13)	0.49155 (10)	0.0163 (2)
C1	0.35619 (14)	0.81164 (14)	0.16438 (11)	0.0142 (2)
C2	0.52565 (15)	0.72803 (15)	0.10450 (11)	0.0158 (2)
H2A	0.5443	0.7178	0.0078	0.019*
C3	0.66759 (14)	0.65957 (15)	0.19168 (11)	0.0148 (2)
C4	0.64415 (14)	0.67699 (14)	0.33540 (11)	0.0143 (2)
H4A	0.7406	0.6318	0.3917	0.017*
C5	0.47341 (14)	0.76345 (14)	0.39383 (10)	0.0133 (2)
C6	0.32381 (14)	0.83151 (14)	0.31125 (11)	0.0134 (2)
C7	0.14037 (15)	0.91807 (15)	0.36584 (11)	0.0147 (2)
C8	0.22823 (17)	0.85390 (17)	-0.05710 (12)	0.0205 (2)
H8A	0.1127	0.9008	-0.0954	0.031*
H8B	0.2601	0.7199	-0.0499	0.031*
H8C	0.3251	0.9241	-0.1234	0.031*
C9	0.96932 (16)	0.47848 (17)	0.21582 (12)	0.0212 (2)
H9A	1.0697	0.4173	0.1585	0.032*
H9B	0.9173	0.3834	0.3022	0.032*
H9C	1.0157	0.5698	0.2473	0.032*
C10	0.59168 (15)	0.72427 (16)	0.61730 (11)	0.0167 (2)
H10A	0.5516	0.7495	0.7123	0.025*

supplementary materials

H10B	0.6945	0.7939	0.5631	0.025*
H10C	0.6302	0.5892	0.6330	0.025*
H7	0.0407 (19)	0.9430 (19)	0.3064 (15)	0.022 (3)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0168 (4)	0.0235 (4)	0.0134 (3)	0.0022 (3)	-0.0047 (3)	-0.0080 (3)
O2	0.0149 (4)	0.0236 (4)	0.0163 (4)	0.0049 (3)	0.0002 (3)	-0.0082 (3)
O3	0.0145 (4)	0.0196 (4)	0.0124 (3)	0.0013 (3)	-0.0034 (3)	-0.0073 (3)
N1	0.0125 (4)	0.0186 (4)	0.0173 (4)	-0.0003 (3)	-0.0004 (3)	-0.0069 (3)
C1	0.0156 (5)	0.0131 (4)	0.0133 (4)	-0.0010 (4)	-0.0031 (4)	-0.0032 (4)
C2	0.0175 (5)	0.0173 (5)	0.0122 (4)	-0.0009 (4)	-0.0001 (4)	-0.0058 (4)
C3	0.0134 (5)	0.0145 (5)	0.0150 (4)	-0.0007 (4)	0.0012 (4)	-0.0049 (4)
C4	0.0133 (5)	0.0149 (5)	0.0144 (4)	-0.0008 (4)	-0.0028 (4)	-0.0043 (4)
C5	0.0153 (5)	0.0120 (4)	0.0123 (4)	-0.0028 (4)	-0.0003 (4)	-0.0040 (3)
C6	0.0140 (5)	0.0134 (4)	0.0126 (4)	-0.0010 (4)	-0.0014 (3)	-0.0046 (4)
C7	0.0137 (5)	0.0151 (5)	0.0151 (4)	-0.0006 (4)	-0.0027 (4)	-0.0049 (4)
C8	0.0240 (6)	0.0251 (6)	0.0143 (5)	0.0020 (4)	-0.0061 (4)	-0.0093 (4)
C9	0.0159 (5)	0.0242 (5)	0.0198 (5)	0.0043 (4)	-0.0014 (4)	-0.0061 (4)
C10	0.0157 (5)	0.0207 (5)	0.0157 (4)	-0.0010 (4)	-0.0051 (4)	-0.0071 (4)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.3632 (12)	C4—H4A	0.9300
O1—C8	1.4347 (12)	C5—C6	1.4135 (14)
O2—C3	1.3642 (12)	C6—C7	1.4564 (14)
O2—C9	1.4328 (13)	C7—H7	0.976 (14)
O3—C5	1.3528 (11)	C8—H8A	0.9600
O3—C10	1.4322 (12)	C8—H8B	0.9600
N1—C7	1.2882 (13)	C8—H8C	0.9600
N1—N1 ⁱ	1.4117 (18)	C9—H9A	0.9600
C1—C2	1.3866 (14)	C9—H9B	0.9600
C1—C6	1.4226 (13)	C9—H9C	0.9600
C2—C3	1.3944 (15)	C10—H10A	0.9600
C2—H2A	0.9300	C10—H10B	0.9600
C3—C4	1.3909 (13)	C10—H10C	0.9600
C4—C5	1.3974 (14)		
C1—O1—C8	118.01 (8)	N1—C7—C6	125.41 (9)
C3—O2—C9	117.79 (8)	N1—C7—H7	115.8 (8)
C5—O3—C10	117.61 (8)	C6—C7—H7	118.7 (8)
C7—N1—N1 ⁱ	110.66 (11)	O1—C8—H8A	109.5
O1—C1—C2	122.94 (9)	O1—C8—H8B	109.5
O1—C1—C6	115.10 (9)	H8A—C8—H8B	109.5
C2—C1—C6	121.95 (9)	O1—C8—H8C	109.5
C1—C2—C3	118.90 (9)	H8A—C8—H8C	109.5
C1—C2—H2A	120.5	H8B—C8—H8C	109.5
C3—C2—H2A	120.5	O2—C9—H9A	109.5

O2—C3—C4	123.44 (9)	O2—C9—H9B	109.5
O2—C3—C2	115.02 (9)	H9A—C9—H9B	109.5
C4—C3—C2	121.55 (9)	O2—C9—H9C	109.5
C3—C4—C5	118.99 (9)	H9A—C9—H9C	109.5
C3—C4—H4A	120.5	H9B—C9—H9C	109.5
C5—C4—H4A	120.5	O3—C10—H10A	109.5
O3—C5—C4	122.15 (9)	O3—C10—H10B	109.5
O3—C5—C6	116.17 (9)	H10A—C10—H10B	109.5
C4—C5—C6	121.67 (9)	O3—C10—H10C	109.5
C5—C6—C1	116.93 (9)	H10A—C10—H10C	109.5
C5—C6—C7	124.92 (9)	H10B—C10—H10C	109.5
C1—C6—C7	118.15 (9)		
C8—O1—C1—C2	2.86 (15)	C3—C4—C5—C6	0.69 (15)
C8—O1—C1—C6	-176.61 (9)	O3—C5—C6—C1	179.70 (8)
O1—C1—C2—C3	-178.70 (9)	C4—C5—C6—C1	-1.27 (15)
C6—C1—C2—C3	0.74 (16)	O3—C5—C6—C7	-0.86 (15)
C9—O2—C3—C4	8.39 (15)	C4—C5—C6—C7	178.16 (9)
C9—O2—C3—C2	-171.46 (9)	O1—C1—C6—C5	-179.97 (8)
C1—C2—C3—O2	178.49 (9)	C2—C1—C6—C5	0.55 (15)
C1—C2—C3—C4	-1.37 (16)	O1—C1—C6—C7	0.55 (14)
O2—C3—C4—C5	-179.18 (9)	C2—C1—C6—C7	-178.93 (9)
C2—C3—C4—C5	0.67 (16)	N1 ⁱ —N1—C7—C6	179.28 (10)
C10—O3—C5—C4	3.58 (14)	C5—C6—C7—N1	5.52 (17)
C10—O3—C5—C6	-177.40 (8)	C1—C6—C7—N1	-175.05 (10)
C3—C4—C5—O3	179.66 (9)		

Symmetry codes: (i) $-x, -y+2, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

C_g is the centroid of the C1–C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C7—H7 [⋯] O1	0.977 (14)	2.332 (14)	2.6886 (12)	100.6 (10)
C10—H10B [⋯] N1 ⁱⁱ	0.96	2.49	3.3876 (15)	155
C8—H8C [⋯] C _g ⁱⁱⁱ	0.97	2.79	3.6678 (13)	152
C10—H10C [⋯] C _g ^{iv}	0.97	2.63	3.4385 (13)	142

Symmetry codes: (ii) $-x+1, -y+2, -z+1$; (iii) $-x+1, -y+2, -z$; (iv) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

