

Ethyl 7-(4-bromophenyl)-5-trifluoromethyl-4,7-dihydrotetrazolo[1,5-a]-pyrimidine-6-carboxylate

Shi-De Shen,^a Xiao-Dong Feng,^{b,c} Wei-Hua Yang^b and Chang-Sheng Yao^{b,c*}

^aXuzhou Institute of Architectural Technology, Xuzhou 221116, People's Republic of China, ^bSchool of Chemistry and Chemical Engineering, Xuzhou Normal University, Xuzhou 221116, People's Republic of China, and ^cKey Laboratory of Biotechnology for Medicinal Plants, Xuzhou Normal University, Xuzhou 221116, People's Republic of China

Correspondence e-mail: csyao@xznu.edu.cn


Received 18 August 2010; accepted 22 August 2010

Key indicators: single-crystal X-ray study; $T = 113\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.030; wR factor = 0.071; data-to-parameter ratio = 15.9.

In the title compound, $\text{C}_{14}\text{H}_{11}\text{BrF}_3\text{N}_5\text{O}_2$, the pyrimidine ring adopts a flattened envelope conformation with sp^3 -hybridized carbon as the flap [deviation = 0.177 (3) \AA]. The dihedral angle between tetrazole and bromophenyl rings is 84.3 (1) $^\circ$. In the crystal, molecules are linked into centrosymmetric dimers by pairs of N—H \cdots N hydrogen bonds.

Related literature

For the biological activity of tetrazolopyrimidine derivatives, see: Von Nussbaum *et al.* (2010); Abelman *et al.* (2009); Dougherty *et al.* (2007). For ring puckering parameters, see: Cremer & Pople (1975). For the synthesis, see: Pryadeina *et al.* (2004).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{11}\text{BrF}_3\text{N}_5\text{O}_2$

$M_r = 418.19$

Monoclinic, $P2_1/c$
 $a = 18.773 (2)\text{ \AA}$
 $b = 10.4716 (11)\text{ \AA}$
 $c = 7.8700 (8)\text{ \AA}$
 $\beta = 92.27 (3)^\circ$
 $V = 1545.9 (3)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 2.71\text{ mm}^{-1}$
 $T = 113\text{ K}$
 $0.32 \times 0.28 \times 0.18\text{ mm}$

Data collection

Rigaku Saturn diffractometer
Absorption correction: multi-scan
(*CrystalClear*; Rigaku/MSC, 2002)
 $T_{\min} = 0.477$, $T_{\max} = 0.641$

18785 measured reflections
3681 independent reflections
2547 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.065$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.071$
 $S = 0.94$
3681 reflections
231 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.43\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.82\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}1-\text{H}1\cdots\text{N}5^{\dagger}$	0.83 (2)	2.06 (2)	2.862 (2)	163 (2)

Symmetry code: (i) $-x, -y, -z$.

Data collection: *CrystalClear* (Rigaku/MSC, 2002); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors are grateful for financial support from the Major Basic Research Project of the Natural Science Foundation of Jiangsu Higher Education Institutions (No. 09 KJ A430003), the Natural Science Foundation of Xuzhou City (No. XM09B016), the Graduate Foundation of Xuzhou Normal University (No. 2010YLB029) and the Qing Lan Project (No. 08QLT001).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5159).

References

- Abelman, M., Jiang, R. & Zablocki, J. (2009). WO Patent No. 2009006580.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Dougherty, A. M., Guo, H., Westby, G., Liu, Y., Simsek, E., Guo, J. T., Mehta, A., Norton, P., Gu, B., Block, T. & Cuconati, A. (2007). *Antimicrob. Agents Chemother.* **51**, 4427–4437.
- Pryadeina, M. V., Burgart, Ya. V., Saloutin, V. I., Kodess, M. I., Ulomskii, E. N. & Rusinov, V. L. (2004). *Russ. J. Org. Chem.* **40**, 902–907.
- Rigaku/MSC (2002). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Von Nussbaum, F., Karthaus, D., Anlauf, S., Delbeck, M., Li, V. M., Meibom, D. & Lustig, K. (2010). WO Patent No. 2010078953.

supplementary materials

Acta Cryst. (2010). E66, o2421 [doi:10.1107/S1600536810033842]

Ethyl 7-(4-bromophenyl)-5-trifluoromethyl-4,7-dihydrotetrazolo[1,5-*a*]pyrimidine-6-carboxylate

S.-D. Shen, X.-D. Feng, W.-H. Yang and C.-S. Yao

Comment

The tetrazolo[1,5-*a*]pyrimidine core represents an interesting pharmacophore with the feature of biological and pharmacological properties, which has human neutrophil elastase inhibitory (Von Nussbaum *et al.*, 2010), late sodium channel blocker (Abelman *et al.*, 2009) and hepatitis B virus surface antigen secretion inhibitory activities (Dougherty *et al.*, 2007). This led us to pay much attention to the synthesis and bioactivity of compounds containing these two significant fragments. During the synthesis of trifluoromethylated tetrazolo[1,5-*a*]pyrimidine derivatives, the title compound was isolated and its structure was determined by X-ray analysis. The results are presented here.

In the title molecule (Fig.1), the tetrahydropyrimidine ring is in a flattened envelope conformation, with Cremer and Pople (1975) puckering parameters Q , θ , ϕ of 0.125 (2) Å, 109.7 (9)° and 11.7 (9)°, respectively; atom C2 deviates from the N1/N2/C1/C3/C4 plane (r.m.s. deviation 0.018 Å) by 0.177 (3) Å. The dihedral angle between N1/N2/C1/C3/C4 and C5-C10 planes [89.53 (3)°] shows that they are nearly perpendicular.

The crystal packing is stabilized by intermolecular N—H···N hydrogen bonds (Table 1 and Fig.2).

Experimental

The title compound was synthesized according the procedure reported by Pryadeina *et al.* (2004). A mixture of ethyl 4,4,4-trifluoro-3-oxobutanoate (0.01 mol), 4-bromobenzaldehyde (0.01 mol) and 5-aminotetrazole (0.01 mol) in ethanol (20 ml) containing a catalytic amount of hydrochloric acid was heated for 12 h under reflux. Then the solvent was removed under reduced pressure. The residue was added to a solution of p-toluenesulfonic acid (0.05 g) in 100 mL of benzene, and the mixture was heated for 8 h with simultaneous removal of water as azeotrope with benzene. The solution was filtered while hot, the filtrate was evaporated, and the precipitate was recrystallized from ethanol. Cooling the ethanol solution slowly gave single crystals suitable for X-ray diffraction.

Refinement

The N-bound H atom was located in a difference map and was refined freely [refined N—H length, 0.83 (2) Å]. All other H atoms were placed in calculated positions [C—H = 0.95–1.00 Å] and included in the final cycles of refinement using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{parent atom})$.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. A packing diagram of the title compound. Intermolecular hydrogen bonds are shown as dashed lines.

Ethyl 7-(4-bromophenyl)-5-trifluoromethyl-4,7-dihydrotetrazolo[1,5-a]pyrimidine-6-carboxylate

Crystal data

C₁₄H₁₁BrF₃N₅O₂

F(000) = 832

M_r = 418.19

D_x = 1.797 Mg m⁻³

Monoclinic, *P*2₁/c

Mo *K*α radiation, λ = 0.71073 Å

Hall symbol: -P 2ybc

Cell parameters from 4823 reflections

a = 18.773 (2) Å

θ = 2.2–27.9°

b = 10.4716 (11) Å

μ = 2.71 mm⁻¹

c = 7.8700 (8) Å

T = 113 K

β = 92.27 (3)°

Block, colourless

V = 1545.9 (3) Å³

0.32 × 0.28 × 0.18 mm

$Z = 4$

Data collection

Rigaku Saturn diffractometer	3681 independent reflections
Radiation source: rotating anode confocal	2547 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.065$
Detector resolution: 7.31 pixels mm ⁻¹	$\theta_{\text{max}} = 27.9^\circ$, $\theta_{\text{min}} = 2.2^\circ$
ω scans	$h = -24 \rightarrow 24$
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku/MSC, 2002)	$k = -13 \rightarrow 13$
$T_{\text{min}} = 0.477$, $T_{\text{max}} = 0.641$	$l = -10 \rightarrow 10$
18785 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.030$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.071$	H atoms treated by a mixture of independent and constrained refinement
$S = 0.94$	$w = 1/[\sigma^2(F_o^2) + (0.0321P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
3681 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
231 parameters	$\Delta\rho_{\text{max}} = 0.43 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.82 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.468171 (12)	0.24266 (2)	-0.00430 (3)	0.02485 (8)
F1	0.04617 (6)	0.09175 (12)	0.60550 (15)	0.0283 (3)
F2	0.00359 (6)	0.18117 (13)	0.37917 (16)	0.0312 (3)
F3	0.10127 (7)	0.25030 (10)	0.49390 (17)	0.0279 (3)

supplementary materials

H1	0.0275 (12)	0.055 (2)	0.176 (3)	0.033 (7)*
N1	0.06837 (9)	0.02879 (16)	0.1959 (2)	0.0142 (4)
N2	0.16300 (9)	-0.09820 (15)	0.1113 (2)	0.0139 (4)
N3	0.17768 (9)	-0.17880 (16)	-0.0190 (2)	0.0195 (4)
N4	0.12129 (9)	-0.17855 (17)	-0.1192 (2)	0.0196 (4)
N5	0.06960 (9)	-0.09971 (15)	-0.0596 (2)	0.0163 (4)
O1	0.18713 (8)	0.08260 (14)	0.67228 (18)	0.0266 (4)
O2	0.27052 (7)	-0.04022 (14)	0.55633 (17)	0.0209 (3)
C1	0.09759 (10)	-0.05253 (18)	0.0835 (2)	0.0134 (4)
C2	0.21318 (10)	-0.06715 (18)	0.2535 (2)	0.0143 (4)
H2	0.2315	-0.1483	0.3062	0.017*
C3	0.17113 (10)	0.00649 (18)	0.3842 (2)	0.0147 (4)
C4	0.10521 (11)	0.05403 (18)	0.3465 (2)	0.0141 (4)
C5	0.27570 (10)	0.00933 (19)	0.1898 (2)	0.0143 (4)
C6	0.26514 (11)	0.12937 (19)	0.1182 (3)	0.0185 (5)
H6	0.2184	0.1642	0.1091	0.022*
C7	0.32231 (11)	0.1991 (2)	0.0599 (3)	0.0187 (5)
H7	0.3150	0.2809	0.0097	0.022*
C8	0.39009 (11)	0.14732 (19)	0.0761 (2)	0.0172 (5)
C9	0.40180 (11)	0.02870 (19)	0.1482 (3)	0.0196 (5)
H9	0.4487	-0.0054	0.1587	0.024*
C10	0.34446 (11)	-0.03993 (19)	0.2049 (3)	0.0185 (5)
H10	0.3521	-0.1216	0.2548	0.022*
C11	0.20751 (11)	0.02294 (19)	0.5533 (3)	0.0174 (4)
C12	0.31205 (11)	-0.0374 (2)	0.7163 (3)	0.0251 (5)
H12A	0.2901	-0.0933	0.8012	0.030*
H12B	0.3145	0.0506	0.7620	0.030*
C13	0.38555 (12)	-0.0847 (2)	0.6780 (3)	0.0290 (6)
H13A	0.3820	-0.1701	0.6276	0.043*
H13B	0.4150	-0.0884	0.7835	0.043*
H13C	0.4074	-0.0262	0.5979	0.043*
C14	0.06429 (11)	0.1435 (2)	0.4593 (3)	0.0205 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.01555 (14)	0.03165 (14)	0.02782 (14)	-0.00341 (9)	0.00700 (9)	0.00705 (10)
F1	0.0232 (8)	0.0413 (8)	0.0212 (7)	-0.0037 (6)	0.0114 (6)	-0.0076 (6)
F2	0.0184 (7)	0.0410 (8)	0.0338 (8)	0.0131 (6)	-0.0035 (6)	-0.0185 (7)
F3	0.0254 (8)	0.0205 (7)	0.0380 (8)	-0.0022 (5)	0.0034 (6)	-0.0136 (6)
N1	0.0094 (10)	0.0162 (9)	0.0174 (9)	0.0012 (7)	0.0031 (7)	-0.0026 (7)
N2	0.0110 (9)	0.0147 (8)	0.0163 (9)	0.0004 (7)	0.0036 (7)	-0.0035 (7)
N3	0.0184 (10)	0.0208 (10)	0.0194 (9)	0.0007 (8)	0.0033 (8)	-0.0074 (8)
N4	0.0152 (10)	0.0220 (10)	0.0218 (10)	0.0018 (8)	0.0027 (8)	-0.0063 (8)
N5	0.0145 (10)	0.0169 (9)	0.0178 (9)	0.0003 (7)	0.0045 (7)	-0.0036 (7)
O1	0.0224 (9)	0.0357 (9)	0.0217 (8)	0.0044 (7)	0.0022 (7)	-0.0102 (7)
O2	0.0175 (8)	0.0271 (8)	0.0178 (8)	0.0040 (6)	-0.0011 (6)	-0.0042 (6)
C1	0.0111 (11)	0.0130 (9)	0.0163 (10)	-0.0016 (8)	0.0050 (8)	0.0019 (8)

C2	0.0122 (11)	0.0140 (10)	0.0167 (10)	0.0005 (8)	0.0012 (8)	-0.0019 (8)
C3	0.0132 (11)	0.0136 (9)	0.0176 (10)	-0.0029 (8)	0.0051 (8)	-0.0018 (8)
C4	0.0137 (11)	0.0135 (10)	0.0154 (10)	-0.0030 (8)	0.0058 (8)	-0.0013 (8)
C5	0.0137 (11)	0.0164 (10)	0.0130 (10)	-0.0005 (8)	0.0038 (8)	-0.0029 (8)
C6	0.0124 (12)	0.0203 (11)	0.0230 (11)	0.0033 (9)	0.0031 (9)	0.0007 (9)
C7	0.0197 (12)	0.0163 (10)	0.0204 (11)	0.0023 (9)	0.0038 (9)	0.0013 (9)
C8	0.0152 (12)	0.0216 (11)	0.0153 (10)	-0.0036 (9)	0.0052 (8)	-0.0008 (9)
C9	0.0107 (11)	0.0248 (12)	0.0236 (12)	0.0045 (9)	0.0037 (9)	0.0011 (9)
C10	0.0168 (12)	0.0170 (10)	0.0220 (11)	0.0036 (9)	0.0047 (9)	0.0015 (9)
C11	0.0148 (12)	0.0175 (10)	0.0203 (11)	-0.0032 (9)	0.0037 (9)	0.0003 (9)
C12	0.0193 (13)	0.0366 (14)	0.0192 (11)	0.0028 (10)	-0.0031 (9)	0.0003 (10)
C13	0.0206 (13)	0.0371 (14)	0.0292 (13)	0.0054 (11)	-0.0004 (10)	0.0039 (11)
C14	0.0142 (12)	0.0238 (11)	0.0235 (12)	0.0006 (9)	0.0014 (9)	-0.0069 (10)

Geometric parameters (\AA , $^\circ$)

Br1—C8	1.9025 (19)	C3—C4	1.356 (3)
F1—C14	1.328 (2)	C3—C11	1.482 (3)
F2—C14	1.340 (2)	C4—C14	1.520 (3)
F3—C14	1.338 (2)	C5—C6	1.388 (3)
N1—C1	1.359 (2)	C5—C10	1.391 (3)
N1—C4	1.374 (3)	C6—C7	1.392 (3)
N1—H1	0.83 (2)	C6—H6	0.95
N2—C1	1.328 (2)	C7—C8	1.385 (3)
N2—N3	1.365 (2)	C7—H7	0.95
N2—C2	1.470 (2)	C8—C9	1.379 (3)
N3—N4	1.295 (2)	C9—C10	1.383 (3)
N4—N5	1.371 (2)	C9—H9	0.95
N5—C1	1.320 (2)	C10—H10	0.95
O1—C11	1.201 (2)	C12—C13	1.508 (3)
O2—C11	1.355 (2)	C12—H12A	0.99
O2—C12	1.455 (2)	C12—H12B	0.99
C2—C5	1.522 (2)	C13—H13A	0.98
C2—C3	1.530 (3)	C13—H13B	0.98
C2—H2	1.00	C13—H13C	0.98
C1—N1—C4	118.68 (17)	C8—C7—H7	120.6
C1—N1—H1	119.1 (17)	C6—C7—H7	120.6
C4—N1—H1	121.9 (16)	C9—C8—C7	121.45 (19)
C1—N2—N3	108.16 (16)	C9—C8—Br1	119.80 (15)
C1—N2—C2	127.32 (16)	C7—C8—Br1	118.75 (15)
N3—N2—C2	124.50 (16)	C8—C9—C10	119.16 (19)
N4—N3—N2	105.79 (16)	C8—C9—H9	120.4
N3—N4—N5	111.43 (16)	C10—C9—H9	120.4
C1—N5—N4	104.74 (16)	C9—C10—C5	120.74 (19)
C11—O2—C12	116.29 (15)	C9—C10—H10	119.6
N5—C1—N2	109.88 (17)	C5—C10—H10	119.6
N5—C1—N1	129.27 (18)	O1—C11—O2	123.05 (19)
N2—C1—N1	120.84 (18)	O1—C11—C3	127.68 (19)
N2—C2—C5	110.17 (15)	O2—C11—C3	109.27 (17)

supplementary materials

N2—C2—C3	106.92 (15)	O2—C12—C13	106.47 (17)
C5—C2—C3	112.42 (16)	O2—C12—H12A	110.4
N2—C2—H2	109.1	C13—C12—H12A	110.4
C5—C2—H2	109.1	O2—C12—H12B	110.4
C3—C2—H2	109.1	C13—C12—H12B	110.4
C4—C3—C11	122.55 (18)	H12A—C12—H12B	108.6
C4—C3—C2	121.92 (18)	C12—C13—H13A	109.5
C11—C3—C2	115.53 (17)	C12—C13—H13B	109.5
C3—C4—N1	122.81 (17)	H13A—C13—H13B	109.5
C3—C4—C14	125.21 (18)	C12—C13—H13C	109.5
N1—C4—C14	111.95 (17)	H13A—C13—H13C	109.5
C6—C5—C10	119.19 (18)	H13B—C13—H13C	109.5
C6—C5—C2	120.69 (18)	F1—C14—F3	108.26 (17)
C10—C5—C2	120.11 (18)	F1—C14—F2	106.58 (17)
C5—C6—C7	120.64 (19)	F3—C14—F2	105.95 (17)
C5—C6—H6	119.7	F1—C14—C4	114.01 (17)
C7—C6—H6	119.7	F3—C14—C4	111.34 (17)
C8—C7—C6	118.80 (19)	F2—C14—C4	110.28 (17)
C1—N2—N3—N4	0.1 (2)	C3—C2—C5—C6	-55.7 (2)
C2—N2—N3—N4	-178.49 (17)	N2—C2—C5—C10	-117.49 (19)
N2—N3—N4—N5	0.2 (2)	C3—C2—C5—C10	123.4 (2)
N3—N4—N5—C1	-0.4 (2)	C10—C5—C6—C7	1.0 (3)
N4—N5—C1—N2	0.4 (2)	C2—C5—C6—C7	-179.92 (18)
N4—N5—C1—N1	-178.44 (19)	C5—C6—C7—C8	-0.7 (3)
N3—N2—C1—N5	-0.3 (2)	C6—C7—C8—C9	0.2 (3)
C2—N2—C1—N5	178.17 (17)	C6—C7—C8—Br1	179.84 (15)
N3—N2—C1—N1	178.65 (16)	C7—C8—C9—C10	0.2 (3)
C2—N2—C1—N1	-2.8 (3)	Br1—C8—C9—C10	-179.52 (15)
C4—N1—C1—N5	172.99 (19)	C8—C9—C10—C5	0.1 (3)
C4—N1—C1—N2	-5.8 (3)	C6—C5—C10—C9	-0.7 (3)
C1—N2—C2—C5	-111.0 (2)	C2—C5—C10—C9	-179.76 (18)
N3—N2—C2—C5	67.3 (2)	C12—O2—C11—O1	2.7 (3)
C1—N2—C2—C3	11.5 (2)	C12—O2—C11—C3	-177.93 (16)
N3—N2—C2—C3	-170.26 (16)	C4—C3—C11—O1	-4.0 (3)
N2—C2—C3—C4	-13.1 (2)	C2—C3—C11—O1	175.58 (19)
C5—C2—C3—C4	108.0 (2)	C4—C3—C11—O2	176.71 (17)
N2—C2—C3—C11	167.39 (16)	C2—C3—C11—O2	-3.7 (2)
C5—C2—C3—C11	-71.6 (2)	C11—O2—C12—C13	-166.71 (17)
C11—C3—C4—N1	-173.55 (17)	C3—C4—C14—F1	-65.4 (3)
C2—C3—C4—N1	6.9 (3)	N1—C4—C14—F1	116.58 (19)
C11—C3—C4—C14	8.6 (3)	C3—C4—C14—F3	57.5 (3)
C2—C3—C4—C14	-170.92 (18)	N1—C4—C14—F3	-120.58 (19)
C1—N1—C4—C3	3.5 (3)	C3—C4—C14—F2	174.78 (18)
C1—N1—C4—C14	-178.41 (17)	N1—C4—C14—F2	-3.3 (2)
N2—C2—C5—C6	63.4 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A

D—H


H···A

D···A

D—H···A

N1—H1…N5ⁱ 0.83 (2) 2.06 (2) 2.862 (2) 163 (2)
Symmetry codes: (i) $-x, -y, -z$.

Fig. 1



supplementary materials

Fig. 2

