

5'-Amino-2-oxo-2',3'-dihydrospiro-[indoline-3,7'-thieno[3,2-*b*]pyran]-6'-carbonitrile 1',1'-dioxide

Shi-De Shen,^a Xiao-Dong Feng,^{b,c} Wei-Hua Yang^b and Chang-Sheng Yao^{b,c*}

^aXuzhou Institute of Architectural Technology, Xuzhou 221116, People's Republic of China, ^bSchool of Chemistry and Chemical Engineering, Xuzhou Normal University, Xuzhou 221116, People's Republic of China, and ^cKey Laboratory of Biotechnology for Medicinal Plants, Xuzhou Normal University, Xuzhou 221116, People's Republic of China

Correspondence e-mail: csyao@xznz.edu.cn


Received 30 July 2010; accepted 16 August 2010

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.036; wR factor = 0.099; data-to-parameter ratio = 18.9.

In the title compound, $\text{C}_{15}\text{H}_{11}\text{N}_3\text{O}_4\text{S}$, the dihedral angle between the mean planes of the dihydroindol-2-one (r.m.s. deviation = 0.015 Å) and dihydrothieno[3,2-*b*]pyran (r.m.s. deviation = 0.011 Å) ring systems is 89.53 (3)°. The crystal packing is consolidated by intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds, which link the molecules into a two-dimensional network into sheets lying parallel to (100).

Related literature

For the antiviral and $\alpha 2$ -adrenoreceptor agonist activity of thieno[3,2-*b*]pyran derivatives, see: Chao *et al.* (2009); Friary *et al.* (1991). For the biological and pharmacological properties of indole derivatives, see: Sundberg (1996).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{11}\text{N}_3\text{O}_4\text{S}$

$M_r = 329.33$

Monoclinic, $C2/c$
 $a = 30.669$ (4) Å
 $b = 8.1760$ (14) Å
 $c = 12.229$ (2) Å
 $\beta = 112.611$ (8)°
 $V = 2830.7$ (8) Å³

$Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.25$ mm⁻¹
 $T = 113$ K
 $0.25 \times 0.22 \times 0.20$ mm

Data collection

Rigaku Saturn724 CCD diffractometer
 Absorption correction: multi-scan (*CrystalClear-SM Expert*; Rigaku/MS, 2009)
 $T_{\min} = 0.939$, $T_{\max} = 0.951$

19659 measured reflections
 4149 independent reflections
 3670 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.099$
 $S = 1.08$
 4149 reflections
 220 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.44$ e Å⁻³
 $\Delta\rho_{\min} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{N3}^{\text{i}}$	0.89 (2)	2.21 (2)	3.0878 (15)	170 (2)
$\text{N1}-\text{H2}\cdots\text{O4}^{\text{ii}}$	0.88 (2)	2.01 (2)	2.8790 (14)	172 (1)
$\text{N2}-\text{H3}\cdots\text{O3}^{\text{iii}}$	0.84 (2)	2.13 (2)	2.9255 (14)	159 (2)

Symmetry codes: (i) $-x, -y + 2, -z$; (ii) $-x, -y + 1, -z$; (iii) $x, -y + 1, z - \frac{1}{2}$.

Data collection: *CrystalClear-SM Expert* (Rigaku/MS, 2009); cell refinement: *CrystalClear-SM Expert*; data reduction: *CrystalClear-SM Expert*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors are grateful for financial support from the Major Basic Research Project of the Natural Science Foundation of the Jiangsu Higher Education Institutions (No. 09 KJ A430003), the Natural Science Foundation of Xuzhou City (No. XM09B016), the Graduate Foundation of Xuzhou Normal University (No. 2010YLB029) and the Qing Lan Project (No. 08QLT001).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5148).

References

- Chao, J. H., Zheng, J. Y. & Aslanian, R. G. (2009). WO Patent No. 2009020578.
 Friary, R. J., Schwerdt, J. H. & Ganguly, A. K. (1991). US Patent No. 5034531.
 Rigaku/MS (2009). *CrystalClear-SM Expert*. Rigaku/MS, The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Sundberg, R. J. (1996). *The Chemistry of Indoles*. New York: Academic Press.

supplementary materials

Acta Cryst. (2010). E66, o2448 [doi:10.1107/S1600536810032861]

5'-Amino-2-oxo-2',3'-dihydrospiro[indoline-3,7'-thieno[3,2-*b*]pyran]-6'-carbonitrile 1',1'-dioxide

S.-D. Shen, X.-D. Feng, W.-H. Yang and C.-S. Yao

Comment

The indole core represents an interesting pharmacophore with the feature of biological and pharmacological properties (Sundberg, 1996). Thienopyranyl compounds, such as thieno[3,2-*b*]pyran derivatives, can be used as antiviral agents (Friary *et al.*, 1991) and α -2 C adrenoreceptor agonists (Chao *et al.*, 2009). This led us to pay much attention to the synthesis and bioactivity of compounds containing these two significant fragments. During the synthesis of spiro[indoline-3,7'-thieno[3,2-*b*]pyran] derivatives, the title compound was isolated and its structure was determined by X-ray analysis. The results are presented here.

In the title molecule (Fig. 1), the dihydroindole-2-one ring system is planar (r.m.s. deviation 0.015 Å); the largest deviation from the mean plane is 0.023 (1) Å for atom C3. The dihydrothieno[3,2-*b*]pyran unit is also planar with an r.m.s. deviation of 0.011 Å (maximum deviation from the plane is 0.022 Å for atom C2). The dihedral angle between the two units is 89.53 (3)°.

The crystal packing is stabilized by intermolecular N—H \cdots O and N—H \cdots N hydrogen bonds (Table 1) which link the molecules into a two-dimensional network parallel to the (100) [Fig.2].

Experimental

The title compound was synthesized by the reaction of dihydrothiophen-3(2*H*)-one-1,1-dioxide (1 mmol) and 2-(2-oxindolin-3-ylidene)malononitrile (1 mmol) catalyzed by piperidine (0.02 g) in 10 ml ethanol under reflux until completion (monitored by TLC). Cooling the reaction mixture slowly gave single crystals suitable for X-ray diffraction.

Refinement

The H atoms bound to N atoms were located in a difference map and were refined freely [refined N—H lengths, 0.89 (2), 0.88 (2) and 0.84 (2) Å]. All other H atoms were placed in calculated positions, with C—H = 0.95, or 0.99 Å, and included in the final cycles of refinement using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{parent atom})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. A packing diagram of the title compound. Intermolecular hydrogen bonds are shown as dashed lines.

5'-Amino-2-oxo-2',3'-dihydrospiro[indoline-3,7'-thieno[3,2-*b*]pyran]-6'- carbonitrile 1',1'-dioxide

Crystal data

$C_{15}H_{11}N_3O_4S$

$M_r = 329.33$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 30.669 (4) \text{ \AA}$

$b = 8.1760 (14) \text{ \AA}$

$c = 12.229 (2) \text{ \AA}$

$\beta = 112.611 (8)^\circ$

$V = 2830.7 (8) \text{ \AA}^3$

$Z = 8$

$F(000) = 1360$

$D_x = 1.546 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71075 \text{ \AA}$

Cell parameters from 6345 reflections

$\theta = 1.4\text{--}33.4^\circ$

$\mu = 0.25 \text{ mm}^{-1}$

$T = 113 \text{ K}$

Prism, colourless

$0.25 \times 0.22 \times 0.20 \text{ mm}$

Data collection

Rigaku Saturn724 CCD
diffractometer

Radiation source: rotating anode
multilayer

Detector resolution: $14.222 \text{ pixels mm}^{-1}$

ω scans

Absorption correction: multi-scan
(*CrystalClear-SM Expert*; Rigaku/MS, 2009)

$T_{\min} = 0.939$, $T_{\max} = 0.951$

19659 measured reflections

4149 independent reflections

3670 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

$\theta_{\max} = 30.1^\circ$, $\theta_{\min} = 1.4^\circ$

$h = -42 \rightarrow 43$

$k = -11 \rightarrow 11$

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.036$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.099$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.0595P)^2 + 0.5864P]$
4149 reflections	where $P = (F_o^2 + 2F_c^2)/3$
220 parameters	$(\Delta/\sigma)_{\max} = 0.001$
0 restraints	$\Delta\rho_{\max} = 0.44 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.147537 (9)	0.29057 (3)	0.24536 (2)	0.01647 (9)
O1	0.02172 (3)	0.46422 (10)	0.16158 (7)	0.01616 (16)
O2	0.16145 (3)	0.24351 (12)	0.15034 (8)	0.0273 (2)
O3	0.18485 (3)	0.33938 (10)	0.35503 (7)	0.01955 (17)
O4	0.10067 (3)	0.52484 (10)	-0.05002 (7)	0.02012 (18)
N1	-0.02109 (3)	0.68754 (13)	0.09053 (9)	0.0183 (2)
N2	0.16403 (3)	0.67827 (13)	0.06897 (9)	0.0202 (2)
N3	0.05132 (3)	0.96181 (13)	0.00106 (9)	0.0214 (2)
C1	0.02156 (4)	0.62030 (13)	0.12026 (9)	0.0145 (2)
C2	0.06085 (4)	0.68905 (13)	0.11213 (9)	0.0145 (2)
C3	0.10896 (3)	0.60632 (13)	0.15197 (9)	0.0142 (2)
C4	0.10317 (4)	0.43896 (13)	0.19455 (9)	0.0143 (2)
C5	0.06325 (4)	0.37984 (13)	0.19645 (9)	0.0145 (2)
C6	0.06143 (4)	0.21074 (13)	0.23920 (10)	0.0180 (2)
H6A	0.0382	0.1442	0.1757	0.022*
H6B	0.0520	0.2125	0.3081	0.022*
C7	0.11110 (4)	0.13877 (14)	0.27463 (11)	0.0209 (2)

supplementary materials

H7A	0.1100	0.0386	0.2282	0.025*
H7B	0.1241	0.1098	0.3598	0.025*
C8	0.05546 (4)	0.84058 (13)	0.05249 (9)	0.0159 (2)
C9	0.12351 (4)	0.59395 (14)	0.04311 (9)	0.0164 (2)
C10	0.18025 (4)	0.74686 (15)	0.18382 (10)	0.0191 (2)
C11	0.14933 (4)	0.70594 (13)	0.23842 (10)	0.0160 (2)
C12	0.15829 (4)	0.75477 (14)	0.35300 (10)	0.0192 (2)
H12	0.1377	0.7253	0.3911	0.023*
C13	0.19871 (4)	0.84902 (15)	0.41138 (11)	0.0246 (3)
H13	0.2057	0.8843	0.4903	0.029*
C14	0.22859 (4)	0.89130 (17)	0.35519 (12)	0.0290 (3)
H14	0.2556	0.9564	0.3964	0.035*
C15	0.22021 (4)	0.84121 (17)	0.24027 (12)	0.0278 (3)
H15	0.2409	0.8702	0.2022	0.033*
H1	-0.0266 (6)	0.792 (2)	0.0714 (15)	0.032 (4)*
H2	-0.0445 (6)	0.623 (2)	0.0855 (14)	0.030 (4)*
H3	0.1762 (6)	0.6867 (19)	0.0186 (15)	0.030 (4)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.01688 (14)	0.01640 (14)	0.01632 (15)	0.00426 (9)	0.00659 (11)	0.00220 (9)
O1	0.0148 (3)	0.0163 (4)	0.0191 (4)	0.0019 (3)	0.0083 (3)	0.0031 (3)
O2	0.0321 (5)	0.0310 (5)	0.0228 (4)	0.0128 (4)	0.0148 (4)	0.0019 (4)
O3	0.0151 (3)	0.0225 (4)	0.0195 (4)	0.0017 (3)	0.0049 (3)	0.0025 (3)
O4	0.0215 (4)	0.0236 (4)	0.0169 (4)	0.0021 (3)	0.0092 (3)	0.0017 (3)
N1	0.0147 (4)	0.0185 (5)	0.0227 (5)	0.0022 (4)	0.0083 (4)	0.0021 (4)
N2	0.0168 (4)	0.0264 (5)	0.0208 (5)	0.0008 (4)	0.0110 (4)	0.0044 (4)
N3	0.0212 (5)	0.0191 (5)	0.0243 (5)	0.0030 (4)	0.0092 (4)	0.0034 (4)
C1	0.0163 (4)	0.0159 (5)	0.0120 (5)	0.0013 (4)	0.0061 (4)	-0.0001 (4)
C2	0.0143 (5)	0.0152 (5)	0.0143 (5)	0.0024 (4)	0.0058 (4)	0.0012 (4)
C3	0.0128 (4)	0.0152 (5)	0.0151 (5)	0.0018 (4)	0.0057 (4)	0.0019 (4)
C4	0.0155 (5)	0.0148 (5)	0.0128 (5)	0.0022 (4)	0.0055 (4)	0.0021 (4)
C5	0.0163 (5)	0.0150 (5)	0.0125 (5)	0.0019 (4)	0.0057 (4)	0.0005 (4)
C6	0.0206 (5)	0.0158 (5)	0.0185 (5)	-0.0005 (4)	0.0085 (4)	0.0019 (4)
C7	0.0206 (5)	0.0150 (5)	0.0228 (6)	0.0002 (4)	0.0036 (4)	0.0034 (4)
C8	0.0137 (4)	0.0177 (5)	0.0165 (5)	0.0019 (4)	0.0060 (4)	-0.0012 (4)
C9	0.0167 (5)	0.0173 (5)	0.0171 (5)	0.0049 (4)	0.0088 (4)	0.0054 (4)
C10	0.0152 (5)	0.0200 (5)	0.0215 (6)	0.0022 (4)	0.0064 (4)	0.0050 (4)
C11	0.0131 (5)	0.0144 (5)	0.0186 (5)	0.0014 (3)	0.0040 (4)	0.0033 (4)
C12	0.0192 (5)	0.0163 (5)	0.0195 (5)	0.0017 (4)	0.0047 (4)	0.0023 (4)
C13	0.0216 (5)	0.0206 (6)	0.0231 (6)	0.0000 (4)	-0.0008 (5)	0.0007 (4)
C14	0.0179 (5)	0.0263 (6)	0.0339 (7)	-0.0048 (5)	0.0001 (5)	0.0032 (5)
C15	0.0157 (5)	0.0304 (7)	0.0344 (7)	-0.0034 (5)	0.0065 (5)	0.0075 (5)

Geometric parameters (\AA , $^\circ$)

S1—O2	1.4359 (9)	C3—C9	1.5617 (15)
S1—O3	1.4448 (9)	C4—C5	1.3248 (14)

S1—C4	1.7491 (11)	C5—C6	1.4867 (15)
S1—C7	1.7957 (12)	C6—C7	1.5319 (16)
O1—C5	1.3648 (12)	C6—H6A	0.99
O1—C1	1.3719 (13)	C6—H6B	0.99
O4—C9	1.2221 (14)	C7—H7A	0.99
N1—C1	1.3340 (14)	C7—H7B	0.99
N1—H1	0.887 (17)	C10—C15	1.3869 (16)
N1—H2	0.877 (17)	C10—C11	1.3942 (16)
N2—C9	1.3480 (15)	C11—C12	1.3792 (16)
N2—C10	1.4133 (16)	C12—C13	1.3998 (16)
N2—H3	0.838 (18)	C12—H12	0.95
N3—C8	1.1543 (15)	C13—C14	1.3842 (19)
C1—C2	1.3673 (15)	C13—H13	0.95
C2—C8	1.4148 (15)	C14—C15	1.390 (2)
C2—C3	1.5228 (14)	C14—H14	0.95
C3—C4	1.4987 (14)	C15—H15	0.95
C3—C11	1.5190 (15)		
O2—S1—O3	116.71 (6)	C7—C6—H6A	110.4
O2—S1—C4	109.33 (5)	C5—C6—H6B	110.4
O3—S1—C4	111.62 (5)	C7—C6—H6B	110.4
O2—S1—C7	112.17 (6)	H6A—C6—H6B	108.6
O3—S1—C7	109.72 (5)	C6—C7—S1	107.79 (8)
C4—S1—C7	95.25 (5)	C6—C7—H7A	110.1
C5—O1—C1	117.12 (8)	S1—C7—H7A	110.1
C1—N1—H1	122.4 (11)	C6—C7—H7B	110.1
C1—N1—H2	117.5 (11)	S1—C7—H7B	110.1
H1—N1—H2	119.9 (15)	H7A—C7—H7B	108.5
C9—N2—C10	112.12 (10)	N3—C8—C2	177.90 (12)
C9—N2—H3	120.1 (11)	O4—C9—N2	126.97 (10)
C10—N2—H3	127.8 (11)	O4—C9—C3	125.10 (10)
N1—C1—C2	127.58 (10)	N2—C9—C3	107.86 (9)
N1—C1—O1	110.36 (9)	C15—C10—C11	121.93 (12)
C2—C1—O1	122.06 (9)	C15—C10—N2	128.42 (11)
C1—C2—C8	117.77 (9)	C11—C10—N2	109.66 (10)
C1—C2—C3	124.51 (10)	C12—C11—C10	120.70 (10)
C8—C2—C3	117.44 (9)	C12—C11—C3	130.77 (10)
C4—C3—C11	114.92 (9)	C10—C11—C3	108.54 (10)
C4—C3—C2	106.69 (8)	C11—C12—C13	117.97 (11)
C11—C3—C2	114.50 (9)	C11—C12—H12	121.0
C4—C3—C9	110.35 (9)	C13—C12—H12	121.0
C11—C3—C9	101.79 (8)	C14—C13—C12	120.66 (12)
C2—C3—C9	108.38 (8)	C14—C13—H13	119.7
C5—C4—C3	124.95 (9)	C12—C13—H13	119.7
C5—C4—S1	109.79 (8)	C13—C14—C15	121.88 (12)
C3—C4—S1	125.24 (8)	C13—C14—H14	119.1
C4—C5—O1	124.61 (10)	C15—C14—H14	119.1
C4—C5—C6	120.35 (10)	C10—C15—C14	116.85 (12)
O1—C5—C6	115.04 (9)	C10—C15—H15	121.6
C5—C6—C7	106.80 (9)	C14—C15—H15	121.6

supplementary materials

C5—C6—H6A	110.4		
C5—O1—C1—N1	-179.52 (9)	C5—C6—C7—S1	-1.03 (11)
C5—O1—C1—C2	0.99 (15)	O2—S1—C7—C6	-112.07 (8)
N1—C1—C2—C8	-8.53 (18)	O3—S1—C7—C6	116.50 (8)
O1—C1—C2—C8	170.86 (9)	C4—S1—C7—C6	1.28 (9)
N1—C1—C2—C3	177.78 (10)	C10—N2—C9—O4	178.17 (11)
O1—C1—C2—C3	-2.82 (17)	C10—N2—C9—C3	0.86 (12)
C1—C2—C3—C4	2.39 (14)	C4—C3—C9—O4	58.53 (13)
C8—C2—C3—C4	-171.31 (9)	C11—C3—C9—O4	-179.02 (10)
C1—C2—C3—C11	-125.94 (11)	C2—C3—C9—O4	-57.96 (14)
C8—C2—C3—C11	60.36 (13)	C4—C3—C9—N2	-124.10 (10)
C1—C2—C3—C9	121.23 (11)	C11—C3—C9—N2	-1.65 (11)
C8—C2—C3—C9	-52.48 (12)	C2—C3—C9—N2	119.41 (10)
C11—C3—C4—C5	127.60 (11)	C9—N2—C10—C15	-179.69 (12)
C2—C3—C4—C5	-0.48 (15)	C9—N2—C10—C11	0.40 (13)
C9—C3—C4—C5	-118.03 (11)	C15—C10—C11—C12	-1.83 (17)
C11—C3—C4—S1	-54.22 (13)	N2—C10—C11—C12	178.09 (10)
C2—C3—C4—S1	177.70 (7)	C15—C10—C11—C3	178.57 (11)
C9—C3—C4—S1	60.16 (12)	N2—C10—C11—C3	-1.51 (12)
O2—S1—C4—C5	114.52 (9)	C4—C3—C11—C12	-58.40 (15)
O3—S1—C4—C5	-114.83 (8)	C2—C3—C11—C12	65.64 (15)
C7—S1—C4—C5	-1.19 (9)	C9—C3—C11—C12	-177.67 (11)
O2—S1—C4—C3	-63.89 (11)	C4—C3—C11—C10	121.14 (10)
O3—S1—C4—C3	66.76 (10)	C2—C3—C11—C10	-114.81 (10)
C7—S1—C4—C3	-179.61 (10)	C9—C3—C11—C10	1.88 (11)
C3—C4—C5—O1	-1.14 (18)	C10—C11—C12—C13	1.27 (16)
S1—C4—C5—O1	-179.56 (8)	C3—C11—C12—C13	-179.23 (11)
C3—C4—C5—C6	179.18 (10)	C11—C12—C13—C14	-0.03 (17)
S1—C4—C5—C6	0.76 (13)	C12—C13—C14—C15	-0.73 (19)
C1—O1—C5—C4	0.96 (15)	C11—C10—C15—C14	1.04 (18)
C1—O1—C5—C6	-179.33 (9)	N2—C10—C15—C14	-178.86 (12)
C4—C5—C6—C7	0.21 (14)	C13—C14—C15—C10	0.22 (19)
O1—C5—C6—C7	-179.51 (9)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots N3 ⁱ	0.89 (2)	2.21 (2)	3.0878 (15)	170 (2)
N1—H2 \cdots O4 ⁱⁱ	0.88 (2)	2.01 (2)	2.8790 (14)	172 (1)
N2—H3 \cdots O3 ⁱⁱⁱ	0.84 (2)	2.13 (2)	2.9255 (14)	159 (2)

Symmetry codes: (i) $-x, -y+2, -z$; (ii) $-x, -y+1, -z$; (iii) $x, -y+1, z-1/2$.

Fig. 1




Fig. 2

