

## **rac-(E)-3-[1-(2-Chlorophenyl)ethyl]-5-methyl-N-nitro-1,3,5-oxadiazinan-4-imine**

Cong-Cong Li, Yuan-Yuan Zhong and Liang-Zhong Xu\*

College of Chemistry and Molecular Engineering, Qingdao University of Science and Technology, Qingdao 266042, People's Republic of China  
Correspondence e-mail: qknhs@yahoo.com.cn


Received 19 August 2010; accepted 25 August 2010

Key indicators: single-crystal X-ray study;  $T = 113\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.039;  $wR$  factor = 0.104; data-to-parameter ratio = 14.1.

In the title compound,  $\text{C}_{12}\text{H}_{15}\text{ClN}_4\text{O}_3$ , which has potential insecticidal activity, the oxadiazine ring and the benzene ring make a dihedral angle of  $84.63(2)^\circ$  to one another. The crystal packing involves weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds.

### Related literature

For the biological activity of oxadiazine derivatives, see: Maienfisch & Huerlimann (1994); Gsell & Maienfisch (1998). For the synthesis, see: Gottfied *et al.* (2001). For related structures, see: Chopra *et al.* (2004); Kang *et al.* (2008); Zhong *et al.* (2010). For puckering parameters, see: Cremer & Pople (1975).



### Experimental

#### Crystal data

$\text{C}_{12}\text{H}_{15}\text{ClN}_4\text{O}_3$	$V = 1368.0(5)\text{ \AA}^3$
$M_r = 298.73$	$Z = 4$
Monoclinic, $P2_1/c$	$\text{Cu } K\alpha$ radiation
$a = 17.259(4)\text{ \AA}$	$\mu = 2.61\text{ mm}^{-1}$
$b = 6.9157(14)\text{ \AA}$	$T = 113\text{ K}$
$c = 12.169(2)\text{ \AA}$	$0.26 \times 0.22 \times 0.18\text{ mm}$
$\beta = 109.63(3)^\circ$	

#### Data collection

Rigaku Saturn diffractometer	12087 measured reflections
Absorption correction: multi-scan ( <i>CrystalClear</i> ; Rigaku, 2005)	2590 independent reflections
$T_{\min} = 0.550$ , $T_{\max} = 0.651$	2562 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.054$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	184 parameters
$wR(F^2) = 0.104$	H-atom parameters constrained
$S = 1.08$	$\Delta\rho_{\max} = 0.26\text{ e \AA}^{-3}$
2590 reflections	$\Delta\rho_{\min} = -0.37\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C1}-\text{H1A}\cdots\text{O2}^{\text{i}}$	0.99	2.52	3.2817 (18)	134
$\text{C1}-\text{H1A}\cdots\text{O3}^{\text{i}}$	0.99	2.50	3.396 (2)	150
$\text{C1}-\text{H1B}\cdots\text{O2}^{\text{ii}}$	0.99	2.56	3.2555 (18)	127
$\text{C3}-\text{H3A}\cdots\text{O3}^{\text{iii}}$	0.99	2.49	3.2294 (19)	131
$\text{C4}-\text{H4B}\cdots\text{O2}^{\text{j}}$	0.98	2.57	3.3070 (19)	132
$\text{C4}-\text{H4C}\cdots\text{O1}^{\text{ii}}$	0.98	2.49	3.377 (2)	150
$\text{C6}-\text{H6C}\cdots\text{O3}^{\text{iii}}$	0.98	2.35	3.3020 (19)	164

Symmetry codes: (i)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (ii)  $-x, -y + 1, -z$ ; (iii)  $x, y + 1, z$ .

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2061).

### References

- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.  
Chopra, D., Mohan, T. P., Rao, K. S. & Guru Row, T. N. (2004). *Acta Cryst. E60*, o2413–o2414.  
Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.  
Gottfied, S., Thomas, R. & Verena, G. (2001). WO Patent 0100623.  
Gsell, L. & Maienfisch, P. (1998). WO Patent 9806710.  
Kang, T.-N., Zhang, L., Ling, Y. & Yang, X.-L. (2008). *Acta Cryst. E64*, o1154.  
Maienfisch, P. & Huerlimann, H. (1994). CN Patent 1084171.  
Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.  
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.  
Zhong, Y., Li, C. & Xu, L. (2010). *Acta Cryst. E66*, o1981.

## **supplementary materials**

Acta Cryst. (2010). E66, o2456 [doi:10.1107/S1600536810034343]

### **rac-(E)-3-[1-(2-Chlorophenyl)ethyl]-5-methyl-N-nitro-1,3,5-oxadiazinan-4-imine**

**C.-C. Li, Y.-Y. Zhong and L.-Z. Xu**

#### **Comment**

Currently, studies on oxadiazine derivatives have mainly concentrated on compounds with oxadiazine as the only active group (Gsell, *et al.*, 1998) and a number of highly insecticidal compounds of this type have been synthesized (Maienfisch *et al.*, 1994). We report here the synthesis and crystal structure of the title compound C<sub>12</sub>H<sub>15</sub>Cl<sub>1</sub>N<sub>4</sub>O<sub>3</sub> (I).

In (I) (Fig. 1) the bond lengths and angles of the oxadiazine rings are in agreement with those in previous reported structures (Chopra *et al.*, 2004). The 1,3,5-oxadiazinane ring is in a half-chair conformation and the ring-puckering parameters (Cremer & Pople, 1975;) were calculated as Q = 0.05126 (12) Å; θ = 121.33 (13)°; φ = 166.3676 (15)°. The N3=O bondlength [1.3904 (17) Å] is close to the value reported in the literature (Zhong *et al.*, 2010). The oxadiazine ring and the benzene ring make a dihedral angle of 84.63 (2)°. Weak intermolecular C—H···O hydrogen bonds give a three-dimensional network (Table 1).

#### **Experimental**

A solution of 1-(1-bromoethyl)-2-chlorobenzene (4.3 g, 20 mmol), *N*-nitro-1,3,5-oxadiazinan-4-imine (3.2 g, 20 mmol) and potassium carbonate (2.8 g, 20 mmol) in 20 g of acetonitrile was heated under reflux for 4 h. Upon cooling to room temperature the solution was filtered and then concentrated under reduced pressure to give the title compound (I) (7.89 g, 90% yield) (Gottfried, *et al.*, 2001). Single crystals suitable for X-ray measurement were obtained by recrystallization from ethanol at room temperature.

#### **Refinement**

All C-bound H atoms were placed in calculated positions, with C—H = 0.95–0.97 Å, and included in the final cycles of refinement using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

#### **Figures**




Fig. 1. Molecular configuration and atom numbering scheme for the title compound (I), with displacement ellipsoids drawn at the 50% probability level.

# supplementary materials

---

## ***rac-(E)-3-[1-(2-Chlorophenyl)ethyl]-5-methyl-N- nitro-1,3,5-oxadiazinan-4-imine***

### *Crystal data*

C <sub>12</sub> H <sub>15</sub> ClN <sub>4</sub> O <sub>3</sub>	F(000) = 624
M <sub>r</sub> = 298.73	D <sub>x</sub> = 1.450 Mg m <sup>-3</sup>
Monoclinic, P2 <sub>1</sub> /c	Cu K $\alpha$ radiation, $\lambda$ = 1.54187 Å
Hall symbol: -P 2ybc	Cell parameters from 1058 reflections
a = 17.259 (4) Å	$\theta$ = 27.5–71.9°
b = 6.9157 (14) Å	$\mu$ = 2.61 mm <sup>-1</sup>
c = 12.169 (2) Å	T = 113 K
$\beta$ = 109.63 (3)°	Block, colorless
V = 1368.0 (5) Å <sup>3</sup>	0.26 × 0.22 × 0.18 mm
Z = 4	

### *Data collection*

Rigaku Saturn diffractometer	2590 independent reflections
Radiation source: fine-focus sealed tube graphite	2562 reflections with $I > 2\sigma(I)$
Detector resolution: 14.63 pixels mm <sup>-1</sup>	$R_{\text{int}} = 0.054$
$\omega$ scans	$\theta_{\text{max}} = 72.1^\circ$ , $\theta_{\text{min}} = 2.7^\circ$
Absorption correction: multi-scan ( <i>CrystalClear</i> ; Rigaku, 2005)	$h = -20 \rightarrow 21$
$T_{\text{min}} = 0.550$ , $T_{\text{max}} = 0.651$	$k = -8 \rightarrow 8$
12087 measured reflections	$l = -13 \rightarrow 14$

### *Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.039$	H-atom parameters constrained
$wR(F^2) = 0.104$	$w = 1/[\sigma^2(F_o^2) + (0.058P)^2 + 0.6637P]$
$S = 1.08$	where $P = (F_o^2 + 2F_c^2)/3$
2590 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
184 parameters	$\Delta\rho_{\text{max}} = 0.26 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.37 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{1/4}$
	Extinction coefficient: 0.0057 (11)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.36163 (2)	0.37729 (6)	-0.00540 (3)	0.02744 (17)
N1	0.09670 (7)	0.33360 (16)	-0.04098 (10)	0.0152 (3)
N2	0.21260 (7)	0.48468 (16)	0.08081 (10)	0.0146 (3)
N3	0.19653 (7)	0.15435 (16)	0.10592 (10)	0.0165 (3)
N4	0.15382 (7)	0.09523 (17)	0.17188 (10)	0.0172 (3)
O1	0.09734 (6)	0.67120 (14)	-0.01841 (9)	0.0180 (2)
O2	0.09602 (7)	0.19372 (16)	0.18398 (9)	0.0247 (3)
O3	0.17457 (7)	-0.06237 (15)	0.22426 (9)	0.0238 (3)
C1	0.06227 (8)	0.5189 (2)	-0.09525 (12)	0.0171 (3)
H1A	0.0740	0.5363	-0.1689	0.021*
H1B	0.0018	0.5188	-0.1139	0.021*
C2	0.16607 (8)	0.32676 (19)	0.04897 (12)	0.0138 (3)
C3	0.18404 (8)	0.66306 (19)	0.01450 (13)	0.0172 (3)
H3A	0.2089	0.7767	0.0630	0.021*
H3B	0.2007	0.6644	-0.0558	0.021*
C4	0.04698 (9)	0.1625 (2)	-0.08861 (13)	0.0196 (3)
H4A	0.0801	0.0459	-0.0611	0.029*
H4B	0.0285	0.1670	-0.1740	0.029*
H4C	-0.0010	0.1599	-0.0626	0.029*
C5	0.28920 (8)	0.48416 (19)	0.18389 (12)	0.0171 (3)
H5	0.3141	0.3524	0.1894	0.020*
C6	0.26984 (10)	0.5185 (2)	0.29596 (12)	0.0225 (3)
H6A	0.3212	0.5198	0.3627	0.034*
H6B	0.2342	0.4147	0.3061	0.034*
H6C	0.2418	0.6430	0.2910	0.034*
C7	0.34952 (8)	0.6269 (2)	0.16156 (13)	0.0184 (3)
C8	0.38573 (9)	0.5900 (2)	0.07689 (13)	0.0209 (3)
C9	0.44068 (9)	0.7161 (2)	0.05401 (14)	0.0257 (4)
H9	0.4640	0.6869	-0.0045	0.031*
C10	0.46123 (10)	0.8863 (2)	0.11817 (16)	0.0305 (4)
H10	0.4997	0.9733	0.1048	0.037*
C11	0.42544 (10)	0.9287 (2)	0.20160 (15)	0.0308 (4)
H11	0.4389	1.0459	0.2445	0.037*

## supplementary materials

---

C12	0.37004 (9)	0.8007 (2)	0.22283 (13)	0.0247 (3)
H12	0.3457	0.8320	0.2800	0.030*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0282 (2)	0.0206 (2)	0.0376 (3)	-0.00083 (13)	0.01641 (18)	-0.00617 (14)
N1	0.0166 (6)	0.0106 (5)	0.0167 (6)	-0.0008 (4)	0.0035 (4)	-0.0001 (4)
N2	0.0146 (5)	0.0095 (5)	0.0179 (6)	0.0011 (4)	0.0029 (4)	0.0012 (4)
N3	0.0171 (6)	0.0104 (5)	0.0226 (6)	0.0015 (4)	0.0075 (5)	0.0038 (4)
N4	0.0216 (6)	0.0110 (5)	0.0182 (6)	0.0015 (5)	0.0058 (5)	0.0019 (4)
O1	0.0167 (5)	0.0119 (5)	0.0225 (5)	0.0038 (4)	0.0030 (4)	-0.0010 (4)
O2	0.0286 (6)	0.0209 (6)	0.0301 (6)	0.0107 (4)	0.0170 (5)	0.0072 (4)
O3	0.0338 (6)	0.0117 (5)	0.0275 (6)	0.0059 (4)	0.0123 (4)	0.0083 (4)
C1	0.0168 (6)	0.0134 (6)	0.0183 (7)	0.0022 (5)	0.0021 (5)	0.0009 (5)
C2	0.0156 (6)	0.0100 (6)	0.0173 (6)	0.0016 (5)	0.0077 (5)	-0.0003 (5)
C3	0.0165 (7)	0.0099 (6)	0.0226 (7)	0.0010 (5)	0.0032 (5)	0.0035 (5)
C4	0.0204 (7)	0.0144 (6)	0.0222 (7)	-0.0049 (5)	0.0048 (6)	-0.0024 (5)
C5	0.0163 (6)	0.0123 (6)	0.0190 (7)	0.0019 (5)	0.0011 (5)	0.0014 (5)
C6	0.0274 (7)	0.0183 (7)	0.0192 (7)	0.0004 (6)	0.0042 (6)	0.0019 (5)
C7	0.0149 (6)	0.0136 (7)	0.0216 (7)	0.0010 (5)	-0.0006 (5)	0.0026 (5)
C8	0.0157 (6)	0.0157 (7)	0.0270 (8)	0.0018 (5)	0.0015 (6)	0.0030 (6)
C9	0.0182 (7)	0.0267 (8)	0.0298 (8)	0.0013 (6)	0.0047 (6)	0.0093 (6)
C10	0.0211 (8)	0.0248 (8)	0.0372 (9)	-0.0083 (6)	-0.0013 (7)	0.0108 (7)
C11	0.0302 (8)	0.0187 (7)	0.0325 (9)	-0.0086 (7)	-0.0042 (7)	0.0003 (6)
C12	0.0249 (7)	0.0185 (7)	0.0242 (7)	-0.0029 (6)	-0.0004 (6)	-0.0013 (6)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Cl1—C8	1.7492 (16)	C4—H4B	0.9800
N1—C2	1.324 (2)	C4—H4C	0.9800
N1—C4	1.4621 (17)	C5—C7	1.524 (2)
N1—C1	1.4721 (17)	C5—C6	1.528 (2)
N2—C2	1.3336 (18)	C5—H5	1.0000
N2—C3	1.4658 (16)	C6—H6A	0.9800
N2—C5	1.4856 (17)	C6—H6B	0.9800
N3—N4	1.3237 (17)	C6—H6C	0.9800
N3—C2	1.3904 (17)	C7—C12	1.396 (2)
N4—O3	1.2526 (15)	C7—C8	1.396 (2)
N4—O2	1.2567 (16)	C8—C9	1.384 (2)
O1—C1	1.4037 (17)	C9—C10	1.391 (2)
O1—C3	1.4143 (18)	C9—H9	0.9500
C1—H1A	0.9900	C10—C11	1.386 (3)
C1—H1B	0.9900	C10—H10	0.9500
C3—H3A	0.9900	C11—C12	1.389 (2)
C3—H3B	0.9900	C11—H11	0.9500
C4—H4A	0.9800	C12—H12	0.9500
C2—N1—C4	123.09 (11)	H4B—C4—H4C	109.5

C2—N1—C1	121.23 (11)	N2—C5—C7	108.38 (11)
C4—N1—C1	115.58 (11)	N2—C5—C6	110.73 (11)
C2—N2—C3	118.10 (11)	C7—C5—C6	115.21 (12)
C2—N2—C5	121.43 (11)	N2—C5—H5	107.4
C3—N2—C5	120.40 (11)	C7—C5—H5	107.4
N4—N3—C2	111.90 (11)	C6—C5—H5	107.4
O3—N4—O2	120.96 (12)	C5—C6—H6A	109.5
O3—N4—N3	117.07 (11)	C5—C6—H6B	109.5
O2—N4—N3	121.95 (11)	H6A—C6—H6B	109.5
C1—O1—C3	109.70 (10)	C5—C6—H6C	109.5
O1—C1—N1	109.62 (11)	H6A—C6—H6C	109.5
O1—C1—H1A	109.7	H6B—C6—H6C	109.5
N1—C1—H1A	109.7	C12—C7—C8	117.08 (14)
O1—C1—H1B	109.7	C12—C7—C5	121.90 (14)
N1—C1—H1B	109.7	C8—C7—C5	121.02 (13)
H1A—C1—H1B	108.2	C9—C8—C7	122.70 (14)
N1—C2—N2	119.99 (12)	C9—C8—Cl1	117.52 (13)
N1—C2—N3	121.87 (12)	C7—C8—Cl1	119.78 (11)
N2—C2—N3	117.88 (12)	C8—C9—C10	118.89 (16)
O1—C3—N2	108.51 (11)	C8—C9—H9	120.6
O1—C3—H3A	110.0	C10—C9—H9	120.6
N2—C3—H3A	110.0	C11—C10—C9	119.87 (15)
O1—C3—H3B	110.0	C11—C10—H10	120.1
N2—C3—H3B	110.0	C9—C10—H10	120.1
H3A—C3—H3B	108.4	C10—C11—C12	120.38 (15)
N1—C4—H4A	109.5	C10—C11—H11	119.8
N1—C4—H4B	109.5	C12—C11—H11	119.8
H4A—C4—H4B	109.5	C11—C12—C7	121.07 (16)
N1—C4—H4C	109.5	C11—C12—H12	119.5
H4A—C4—H4C	109.5	C7—C12—H12	119.5
C2—N3—N4—O3	177.78 (11)	C3—N2—C5—C7	-33.89 (16)
C2—N3—N4—O2	-3.68 (18)	C2—N2—C5—C6	-83.61 (15)
C3—O1—C1—N1	55.36 (14)	C3—N2—C5—C6	93.38 (14)
C2—N1—C1—O1	-18.08 (18)	N2—C5—C7—C12	110.06 (14)
C4—N1—C1—O1	158.55 (12)	C6—C5—C7—C12	-14.60 (19)
C4—N1—C2—N2	173.37 (12)	N2—C5—C7—C8	-68.88 (16)
C1—N1—C2—N2	-10.3 (2)	C6—C5—C7—C8	166.46 (13)
C4—N1—C2—N3	-0.6 (2)	C12—C7—C8—C9	0.9 (2)
C1—N1—C2—N3	175.73 (12)	C5—C7—C8—C9	179.93 (13)
C3—N2—C2—N1	0.66 (19)	C12—C7—C8—Cl1	-178.83 (10)
C5—N2—C2—N1	177.72 (12)	C5—C7—C8—Cl1	0.15 (18)
C3—N2—C2—N3	174.91 (11)	C7—C8—C9—C10	0.4 (2)
C5—N2—C2—N3	-8.03 (19)	Cl1—C8—C9—C10	-179.86 (11)
N4—N3—C2—N1	-73.97 (16)	C8—C9—C10—C11	-1.3 (2)
N4—N3—C2—N2	111.90 (14)	C9—C10—C11—C12	1.0 (2)
C1—O1—C3—N2	-64.67 (14)	C10—C11—C12—C7	0.4 (2)
C2—N2—C3—O1	36.13 (16)	C8—C7—C12—C11	-1.3 (2)
C5—N2—C3—O1	-140.96 (12)	C5—C7—C12—C11	179.72 (13)
C2—N2—C5—C7	149.11 (13)		

## supplementary materials

---

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C1—H1A···O2 <sup>i</sup>	0.99	2.52	3.2817 (18)	134.
C1—H1A···O3 <sup>i</sup>	0.99	2.50	3.396 (2)	150.
C1—H1B···O2 <sup>ii</sup>	0.99	2.56	3.2555 (18)	127.
C3—H3A···O3 <sup>iii</sup>	0.99	2.49	3.2294 (19)	131.
C4—H4B···O2 <sup>i</sup>	0.98	2.57	3.3070 (19)	132.
C4—H4C···O1 <sup>ii</sup>	0.98	2.49	3.377 (2)	150.
C6—H6C···O3 <sup>iii</sup>	0.98	2.35	3.3020 (19)	164.

Symmetry codes: (i)  $x, -y+1/2, z-1/2$ ; (ii)  $-x, -y+1, -z$ ; (iii)  $x, y+1, z$ .

Fig. 1

