

Acetato(*N*-[(*E*)-1-(6-methyl-2-pyridyl)methylidene]-2-[2-[(*E*)-1-(6-methyl-2-pyridyl)methylideneamino]phenethyl]-aniline)nickel(II) perchlorate

Shuranjan Sarkar,^a Hajin Lee^b and Hong-In Lee^{a*}

^aDepartment of Chemistry, Kyungpook National University, Daegu, 702-701, Republic of Korea, and ^bKorea Basic Science Institute, 664-14 Dukjin dong 1-ga, Dukjin-gu, Jeonju, 561-756, Republic of Korea
Correspondence e-mail: lee@knu.ac.kr


Received 17 August 2010; accepted 26 August 2010

Key indicators: single-crystal X-ray study; *T* = 200 K; mean $\sigma(\text{C}-\text{C})$ = 0.008 Å; *R* factor = 0.068; *wR* factor = 0.149; data-to-parameter ratio = 17.9.

In the title complex, $[\text{Ni}(\text{CH}_3\text{COO})(\text{C}_{28}\text{H}_{26}\text{N}_4)]\text{ClO}_4$, the Ni^{II} atom is coordinated by two imine N atoms and two pyridine N atoms of the *N*-[(*E*)-1-(6-methyl-2-pyridyl)methylidene]-2-(2-[(*E*)-1-(6-methyl-2-pyridyl)methylideneamino]phenethyl)-aniline donor ligand and two O atoms of the acetate ion in a distorted octahedral coordination. The average Ni—N and Ni—O bond lengths are 2.131 (13) and 2.098 (11) Å, respectively. An intramolecular N—H···O interaction occurs. Relatively weak intermolecular C—H···O interactions between the ligands and the ClO₄[−] ions result in a chain extending along the *b* axis.

Related literature

For structures of Ni complexes with ligands formed by the condensation of 2-pyridyl aldehydes and a variety of diamines, see: Banerjee *et al.* (2004). For comparison Ni—N bond distances, see: Martin *et al.* (1977).



Experimental

Crystal data

$[\text{Ni}(\text{C}_2\text{H}_3\text{O}_2)(\text{C}_{28}\text{H}_{26}\text{N}_4)]\text{ClO}_4$
M_r = 635.73
Triclinic, *P* $\bar{1}$
a = 8.5759 (8) Å
b = 11.4975 (10) Å
c = 14.8322 (13) Å
 α = 79.392 (2)°
 β = 78.102 (2)°
 γ = 81.327 (2)°
V = 1396.9 (2) Å³
Z = 2
Mo *K*α radiation
 μ = 0.84 mm^{−1}
T = 200 K
0.23 × 0.11 × 0.10 mm

Data collection

Bruker APEX CCD area-detector diffractometer
10545 measured reflections
6856 independent reflections
3311 reflections with *I* > 2σ(*I*)
R_{int} = 0.049

Refinement

$R[F^2 > 2\sigma(F^2)]$ = 0.068
 $wR(F^2)$ = 0.149
S = 1.05
6856 reflections
382 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}}$ = 1.04 e Å^{−3}
 $\Delta\rho_{\text{min}}$ = −1.84 e Å^{−3}

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C4—H4···O1 ⁱ	0.95	2.45	3.274 (6)	144
C7—H7···O4 ⁱⁱ	0.95	2.41	3.290 (7)	155
C20—H20···O3 ⁱⁱⁱ	0.95	2.43	3.366 (8)	168
C15—H15B···O2	0.99	2.54	3.523 (7)	174
C15—H15B···N3	0.99	2.47	2.935 (7)	108

Symmetry codes: (i) $-x + 2, -y + 2, -z + 1$; (ii) $x + 1, y + 1, z$; (iii) $x + 1, y, z$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by a Korea Research Foundation Grant funded by the Korean Government (KRF-2008-313-C00433).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2321).

References

Banerjee, S., Gangopadhyay, J., Lu, C.-Z., Chen, J.-T. & Ghosh, A. (2004). *Eur. J. Inorg. Chem.* pp. 2533–2541.
Bruker (2000). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Martin, L. Y., Sperati, C. R. & Busch, D. H. (1977). *J. Am. Chem. Soc.* **99**, 2968–2981.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, m1184 [doi:10.1107/S1600536810034446]

Acetato(*N*-[(*E*)-1-(6-methyl-2-pyridyl)methylidene]-2-{2-[(*E*)-1-(6-methyl-2-pyridyl)methylideneamino]phenethyl}aniline)nickel(II) perchlorate

S. Sarkar, H. Lee and H.-I. Lee

Comment

The coordination chemistry of Schiff-base ligands formed by the condensation of 2-pyridyl aldehydes with a variety of diamines has been reported recently. These ligands bind tetradentately to metal ions to form a planar arrangement around the metals (Banerjee *et al.*, 2004). We are quite interested in the synthesis of this type of Schiff-base tetraamine nickel(II) complexes and have obtained a novel nickel(II) compound with an acetate group. In this paper, we report the synthesis and crystal structure of the title complex.

The title complex consists of $[\text{Ni}(\text{mpma})(\eta^2\text{-CH}_3\text{CO}_2)]^+$ (mpma = *N*1-[(*E*)-1-(6-methyl-2-pyridyl)-methylidene]-2-(2-[(*E*)-1-(6-methyl-2-pyridyl)methylidene]aminophenethyl)aniline) cation and ClO_4^- anion (Fig. 1) wherein the Nickel(II) ion is six-coordinated with four N atoms of mpma and two O atoms of the acetate group, giving a distorted octahedral geometry. The average Ni—N and Ni—O bond lengths are 2.131 (13) and 2.098 (11) Å, respectively. It is in good agreement with the general trend that the nickel(II)-nitrogen bonds are longer (or weaker) in the octahedral species than in square planar species (Ni—N = 1.88–1.91 Å) (Martin *et al.*, 1977). The N1—Ni1—N2, N3—Ni1—N4, and O1—Ni1—O2 bond angles are 79.57 (16), 79.87 (18), and 63.16 (14)°, respectively. The deviation of these angles from the ideal octahedral geometry is due to the constraints of the five membered chelate rings (N1—Ni1—N2—C7—C6 and N3—Ni1—N4—N23—N22) and four membered chelate ring (O1—Ni1—O2—C29). The distortion is also reflected in the bond angles N2—Ni1—N4, O2—Ni1—N1, and O1—Ni1—N3, which are 171.81 (19), 164.65 (15), and 158.19 (15)°, respectively. Relatively weak, intermolecular, C—H···O distances are found between the ligand and ClO_4^- ion, forming an one-dimensional chain extended along the *b*-axis (Fig. 2 and Tab. 1).

Experimental

Nickel(II)acetate tetrahydrate (0.5 g, 2.0 mmol) dissolved in dry methanol (25 ml) was added dropwise to a methanol solution (10 ml) of mpma (0.84 g, 2.0 mmol) and stirred. A green color solution appeared. Then methanol solution (5 ml) of sodium perchlorate (0.25 g, 2.0 mmol) was added. After 1 h, a crystalline powder (1.15 g) was collected by filtration and dried in vacuum. The powder (*ca.* 0.7 g) was dissolved in dry methanol (5 ml) and then diethyl ether (5 ml) was added slowly into the methanol solution. Suitable crystals for X-ray analysis were obtained from the solution after one day.

Refinement

All H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with distances C—H = 0.95, 0.98 and 0.99 Å for aryl, methyl and methylene H-atoms and $U_{\text{iso}}(\text{H}) = 1.5$ (methyl) and 1.2 (the rest) \times the U_{eq} of the parent C-atoms. A relatively large residual density on Ni ion is a ghost peak residing less than 1 Å from the ion. An absorption correction did not improve the refinement.

Figures




Fig. 1. Structure of the title compound showing 33% probability displacement ellipsoids and the atom-numbering scheme; H atoms have been omitted for clarity.




Fig. 2. One-dimensional chain of the title complex formed by intermolecular interactions (dotted lines); H atoms have been omitted except for those involved in the C—H...O interactions.

Acetato(*N*-[*(E)*-1-(6-methyl-2-pyridyl)methylidene]-2-{2-[*(E)*-1-(6-methyl-2-pyridyl)methylideneamino]phenethyl}aniline)nickel(II) perchlorate

Crystal data

[Ni(C₂H₃O₂)(C₂₈H₂₆N₄)]ClO₄

M_r = 635.73

Triclinic, *PT*

Hall symbol: -P 1

a = 8.5759 (8) Å

b = 11.4975 (10) Å

c = 14.8322 (13) Å

α = 79.392 (2)°

β = 78.102 (2)°

γ = 81.327 (2)°

V = 1396.9 (2) Å³

Z = 2

F(000) = 660

D_x = 1.511 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 2599 reflections

θ = 2.4–26.0°

μ = 0.84 mm⁻¹

T = 200 K

Block, yellow

0.23 × 0.11 × 0.10 mm

Data collection

Bruker APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

graphite

φ & ω scans

10545 measured reflections

6856 independent reflections

3311 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.049

θ_{max} = 28.3°, θ_{min} = 1.4°

h = -9→11

k = -15→15

l = -18→19

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.068

wR(*F*²) = 0.149

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + 2.2621P]$
6856 reflections	where $P = (F_o^2 + 2F_c^2)/3$
382 parameters	$(\Delta/\sigma)_{\max} < 0.001$
0 restraints	$\Delta\rho_{\max} = 1.04 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -1.84 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.81516 (9)	0.76280 (6)	0.72685 (5)	0.02685 (19)
N1	1.0115 (5)	0.8119 (4)	0.6208 (3)	0.0245 (10)
C1	1.1033 (7)	0.7543 (5)	0.5523 (4)	0.0314 (14)
C2	1.0751 (7)	0.6312 (5)	0.5466 (4)	0.0404 (16)
H2A	1.0736	0.5815	0.6080	0.061*
H2B	1.1614	0.5975	0.5010	0.061*
H2C	0.9719	0.6338	0.5273	0.061*
C3	1.2207 (7)	0.8104 (5)	0.4861 (4)	0.0382 (16)
H3	1.2797	0.7698	0.4372	0.046*
C4	1.2535 (7)	0.9230 (5)	0.4898 (4)	0.0342 (15)
H4	1.3343	0.9602	0.4447	0.041*
C5	1.1637 (7)	0.9801 (5)	0.5622 (4)	0.0351 (15)
H5	1.1824	1.0574	0.5681	0.042*
C6	1.0472 (7)	0.9219 (5)	0.6249 (4)	0.0283 (13)
C7	0.9486 (7)	0.9817 (5)	0.7008 (4)	0.0289 (13)
H7	0.9575	1.0627	0.7020	0.035*
N2	0.8519 (5)	0.9256 (4)	0.7645 (3)	0.0264 (11)
C8	0.7531 (7)	0.9948 (5)	0.8311 (4)	0.0294 (14)
C9	0.6261 (7)	1.0765 (4)	0.8044 (4)	0.0329 (15)
H9	0.6029	1.0833	0.7436	0.039*
C10	0.5334 (7)	1.1478 (5)	0.8670 (4)	0.0353 (15)
H10	0.4468	1.2029	0.8490	0.042*
C11	0.5679 (7)	1.1381 (5)	0.9551 (4)	0.0373 (16)
H11	0.5057	1.1872	0.9976	0.045*
C12	0.6942 (7)	1.0560 (5)	0.9817 (4)	0.0364 (15)
H12	0.7180	1.0502	1.0423	0.044*
C13	0.7850 (7)	0.9829 (4)	0.9209 (4)	0.0275 (13)

supplementary materials

C14	0.9193 (7)	0.8913 (5)	0.9504 (4)	0.0347 (15)
H14A	0.9391	0.9050	1.0109	0.042*
H14B	1.0188	0.9024	0.9037	0.042*
C15	0.8821 (7)	0.7607 (4)	0.9605 (4)	0.0290 (13)
H15A	0.8297	0.7337	1.0254	0.035*
H15B	0.8075	0.7567	0.9186	0.035*
C16	1.0336 (7)	0.6814 (4)	0.9361 (4)	0.0252 (12)
C17	1.1568 (7)	0.6698 (5)	0.9882 (4)	0.0308 (14)
H17	1.1402	0.7108	1.0401	0.037*
C18	1.2997 (7)	0.6012 (5)	0.9660 (4)	0.0393 (15)
H18	1.3804	0.5954	1.0024	0.047*
C19	1.3277 (8)	0.5402 (5)	0.8913 (4)	0.0418 (16)
H19	1.4274	0.4930	0.8759	0.050*
C20	1.2097 (7)	0.5483 (5)	0.8389 (4)	0.0360 (15)
H20	1.2283	0.5062	0.7876	0.043*
C21	1.0633 (7)	0.6177 (5)	0.8609 (4)	0.0272 (13)
N3	0.9426 (5)	0.6226 (4)	0.8063 (3)	0.0253 (11)
C22	0.9007 (7)	0.5194 (5)	0.8015 (4)	0.0308 (14)
H22	0.9467	0.4488	0.8354	0.037*
C23	0.7849 (7)	0.5120 (4)	0.7447 (4)	0.0264 (13)
C24	0.7234 (7)	0.4057 (5)	0.7518 (4)	0.0359 (15)
H24	0.7571	0.3367	0.7924	0.043*
C25	0.6106 (7)	0.4033 (5)	0.6974 (4)	0.0387 (16)
H25	0.5607	0.3333	0.7032	0.046*
C26	0.5713 (7)	0.5021 (5)	0.6354 (4)	0.0391 (16)
H26	0.4949	0.5008	0.5975	0.047*
C27	0.6460 (7)	0.6062 (5)	0.6284 (4)	0.0296 (13)
C28	0.6155 (8)	0.7094 (5)	0.5526 (4)	0.0491 (18)
H28A	0.5150	0.7579	0.5738	0.074*
H28B	0.6074	0.6790	0.4964	0.074*
H28C	0.7043	0.7584	0.5384	0.074*
N4	0.7454 (5)	0.6115 (4)	0.6852 (3)	0.0267 (11)
O1	0.6159 (5)	0.8708 (3)	0.6811 (3)	0.0337 (10)
O2	0.5968 (4)	0.7613 (3)	0.8201 (2)	0.0291 (9)
C29	0.5314 (7)	0.8311 (5)	0.7580 (4)	0.0317 (14)
C30	0.3545 (7)	0.8655 (5)	0.7740 (4)	0.0455 (17)
H30A	0.3315	0.9522	0.7700	0.068*
H30B	0.3063	0.8271	0.8361	0.068*
H30C	0.3092	0.8397	0.7265	0.068*
Cl1	0.1830 (2)	0.26066 (14)	0.70157 (11)	0.0417 (4)
O3	0.2297 (6)	0.3780 (4)	0.6741 (3)	0.0703 (16)
O4	0.0681 (6)	0.2487 (4)	0.6474 (3)	0.0619 (14)
O5	0.3214 (6)	0.1768 (4)	0.6830 (3)	0.0754 (17)
O6	0.1128 (7)	0.2433 (5)	0.7980 (3)	0.0837 (19)

Atomic displacement parameters (\AA^2)

U^{11}

U^{22}

U^{33}

U^{12}

U^{13}

U^{23}

Ni1	0.0292 (4)	0.0244 (4)	0.0265 (4)	-0.0057 (3)	-0.0023 (3)	-0.0036 (3)
N1	0.030 (3)	0.019 (2)	0.024 (2)	-0.006 (2)	-0.003 (2)	-0.0036 (19)
C1	0.044 (4)	0.028 (3)	0.020 (3)	-0.005 (3)	-0.001 (3)	-0.004 (3)
C2	0.045 (4)	0.041 (4)	0.033 (4)	-0.006 (3)	0.005 (3)	-0.011 (3)
C3	0.042 (4)	0.043 (4)	0.026 (3)	-0.002 (3)	0.000 (3)	-0.003 (3)
C4	0.036 (4)	0.029 (3)	0.033 (3)	-0.006 (3)	0.002 (3)	-0.002 (3)
C5	0.035 (4)	0.035 (3)	0.032 (3)	-0.009 (3)	-0.007 (3)	0.006 (3)
C6	0.028 (3)	0.032 (3)	0.027 (3)	-0.008 (3)	-0.005 (3)	-0.008 (3)
C7	0.031 (3)	0.026 (3)	0.032 (3)	-0.008 (3)	-0.004 (3)	-0.009 (3)
N2	0.034 (3)	0.026 (2)	0.020 (2)	-0.005 (2)	0.000 (2)	-0.010 (2)
C8	0.036 (4)	0.020 (3)	0.030 (3)	-0.009 (3)	0.000 (3)	-0.001 (3)
C9	0.042 (4)	0.019 (3)	0.035 (3)	-0.007 (3)	0.001 (3)	-0.001 (3)
C10	0.033 (4)	0.024 (3)	0.044 (4)	-0.002 (3)	0.000 (3)	-0.003 (3)
C11	0.039 (4)	0.031 (3)	0.038 (4)	-0.012 (3)	0.011 (3)	-0.009 (3)
C12	0.036 (4)	0.035 (3)	0.038 (4)	-0.008 (3)	-0.004 (3)	-0.006 (3)
C13	0.025 (3)	0.019 (3)	0.036 (3)	-0.001 (2)	-0.001 (3)	-0.006 (3)
C14	0.039 (4)	0.034 (3)	0.031 (3)	-0.007 (3)	-0.004 (3)	-0.007 (3)
C15	0.034 (3)	0.029 (3)	0.025 (3)	-0.009 (3)	0.001 (3)	-0.009 (3)
C16	0.029 (3)	0.024 (3)	0.023 (3)	-0.008 (2)	-0.004 (2)	0.001 (2)
C17	0.029 (3)	0.032 (3)	0.032 (3)	-0.004 (3)	-0.010 (3)	-0.005 (3)
C18	0.039 (4)	0.040 (4)	0.042 (4)	0.000 (3)	-0.019 (3)	-0.005 (3)
C19	0.036 (4)	0.045 (4)	0.046 (4)	0.007 (3)	-0.014 (3)	-0.013 (3)
C20	0.033 (4)	0.039 (4)	0.036 (4)	-0.002 (3)	-0.008 (3)	-0.006 (3)
C21	0.024 (3)	0.035 (3)	0.023 (3)	-0.004 (3)	-0.009 (2)	0.000 (3)
N3	0.024 (3)	0.026 (2)	0.019 (2)	0.008 (2)	-0.001 (2)	0.002 (2)
C22	0.041 (4)	0.018 (3)	0.033 (3)	-0.008 (3)	-0.006 (3)	-0.001 (2)
C23	0.029 (3)	0.019 (3)	0.029 (3)	-0.007 (2)	0.000 (3)	-0.003 (2)
C24	0.033 (4)	0.039 (4)	0.035 (4)	-0.007 (3)	-0.001 (3)	-0.007 (3)
C25	0.041 (4)	0.031 (3)	0.044 (4)	-0.013 (3)	0.004 (3)	-0.013 (3)
C26	0.039 (4)	0.041 (4)	0.040 (4)	-0.009 (3)	-0.003 (3)	-0.015 (3)
C27	0.026 (3)	0.036 (3)	0.026 (3)	-0.003 (3)	-0.003 (3)	-0.008 (3)
C28	0.065 (5)	0.049 (4)	0.036 (4)	-0.006 (4)	-0.014 (3)	-0.008 (3)
N4	0.033 (3)	0.025 (2)	0.025 (3)	-0.006 (2)	-0.005 (2)	-0.009 (2)
O1	0.036 (3)	0.034 (2)	0.028 (2)	-0.0085 (19)	0.0022 (19)	-0.0041 (19)
O2	0.034 (2)	0.028 (2)	0.024 (2)	-0.0031 (18)	-0.0057 (18)	-0.0012 (17)
C29	0.037 (4)	0.032 (3)	0.028 (3)	-0.006 (3)	-0.005 (3)	-0.009 (3)
C30	0.032 (4)	0.053 (4)	0.045 (4)	0.003 (3)	-0.002 (3)	-0.002 (3)
C11	0.0513 (11)	0.0365 (9)	0.0393 (9)	-0.0018 (8)	-0.0136 (8)	-0.0081 (7)
O3	0.092 (4)	0.046 (3)	0.081 (4)	-0.033 (3)	-0.013 (3)	-0.015 (3)
O4	0.067 (4)	0.065 (3)	0.068 (3)	-0.028 (3)	-0.032 (3)	-0.008 (3)
O5	0.083 (4)	0.068 (3)	0.065 (4)	0.045 (3)	-0.022 (3)	-0.020 (3)
O6	0.105 (5)	0.099 (4)	0.032 (3)	0.003 (4)	0.001 (3)	0.003 (3)

Geometric parameters (Å, °)

Ni1—O2	2.087 (4)	C15—H15A	0.9900
Ni1—O1	2.109 (4)	C15—H15B	0.9900
Ni1—N3	2.117 (4)	C16—C21	1.405 (7)
Ni1—N1	2.121 (4)	C16—C17	1.410 (7)

supplementary materials

Ni1—N2	2.133 (4)	C17—C18	1.368 (8)
Ni1—N4	2.151 (4)	C17—H17	0.9500
Ni1—C29	2.417 (6)	C18—C19	1.380 (8)
N1—C6	1.360 (6)	C18—H18	0.9500
N1—C1	1.361 (6)	C19—C20	1.380 (8)
C1—C3	1.396 (7)	C19—H19	0.9500
C1—C2	1.491 (7)	C20—C21	1.394 (7)
C2—H2A	0.9800	C20—H20	0.9500
C2—H2B	0.9800	C21—N3	1.428 (7)
C2—H2C	0.9800	N3—C22	1.309 (6)
C3—C4	1.378 (7)	C22—C23	1.449 (8)
C3—H3	0.9500	C22—H22	0.9500
C4—C5	1.394 (7)	C23—N4	1.356 (6)
C4—H4	0.9500	C23—C24	1.381 (7)
C5—C6	1.379 (7)	C24—C25	1.386 (8)
C5—H5	0.9500	C24—H24	0.9500
C6—C7	1.473 (7)	C25—C26	1.370 (8)
C7—N2	1.269 (6)	C25—H25	0.9500
C7—H7	0.9500	C26—C27	1.419 (7)
N2—C8	1.437 (6)	C26—H26	0.9500
C8—C13	1.393 (8)	C27—N4	1.331 (7)
C8—C9	1.399 (8)	C27—C28	1.509 (8)
C9—C10	1.394 (7)	C28—H28A	0.9800
C9—H9	0.9500	C28—H28B	0.9800
C10—C11	1.380 (8)	C28—H28C	0.9800
C10—H10	0.9500	O1—C29	1.265 (6)
C11—C12	1.397 (8)	O2—C29	1.270 (6)
C11—H11	0.9500	C29—C30	1.490 (8)
C12—C13	1.385 (7)	C30—H30A	0.9800
C12—H12	0.9500	C30—H30B	0.9800
C13—C14	1.517 (7)	C30—H30C	0.9800
C14—C15	1.556 (7)	C11—O6	1.422 (4)
C14—H14A	0.9900	C11—O5	1.426 (4)
C14—H14B	0.9900	C11—O3	1.428 (4)
C15—C16	1.490 (7)	C11—O4	1.429 (5)
O2—Ni1—O1	63.17 (14)	C16—C15—C14	109.9 (5)
O2—Ni1—N3	96.41 (15)	C16—C15—H15A	109.7
O1—Ni1—N3	158.19 (15)	C14—C15—H15A	109.7
O2—Ni1—N1	164.65 (15)	C16—C15—H15B	109.7
O1—Ni1—N1	103.58 (15)	C14—C15—H15B	109.7
N3—Ni1—N1	97.63 (17)	H15A—C15—H15B	108.2
O2—Ni1—N2	90.38 (15)	C21—C16—C17	116.9 (5)
O1—Ni1—N2	81.85 (16)	C21—C16—C15	123.1 (5)
N3—Ni1—N2	107.24 (18)	C17—C16—C15	120.0 (5)
N1—Ni1—N2	79.57 (16)	C18—C17—C16	121.8 (6)
O2—Ni1—N4	84.69 (15)	C18—C17—H17	119.1
O1—Ni1—N4	90.08 (17)	C16—C17—H17	119.1
N3—Ni1—N4	79.87 (18)	C17—C18—C19	120.6 (6)
N1—Ni1—N4	103.85 (16)	C17—C18—H18	119.7

N2—Ni1—N4	171.81 (19)	C19—C18—H18	119.7
O2—Ni1—C29	31.69 (15)	C20—C19—C18	119.5 (6)
O1—Ni1—C29	31.52 (16)	C20—C19—H19	120.3
N3—Ni1—C29	127.48 (18)	C18—C19—H19	120.3
N1—Ni1—C29	134.88 (19)	C19—C20—C21	120.5 (6)
N2—Ni1—C29	86.64 (18)	C19—C20—H20	119.7
N4—Ni1—C29	85.73 (18)	C21—C20—H20	119.7
C6—N1—C1	116.9 (4)	C20—C21—C16	120.7 (5)
C6—N1—Ni1	111.2 (3)	C20—C21—N3	119.2 (5)
C1—N1—Ni1	131.9 (4)	C16—C21—N3	120.1 (5)
N1—C1—C3	120.6 (5)	C22—N3—C21	115.2 (5)
N1—C1—C2	119.5 (5)	C22—N3—Ni1	110.9 (4)
C3—C1—C2	119.8 (5)	C21—N3—Ni1	133.9 (4)
C1—C2—H2A	109.5	N3—C22—C23	120.6 (5)
C1—C2—H2B	109.5	N3—C22—H22	119.7
H2A—C2—H2B	109.5	C23—C22—H22	119.7
C1—C2—H2C	109.5	N4—C23—C24	123.5 (6)
H2A—C2—H2C	109.5	N4—C23—C22	116.9 (5)
H2B—C2—H2C	109.5	C24—C23—C22	119.6 (5)
C4—C3—C1	121.9 (5)	C23—C24—C25	117.6 (6)
C4—C3—H3	119.1	C23—C24—H24	121.2
C1—C3—H3	119.1	C25—C24—H24	121.2
C3—C4—C5	117.4 (5)	C26—C25—C24	120.0 (6)
C3—C4—H4	121.3	C26—C25—H25	120.0
C5—C4—H4	121.3	C24—C25—H25	120.0
C6—C5—C4	118.6 (5)	C25—C26—C27	119.2 (6)
C6—C5—H5	120.7	C25—C26—H26	120.4
C4—C5—H5	120.7	C27—C26—H26	120.4
N1—C6—C5	124.5 (5)	N4—C27—C26	120.9 (6)
N1—C6—C7	116.0 (5)	N4—C27—C28	120.5 (5)
C5—C6—C7	119.5 (5)	C26—C27—C28	118.6 (6)
N2—C7—C6	120.5 (5)	C27—C28—H28A	109.5
N2—C7—H7	119.8	C27—C28—H28B	109.5
C6—C7—H7	119.8	H28A—C28—H28B	109.5
C7—N2—C8	115.8 (4)	C27—C28—H28C	109.5
C7—N2—Ni1	111.2 (3)	H28A—C28—H28C	109.5
C8—N2—Ni1	129.7 (3)	H28B—C28—H28C	109.5
C13—C8—C9	120.0 (5)	C27—N4—C23	118.6 (5)
C13—C8—N2	120.6 (5)	C27—N4—Ni1	130.4 (4)
C9—C8—N2	119.4 (5)	C23—N4—Ni1	109.1 (4)
C10—C9—C8	120.1 (6)	C29—O1—Ni1	87.8 (4)
C10—C9—H9	119.9	C29—O2—Ni1	88.7 (3)
C8—C9—H9	119.9	O1—C29—O2	120.2 (6)
C11—C10—C9	119.8 (6)	O1—C29—C30	119.3 (6)
C11—C10—H10	120.1	O2—C29—C30	120.5 (5)
C9—C10—H10	120.1	O1—C29—Ni1	60.7 (3)
C10—C11—C12	119.9 (6)	O2—C29—Ni1	59.7 (3)
C10—C11—H11	120.0	C30—C29—Ni1	175.6 (4)
C12—C11—H11	120.0	C29—C30—H30A	109.5

supplementary materials

C13—C12—C11	120.8 (6)	C29—C30—H30B	109.5
C13—C12—H12	119.6	H30A—C30—H30B	109.5
C11—C12—H12	119.6	C29—C30—H30C	109.5
C12—C13—C8	119.3 (5)	H30A—C30—H30C	109.5
C12—C13—C14	121.0 (5)	H30B—C30—H30C	109.5
C8—C13—C14	119.7 (5)	O6—C11—O5	110.9 (3)
C13—C14—C15	113.1 (5)	O6—C11—O3	109.8 (3)
C13—C14—H14A	109.0	O5—C11—O3	108.6 (3)
C15—C14—H14A	109.0	O6—C11—O4	109.7 (3)
C13—C14—H14B	109.0	O5—C11—O4	109.7 (3)
C15—C14—H14B	109.0	O3—C11—O4	108.1 (3)
H14A—C14—H14B	107.8		
O2—Ni1—N1—C6	43.6 (9)	C15—C16—C21—N3	-3.0 (8)
O1—Ni1—N1—C6	72.7 (4)	C20—C21—N3—C22	56.1 (7)
N3—Ni1—N1—C6	-112.4 (4)	C16—C21—N3—C22	-123.4 (5)
N2—Ni1—N1—C6	-6.2 (4)	C20—C21—N3—Ni1	-122.4 (5)
N4—Ni1—N1—C6	166.2 (4)	C16—C21—N3—Ni1	58.1 (7)
C29—Ni1—N1—C6	68.3 (5)	O2—Ni1—N3—C22	74.9 (4)
O2—Ni1—N1—C1	-136.2 (6)	O1—Ni1—N3—C22	55.2 (6)
O1—Ni1—N1—C1	-107.1 (5)	N1—Ni1—N3—C22	-111.3 (4)
N3—Ni1—N1—C1	67.8 (5)	N2—Ni1—N3—C22	167.3 (3)
N2—Ni1—N1—C1	174.0 (5)	N4—Ni1—N3—C22	-8.5 (4)
N4—Ni1—N1—C1	-13.6 (5)	C29—Ni1—N3—C22	68.1 (4)
C29—Ni1—N1—C1	-111.5 (5)	O2—Ni1—N3—C21	-106.6 (5)
C6—N1—C1—C3	-4.2 (8)	O1—Ni1—N3—C21	-126.3 (5)
Ni1—N1—C1—C3	175.6 (4)	N1—Ni1—N3—C21	67.2 (5)
C6—N1—C1—C2	177.3 (5)	N2—Ni1—N3—C21	-14.2 (5)
Ni1—N1—C1—C2	-2.9 (9)	N4—Ni1—N3—C21	170.0 (5)
N1—C1—C3—C4	3.0 (10)	C29—Ni1—N3—C21	-113.4 (5)
C2—C1—C3—C4	-178.4 (6)	C21—N3—C22—C23	-177.1 (5)
C1—C3—C4—C5	-0.5 (10)	Ni1—N3—C22—C23	1.7 (6)
C3—C4—C5—C6	-0.6 (9)	N3—C22—C23—N4	11.5 (8)
C1—N1—C6—C5	3.2 (9)	N3—C22—C23—C24	-169.8 (5)
Ni1—N1—C6—C5	-176.6 (5)	N4—C23—C24—C25	-2.6 (8)
C1—N1—C6—C7	-178.8 (5)	C22—C23—C24—C25	178.8 (5)
Ni1—N1—C6—C7	1.4 (6)	C23—C24—C25—C26	3.9 (8)
C4—C5—C6—N1	-0.8 (9)	C24—C25—C26—C27	-0.6 (8)
C4—C5—C6—C7	-178.7 (5)	C25—C26—C27—N4	-4.4 (8)
N1—C6—C7—N2	8.9 (8)	C25—C26—C27—C28	173.3 (5)
C5—C6—C7—N2	-173.0 (6)	C26—C27—N4—C23	5.7 (8)
C6—C7—N2—C8	-175.5 (5)	C28—C27—N4—C23	-172.0 (5)
C6—C7—N2—Ni1	-13.9 (7)	C26—C27—N4—Ni1	-156.9 (4)
O2—Ni1—N2—C7	-157.5 (4)	C28—C27—N4—Ni1	25.5 (8)
O1—Ni1—N2—C7	-94.7 (4)	C24—C23—N4—C27	-2.2 (8)
N3—Ni1—N2—C7	105.6 (4)	C22—C23—N4—C27	176.4 (5)
N1—Ni1—N2—C7	10.8 (4)	C24—C23—N4—Ni1	163.8 (4)
C29—Ni1—N2—C7	-126.0 (4)	C22—C23—N4—Ni1	-17.6 (6)
O2—Ni1—N2—C8	0.8 (5)	O2—Ni1—N4—C27	80.4 (5)
O1—Ni1—N2—C8	63.7 (5)	O1—Ni1—N4—C27	17.3 (5)

N3—Ni1—N2—C8	-96.0 (5)	N3—Ni1—N4—C27	177.9 (5)
N1—Ni1—N2—C8	169.2 (5)	N1—Ni1—N4—C27	-86.7 (5)
C29—Ni1—N2—C8	32.3 (5)	C29—Ni1—N4—C27	48.6 (5)
C7—N2—C8—C13	-105.4 (6)	O2—Ni1—N4—C23	-83.4 (3)
Ni1—N2—C8—C13	97.1 (6)	O1—Ni1—N4—C23	-146.5 (3)
C7—N2—C8—C9	73.2 (7)	N3—Ni1—N4—C23	14.1 (3)
Ni1—N2—C8—C9	-84.3 (6)	N1—Ni1—N4—C23	109.5 (3)
C13—C8—C9—C10	1.3 (8)	C29—Ni1—N4—C23	-115.2 (4)
N2—C8—C9—C10	-177.3 (5)	O2—Ni1—O1—C29	-2.3 (3)
C8—C9—C10—C11	0.4 (8)	N3—Ni1—O1—C29	19.7 (6)
C9—C10—C11—C12	-0.7 (9)	N1—Ni1—O1—C29	-174.0 (3)
C10—C11—C12—C13	-0.6 (9)	N2—Ni1—O1—C29	-96.9 (3)
C11—C12—C13—C8	2.3 (8)	N4—Ni1—O1—C29	81.7 (3)
C11—C12—C13—C14	-178.3 (5)	O1—Ni1—O2—C29	2.3 (3)
C9—C8—C13—C12	-2.6 (8)	N3—Ni1—O2—C29	-169.6 (3)
N2—C8—C13—C12	176.0 (5)	N1—Ni1—O2—C29	34.2 (8)
C9—C8—C13—C14	177.9 (5)	N2—Ni1—O2—C29	83.0 (3)
N2—C8—C13—C14	-3.5 (8)	N4—Ni1—O2—C29	-90.5 (3)
C12—C13—C14—C15	111.2 (6)	Ni1—O1—C29—O2	3.9 (5)
C8—C13—C14—C15	-69.3 (7)	Ni1—O1—C29—C30	-175.0 (5)
C13—C14—C15—C16	148.2 (5)	Ni1—O2—C29—O1	-4.0 (5)
C14—C15—C16—C21	-117.4 (6)	Ni1—O2—C29—C30	174.9 (5)
C14—C15—C16—C17	61.1 (6)	O2—Ni1—C29—O1	176.1 (5)
C21—C16—C17—C18	0.9 (8)	N3—Ni1—C29—O1	-170.9 (3)
C15—C16—C17—C18	-177.7 (5)	N1—Ni1—C29—O1	8.2 (4)
C16—C17—C18—C19	-0.2 (9)	N2—Ni1—C29—O1	79.9 (3)
C17—C18—C19—C20	-0.4 (9)	N4—Ni1—C29—O1	-97.1 (3)
C18—C19—C20—C21	0.2 (9)	O1—Ni1—C29—O2	-176.1 (5)
C19—C20—C21—C16	0.5 (9)	N3—Ni1—C29—O2	13.0 (4)
C19—C20—C21—N3	-179.0 (5)	N1—Ni1—C29—O2	-167.9 (3)
C17—C16—C21—C20	-1.0 (8)	N2—Ni1—C29—O2	-96.2 (3)
C15—C16—C21—C20	177.5 (5)	N4—Ni1—C29—O2	86.8 (3)
C17—C16—C21—N3	178.4 (5)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4 \cdots O1 ⁱ	0.95	2.45	3.274 (6)	144.
C7—H7 \cdots O4 ⁱⁱ	0.95	2.41	3.290 (7)	155.
C20—H20 \cdots O3 ⁱⁱⁱ	0.95	2.43	3.366 (8)	168.
C15—H15B \cdots O2	0.99	2.54	3.523 (7)	174.
C15—H15B \cdots N3	0.99	2.47	2.935 (7)	108.

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $x+1, y+1, z$; (iii) $x+1, y, z$.

Fig. 1




Fig. 2

