

Poly[[tris(*N,N*-dimethylformamide)- (μ_4 -5-nitroisophthalato)(μ_3 -5-nitro- isophthalato)dicobalt(II)] *N,N*-dimethyl- formamide monosolvate]

Meng Su,^a Zhan-Dong Huang,^a Huang Sun,^a Guang Yang^a
and Seik Weng Ng^{b*}

^aDepartment of Chemistry, Zhengzhou University, Zhengzhou 450001, People's Republic of China, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 31 August 2010; accepted 1 September 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.048; wR factor = 0.132; data-to-parameter ratio = 16.5.

In the polymeric title compound, $[\text{Co}_2(\text{C}_3\text{H}_7\text{NO})_3(\text{C}_8\text{H}_3\text{NO}_6)_2]\cdot\text{C}_3\text{H}_7\text{NO}$, one 5-nitroisophthalate dianion has its two carboxylate groups chelating to one Co^{II} atom while simultaneously coordinating to another metal atom in a μ_4 -bridging mode. The other 5-nitroisophthalate dianion has one carboxylate group chelating to a metal atom whereas the other bridges two metal atoms in a μ_3 -bridging mode. Both metal atoms show an octahedral coordination. The polymer adopts a layer motif, with the lattice dimethylformamide molecules occupying the space between adjacent layers.

Related literature

For adducts of cobalt 5-nitroisophthalate, see: Chen *et al.* (2006); Du *et al.* (2008); Guo *et al.* (2006); Liu *et al.* (2008); Luo *et al.* (2003); Wang *et al.* (2008, 2009); Xie *et al.* (2006); Ye *et al.* (2008a,b); Yuan *et al.* (2009); Zhou *et al.* (2004).



Experimental

Crystal data

$[\text{Co}_2(\text{C}_3\text{H}_7\text{NO})_3(\text{C}_8\text{H}_3\text{NO}_6)_2]\cdot\text{C}_3\text{H}_7\text{NO}$	$\beta = 92.910(2)^\circ$
	$V = 3626.7(7)\text{ \AA}^3$
$M_r = 828.47$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $\text{K}\alpha$ radiation
$a = 10.0833(12)\text{ \AA}$	$\mu = 0.99\text{ mm}^{-1}$
$b = 17.0887(19)\text{ \AA}$	$T = 293\text{ K}$
$c = 21.074(2)\text{ \AA}$	$0.40 \times 0.30 \times 0.30\text{ mm}$

Data collection

Bruker SMART diffractometer	21143 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	7876 independent reflections
$T_{\min} = 0.796$, $T_{\max} = 1$	5128 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$	54 restraints
$wR(F^2) = 0.132$	H-atom parameters constrained
$S = 0.99$	$\Delta\rho_{\max} = 0.57\text{ e \AA}^{-3}$
7876 reflections	$\Delta\rho_{\min} = -0.45\text{ e \AA}^{-3}$
477 parameters	

Table 1
Selected bond lengths (\AA).

Co1—O1	2.001 (2)	Co2—O2	2.042 (2)
Co1—O3 ⁱ	2.247 (2)	Co2—O7	2.091 (2)
Co1—O4 ⁱ	2.058 (2)	Co2—O9 ⁱⁱ	2.082 (2)
Co1—O7	2.383 (3)	Co2—O13	2.139 (3)
Co1—O8	2.082 (3)	Co2—O14	2.100 (2)
Co1—O10 ⁱⁱ	2.008 (2)	Co2—O15	2.063 (2)

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1999); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX* (Dolomanov *et al.*, 2003) and *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank the National Science Foundation of China (No. J0830412) and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5344).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (1999). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, H.-J., Zhang, J., Feng, W.-L. & Fu, M. (2006). *Inorg. Chem. Commun.* **9**, 300–303.
- Dolomanov, O. V., Blake, A. J., Champness, N. R. & Schröder, M. (2003). *J. Appl. Cryst.* **36**, 1283–1284.
- Du, M., Zhang, Z.-H., You, Y.-P. & Zhao, X.-J. (2008). *CrystEngComm*, **10**, 306–321.
- Guo, J.-Y., Zhang, J.-G. & Zhang, T.-L. (2006). *Acta Chim. Sin.* **64**, 1693–1699.
- Liu, Y., He, Q., Zhang, X., Xue, Z. & Lv, C. (2008). *Acta Cryst. E* **64**, m1605–m1606.
- Luo, J., Hong, M., Wang, R., Cao, R., Han, L., Yuan, D., Lin, Z. & Zhou, Y. (2003). *Inorg. Chem.* **42**, 4486–4488.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

- Wang, H.-D., Li, M.-M. & He, H.-Y. (2009). *Acta Cryst. E* **65**, m510.
- Wang, Q., Wu, M.-J., Yang, E.-C., Wang, X.-G. & Zhao, X.-J. (2008). *J. Coord. Chem.* **61**, 595–604.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Xie, G., Zeng, M.-H., Chen, S.-P. & Gao, S.-L. (2006). *Acta Cryst. E* **62**, m397–m399.
- Ye, J.-W., Li, D., Ye, K.-Q., Liu, Y., Zhao, Y.-F. & Zhang, P. (2008a). *Z. Anorg. Allg. Chem.* **634**, 345–351.
- Ye, J., Wang, J., Wu, Y., Ye, L. & Zhang, P. (2008b). *J. Mol. Struct.* **873**, 35–40.
- Yuan, S., Zhang, W.-Y., Shu, K.-X., Liang, Y.-L. & Yan, G.-F. (2009). *Z. Krist. New Cryst. Struct.* **224**, 203–204.
- Zhou, Y.-L., He, H.-Y. & Zhu, L.-G. (2004). *Chin. J. Inorg. Chem.* **20**, 576–579.

supplementary materials

Acta Cryst. (2010). E66, m1220-m1221 [doi:10.1107/S1600536810035270]

**Poly[[tris(*N,N*-dimethylformamide)(μ_4 -5-nitroisophthalato)(μ_3 -5-nitroisophthalato)dicobalt(II)]
N,N-dimethylformamide monosolvate]**

M. Su, Z.-D. Huang, H. Sun, G. Yang and S. W. Ng

Comment

Cobalt 5-nitroisophthalate forms a number of adducts with neutral ligands (Chen *et al.*, 2006; Du *et al.*, 2008; Guo *et al.*, 2006; Liu *et al.*, 2008; Luo *et al.*, 2003; Wang *et al.*, 2008; Wang *et al.*, 2009; Xie *et al.*, 2006; Ye *et al.*, 2008a; Ye *et al.*, 2008b; Yuan *et al.*, 2009; Zhou *et al.*, 2004). The structure of the parent coordination polymer has not been reported. The attempt to synthesize this compound by using DMF as the solvent gave instead the DMF-coordinated polymer, who crystallizes as a DMF solvate (Scheme I).

In polymeric $\text{Co}_2(\text{DMF})_3(\text{C}_8\text{H}_3\text{NO}_6)_2 \cdot \text{DMF}$, the 5-nitroisophthalate units show different binding modes. With one dianion, each carboxyl $-\text{CO}_2$ fragment chelates to one cobalt(II) atom while simultaneously coordinating to another metal atom, *i.e.*, this dianion functions in a μ_4 -bridging mode. With the other, one carboxyl fragment chelates to a metal atom whereas the other bridges two metal atoms, the dianion functioning in a μ_3 -bridging mode. Both metal atoms show octahedral coordination. Of the two independent metal atoms, one is coordinated by three DMF molecules. The bridging mode exercised by the dianion gives rise to a layer motif (Fig. 1). The lattice DMF molecules occupy the space between adjacent layers.

Experimental

Cobalt(II) nitrate hexahydrate (0.5 mmol, 0.146 g) and 5-nitroisophthalic acid (0.5 mmol, 0.106 g) were dissolved in DMF (2 ml); triethylamine was allowed to diffuse into the solution. Purple crystals were obtained in 30% yield after one week.

Refinement

H-atoms were placed in calculated positions [$\text{C}-\text{H}$ 0.93–0.96 Å, $U(\text{H})$ 1.2–1.5 $U(\text{C})$] and were included in the refinement in the riding model approximation.

For the coordinated and lattice DMF molecules, the $\text{C}-\text{O}$ distance was restrained to 1.25 ± 0.01 Å, the $\text{N}-\text{C}_{\text{carbonyl}}$ distance to 1.35 ± 0.01 and the $\text{N}-\text{C}_{\text{methyl}}$ distances to 1.45 ± 0.01 Å; the molecule was restrained to lie on a plane to within 0.01 Å. For the lattice DMF molecule, the anisotropic displacement parameters of the non-H atoms were restrained to be nearly isotropic.

Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of a portion of the layer structure of $\text{Co}_2(\text{DMF})_3(\text{C}_8\text{H}_3\text{NO}_6)_2 \cdot \text{DMF}$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. Symmetry codes are given in Table 1.

supplementary materials

Poly[[tris(*N,N*-dimethylformamide)(μ_4 -5- \backslash nitroisophthalato)(μ_3 -5-nitroisophthalato)dicobalt(II)] *N,N*-dimethyl-formamide monosolvate]

Crystal data

[Co ₂ (C ₃ H ₇ NO) ₃ (C ₈ H ₃ NO ₆) ₂]·C ₃ H ₇ NO	$F(000) = 1704$
$M_r = 828.47$	$D_x = 1.517 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 1014 reflections
$a = 10.0833 (12) \text{ \AA}$	$\theta = 2.3\text{--}26.7^\circ$
$b = 17.0887 (19) \text{ \AA}$	$\mu = 0.99 \text{ mm}^{-1}$
$c = 21.074 (2) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 92.910 (2)^\circ$	Block, purple
$V = 3626.7 (7) \text{ \AA}^3$	$0.40 \times 0.30 \times 0.30 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART diffractometer	7876 independent reflections
Radiation source: fine-focus sealed tube graphite	5128 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.036$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\max} = 27.0^\circ, \theta_{\min} = 2.0^\circ$
$T_{\min} = 0.796, T_{\max} = 1$	$h = -12 \rightarrow 11$
21143 measured reflections	$k = -21 \rightarrow 20$
	$l = -24 \rightarrow 26$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.048$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.132$	H-atom parameters constrained
$S = 0.99$	$w = 1/[\sigma^2(F_o^2) + (0.0741P)^2 + 0.1674P]$
7876 reflections	where $P = (F_o^2 + 2F_c^2)/3$
477 parameters	$(\Delta/\sigma)_{\max} = 0.001$
54 restraints	$\Delta\rho_{\max} = 0.57 \text{ e \AA}^{-3}$
	$\Delta\rho_{\min} = -0.45 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.54070 (4)	0.62850 (2)	0.36883 (2)	0.03535 (14)

Co2	0.75875 (4)	0.69413 (2)	0.25276 (2)	0.03577 (14)
O1	0.7161 (2)	0.61369 (15)	0.41609 (12)	0.0509 (6)
O2	0.8391 (2)	0.65261 (16)	0.33703 (12)	0.0536 (7)
O3	1.3188 (2)	0.63449 (18)	0.35333 (13)	0.0615 (8)
O4	1.4234 (2)	0.60440 (14)	0.44350 (11)	0.0452 (6)
O5	1.1571 (3)	0.5635 (2)	0.63158 (14)	0.0841 (10)
O6	0.9466 (3)	0.5792 (2)	0.62357 (14)	0.0890 (11)
O7	0.5916 (2)	0.62286 (13)	0.25975 (14)	0.0584 (7)
O8	0.5266 (3)	0.52567 (16)	0.31596 (14)	0.0641 (8)
O9	0.3476 (2)	0.28716 (13)	0.21118 (12)	0.0447 (6)
O10	0.4601 (3)	0.24599 (14)	0.12880 (13)	0.0548 (7)
O11	0.6418 (4)	0.4515 (2)	-0.00474 (17)	0.0982 (12)
O12	0.7008 (6)	0.5587 (3)	0.0369 (2)	0.160 (2)
O13	0.8704 (3)	0.60593 (15)	0.20713 (13)	0.0634 (8)
O14	0.9230 (3)	0.76957 (15)	0.25177 (13)	0.0619 (7)
O15	0.6900 (3)	0.72766 (15)	0.16294 (12)	0.0599 (7)
O16	1.0622 (10)	0.7746 (4)	0.5743 (4)	0.236 (4)
N1	1.0547 (3)	0.5774 (2)	0.60062 (15)	0.0565 (8)
N2	0.6509 (4)	0.4948 (3)	0.03991 (19)	0.0781 (11)
N3	1.0517 (3)	0.53017 (18)	0.20161 (17)	0.0715 (11)
N4	1.1460 (3)	0.7832 (2)	0.24308 (18)	0.0788 (11)
N5	0.6043 (3)	0.82156 (18)	0.09718 (13)	0.0572 (9)
N6	0.9621 (7)	0.8134 (3)	0.4885 (3)	0.1170 (18)
C1	0.8239 (3)	0.62893 (19)	0.39168 (17)	0.0385 (8)
C2	0.9490 (3)	0.61717 (18)	0.43297 (16)	0.0361 (7)
C3	1.0713 (3)	0.62258 (19)	0.40590 (16)	0.0390 (8)
H3	1.0752	0.6346	0.3630	0.047*
C4	1.1879 (3)	0.6102 (2)	0.44235 (16)	0.0381 (8)
C5	1.1835 (3)	0.5940 (2)	0.50582 (16)	0.0411 (8)
H5	1.2612	0.5851	0.5305	0.049*
C6	1.0611 (3)	0.5913 (2)	0.53248 (16)	0.0399 (8)
C7	0.9434 (3)	0.60096 (19)	0.49684 (16)	0.0394 (8)
H7	0.8621	0.5966	0.5155	0.047*
C8	1.3176 (3)	0.6162 (2)	0.41043 (18)	0.0411 (8)
C9	0.5582 (3)	0.5522 (2)	0.2641 (2)	0.0454 (9)
C10	0.5511 (3)	0.50093 (19)	0.20667 (17)	0.0400 (8)
C11	0.4916 (3)	0.42749 (18)	0.21119 (16)	0.0389 (8)
H11	0.4562	0.4127	0.2493	0.047*
C12	0.4843 (3)	0.37624 (18)	0.16031 (16)	0.0374 (7)
C13	0.5376 (3)	0.3984 (2)	0.10396 (17)	0.0458 (9)
H13	0.5358	0.3646	0.0694	0.055*
C14	0.5937 (4)	0.4717 (2)	0.10006 (18)	0.0525 (10)
C15	0.6007 (4)	0.5237 (2)	0.15018 (18)	0.0509 (9)
H15	0.6382	0.5730	0.1458	0.061*
C16	0.4250 (3)	0.29629 (19)	0.16771 (17)	0.0391 (8)
C17	0.9538 (4)	0.5620 (2)	0.2306 (2)	0.0616 (11)
H17	0.9474	0.5499	0.2734	0.074*
C18	1.0674 (8)	0.5465 (4)	0.1361 (3)	0.180 (4)
H18A	1.0040	0.5855	0.1219	0.269*

supplementary materials

H18B	1.0532	0.4996	0.1117	0.269*
H18C	1.1556	0.5656	0.1306	0.269*
C19	1.1493 (5)	0.4793 (3)	0.2333 (3)	0.132 (3)
H19A	1.1219	0.4670	0.2751	0.198*
H19B	1.2337	0.5054	0.2365	0.198*
H19C	1.1569	0.4319	0.2092	0.198*
C20	1.0260 (4)	0.7575 (2)	0.2246 (2)	0.0653 (11)
H20	1.0193	0.7276	0.1877	0.078*
C21	1.1625 (6)	0.8289 (5)	0.2992 (3)	0.167 (4)
H21A	1.0866	0.8224	0.3244	0.250*
H21B	1.1710	0.8831	0.2880	0.250*
H21C	1.2409	0.8123	0.3232	0.250*
C22	1.2638 (5)	0.7666 (4)	0.2097 (3)	0.123 (2)
H22A	1.2399	0.7391	0.1711	0.184*
H22B	1.3228	0.7349	0.2360	0.184*
H22C	1.3071	0.8148	0.1999	0.184*
C23	0.6784 (3)	0.7969 (2)	0.14649 (15)	0.0503 (9)
H23	0.7252	0.8342	0.1707	0.060*
C24	0.5276 (6)	0.7661 (3)	0.0572 (2)	0.099 (2)
H24A	0.5459	0.7138	0.0716	0.149*
H24B	0.5520	0.7713	0.0139	0.149*
H24C	0.4346	0.7769	0.0597	0.149*
C25	0.5940 (4)	0.9038 (2)	0.08090 (19)	0.0641 (11)
H25A	0.6564	0.9332	0.1073	0.096*
H25B	0.5056	0.9218	0.0875	0.096*
H25C	0.6132	0.9108	0.0371	0.096*
C26	0.9403 (11)	0.7817 (4)	0.5443 (5)	0.175 (4)
H26	0.8589	0.7673	0.5596	0.210*
C27	1.0928 (11)	0.8320 (7)	0.4728 (6)	0.267 (6)
H27A	1.1542	0.8143	0.5061	0.400*
H27B	1.1009	0.8876	0.4680	0.400*
H27C	1.1123	0.8066	0.4337	0.400*
C28	0.8669 (10)	0.8325 (5)	0.4374 (4)	0.176 (4)
H28A	0.7826	0.8096	0.4456	0.265*
H28B	0.8972	0.8124	0.3981	0.265*
H28C	0.8577	0.8884	0.4344	0.265*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0234 (2)	0.0368 (2)	0.0459 (3)	-0.00044 (18)	0.00267 (17)	0.0023 (2)
Co2	0.0301 (2)	0.0324 (2)	0.0445 (3)	0.00062 (18)	-0.00140 (18)	0.00308 (19)
O1	0.0218 (12)	0.0657 (16)	0.0653 (17)	-0.0010 (11)	0.0021 (11)	0.0089 (13)
O2	0.0333 (14)	0.0776 (18)	0.0495 (16)	0.0072 (13)	-0.0015 (11)	0.0134 (14)
O3	0.0308 (14)	0.105 (2)	0.0496 (17)	-0.0015 (14)	0.0063 (11)	0.0135 (15)
O4	0.0236 (12)	0.0602 (15)	0.0515 (15)	0.0024 (11)	0.0001 (10)	0.0052 (12)
O5	0.073 (2)	0.128 (3)	0.0500 (18)	0.019 (2)	-0.0080 (15)	0.0166 (18)
O6	0.066 (2)	0.144 (3)	0.059 (2)	0.004 (2)	0.0244 (16)	0.0161 (19)

O7	0.0414 (15)	0.0348 (13)	0.098 (2)	-0.0069 (11)	-0.0058 (14)	-0.0051 (13)
O8	0.075 (2)	0.0554 (17)	0.0625 (19)	-0.0133 (15)	0.0114 (15)	-0.0148 (14)
O9	0.0435 (14)	0.0315 (12)	0.0598 (16)	-0.0068 (10)	0.0106 (12)	-0.0034 (11)
O10	0.0599 (17)	0.0379 (14)	0.0683 (18)	-0.0077 (12)	0.0183 (13)	-0.0083 (13)
O11	0.128 (3)	0.104 (3)	0.066 (2)	-0.006 (2)	0.035 (2)	0.008 (2)
O12	0.243 (6)	0.148 (4)	0.096 (3)	-0.123 (4)	0.055 (3)	0.006 (3)
O13	0.0591 (18)	0.0579 (16)	0.0722 (19)	0.0208 (14)	-0.0057 (14)	-0.0053 (14)
O14	0.0412 (15)	0.0610 (17)	0.085 (2)	-0.0156 (13)	0.0186 (14)	-0.0011 (15)
O15	0.080 (2)	0.0464 (15)	0.0516 (16)	0.0092 (14)	-0.0116 (14)	0.0047 (13)
O16	0.329 (8)	0.147 (5)	0.223 (6)	0.002 (6)	-0.068 (6)	-0.057 (5)
N1	0.054 (2)	0.070 (2)	0.046 (2)	0.0064 (17)	0.0045 (17)	0.0046 (16)
N2	0.087 (3)	0.089 (3)	0.059 (3)	-0.026 (2)	0.013 (2)	0.011 (2)
N3	0.072 (3)	0.068 (2)	0.077 (3)	0.031 (2)	0.028 (2)	0.009 (2)
N4	0.045 (2)	0.097 (3)	0.095 (3)	-0.014 (2)	0.008 (2)	0.003 (2)
N5	0.067 (2)	0.0534 (19)	0.0494 (19)	0.0082 (17)	-0.0100 (16)	0.0036 (15)
N6	0.134 (5)	0.098 (4)	0.120 (4)	-0.012 (3)	0.020 (4)	-0.034 (3)
C1	0.0256 (17)	0.0405 (18)	0.049 (2)	0.0047 (14)	0.0006 (14)	0.0002 (16)
C2	0.0226 (15)	0.0384 (17)	0.047 (2)	0.0009 (13)	0.0020 (13)	0.0032 (15)
C3	0.0296 (17)	0.0478 (19)	0.0393 (19)	-0.0016 (15)	0.0001 (14)	0.0044 (15)
C4	0.0255 (16)	0.0474 (19)	0.041 (2)	-0.0011 (14)	0.0003 (13)	0.0005 (15)
C5	0.0293 (17)	0.0479 (19)	0.046 (2)	0.0005 (15)	-0.0047 (14)	-0.0006 (16)
C6	0.0349 (18)	0.0449 (19)	0.0397 (19)	0.0038 (15)	0.0018 (14)	0.0041 (15)
C7	0.0262 (17)	0.0438 (18)	0.049 (2)	0.0032 (14)	0.0089 (14)	0.0034 (16)
C8	0.0228 (17)	0.049 (2)	0.052 (2)	-0.0035 (14)	0.0025 (15)	0.0001 (17)
C9	0.0286 (18)	0.038 (2)	0.069 (3)	-0.0006 (15)	-0.0030 (17)	-0.0078 (18)
C10	0.0320 (18)	0.0347 (17)	0.053 (2)	-0.0024 (14)	0.0014 (15)	0.0011 (16)
C11	0.0298 (17)	0.0367 (18)	0.050 (2)	-0.0007 (14)	0.0031 (14)	0.0028 (15)
C12	0.0287 (16)	0.0349 (17)	0.048 (2)	0.0004 (14)	-0.0018 (14)	-0.0023 (15)
C13	0.043 (2)	0.047 (2)	0.047 (2)	-0.0009 (17)	-0.0013 (16)	-0.0033 (17)
C14	0.050 (2)	0.057 (2)	0.051 (2)	-0.0101 (19)	0.0047 (18)	0.0104 (19)
C15	0.049 (2)	0.0386 (19)	0.065 (3)	-0.0122 (17)	-0.0025 (18)	0.0105 (18)
C16	0.0316 (18)	0.0357 (18)	0.049 (2)	-0.0005 (14)	-0.0046 (15)	-0.0023 (16)
C17	0.062 (3)	0.062 (3)	0.061 (3)	0.009 (2)	0.009 (2)	0.006 (2)
C18	0.281 (11)	0.150 (7)	0.117 (6)	0.123 (7)	0.099 (6)	0.046 (5)
C19	0.105 (5)	0.165 (6)	0.128 (5)	0.084 (5)	0.029 (4)	0.041 (5)
C20	0.051 (3)	0.071 (3)	0.073 (3)	-0.015 (2)	0.001 (2)	-0.007 (2)
C21	0.084 (5)	0.256 (10)	0.159 (7)	-0.058 (6)	-0.004 (4)	-0.110 (7)
C22	0.049 (3)	0.177 (7)	0.146 (6)	-0.008 (4)	0.026 (3)	-0.008 (5)
C23	0.054 (2)	0.054 (2)	0.043 (2)	0.0032 (19)	0.0003 (17)	0.0010 (18)
C24	0.136 (5)	0.072 (3)	0.085 (4)	0.001 (3)	-0.055 (3)	-0.002 (3)
C25	0.076 (3)	0.059 (3)	0.057 (3)	0.015 (2)	-0.005 (2)	0.009 (2)
C26	0.238 (9)	0.122 (6)	0.169 (7)	-0.015 (6)	0.044 (7)	-0.062 (6)
C27	0.241 (10)	0.278 (10)	0.282 (10)	-0.034 (8)	0.011 (8)	-0.061 (8)
C28	0.197 (8)	0.158 (6)	0.171 (7)	0.023 (6)	-0.014 (6)	-0.016 (6)

Geometric parameters (\AA , $^\circ$)

Co1—O1	2.001 (2)	C2—C3	1.388 (4)
Co1—O3 ⁱ	2.247 (2)	C3—C4	1.387 (4)

supplementary materials

Co1—O4 ⁱ	2.058 (2)	C3—H3	0.9300
Co1—O7	2.383 (3)	C4—C5	1.369 (5)
Co1—C8 ⁱ	2.465 (3)	C4—C8	1.504 (4)
Co1—O8	2.082 (3)	C5—C6	1.383 (5)
Co1—O10 ⁱⁱ	2.008 (2)	C5—H5	0.9300
Co2—O2	2.042 (2)	C6—C7	1.382 (4)
Co2—O7	2.091 (2)	C7—H7	0.9300
Co2—O9 ⁱⁱ	2.082 (2)	C8—Co1 ⁱⁱⁱ	2.465 (3)
Co2—O13	2.139 (3)	C9—C10	1.493 (5)
Co2—O14	2.100 (2)	C10—C15	1.371 (5)
Co2—O15	2.063 (2)	C10—C11	1.396 (4)
O1—C1	1.253 (4)	C11—C12	1.383 (5)
O2—C1	1.238 (4)	C11—H11	0.9300
O3—C8	1.244 (4)	C12—C13	1.381 (5)
O3—Co1 ⁱⁱⁱ	2.247 (2)	C12—C16	1.503 (4)
O4—C8	1.260 (4)	C13—C14	1.379 (5)
O4—Co1 ⁱⁱⁱ	2.058 (2)	C13—H13	0.9300
O5—N1	1.216 (4)	C14—C15	1.379 (5)
O6—N1	1.215 (4)	C15—H15	0.9300
O7—C9	1.259 (4)	C17—H17	0.9300
O8—C9	1.240 (4)	C18—H18A	0.9600
O9—C16	1.243 (4)	C18—H18B	0.9600
O9—Co2 ^{iv}	2.082 (2)	C18—H18C	0.9600
O10—C16	1.251 (4)	C19—H19A	0.9600
O10—Co1 ^{iv}	2.008 (2)	C19—H19B	0.9600
O11—N2	1.196 (5)	C19—H19C	0.9600
O12—N2	1.206 (5)	C20—H20	0.9300
O13—C17	1.214 (4)	C21—H21A	0.9600
O14—C20	1.229 (4)	C21—H21B	0.9600
O15—C23	1.237 (4)	C21—H21C	0.9600
O16—C26	1.359 (8)	C22—H22A	0.9600
N1—C6	1.460 (5)	C22—H22B	0.9600
N2—C14	1.473 (5)	C22—H22C	0.9600
N3—C17	1.305 (5)	C23—H23	0.9300
N3—C18	1.426 (6)	C24—H24A	0.9600
N3—C19	1.450 (5)	C24—H24B	0.9600
N4—C20	1.326 (5)	C24—H24C	0.9600
N4—C21	1.421 (6)	C25—H25A	0.9600
N4—C22	1.439 (5)	C25—H25B	0.9600
N5—C23	1.318 (4)	C25—H25C	0.9600
N5—C25	1.449 (4)	C26—H26	0.9300
N5—C24	1.463 (5)	C27—H27A	0.9600
N6—C26	1.322 (8)	C27—H27B	0.9600
N6—C27	1.411 (8)	C27—H27C	0.9600
N6—C28	1.444 (7)	C28—H28A	0.9600
C1—C2	1.509 (4)	C28—H28B	0.9600
C2—C7	1.378 (5)	C28—H28C	0.9600

O1—Co1—O10 ⁱⁱ	96.82 (11)	O3—C8—C4	120.1 (3)
O1—Co1—O4 ⁱ	97.00 (10)	O4—C8—C4	118.2 (3)
O10 ⁱⁱ —Co1—O4 ⁱ	100.24 (10)	O3—C8—Co1 ⁱⁱⁱ	65.20 (18)
O1—Co1—O8	101.20 (11)	O4—C8—Co1 ⁱⁱⁱ	56.52 (16)
O10 ⁱⁱ —Co1—O8	148.98 (12)	C4—C8—Co1 ⁱⁱⁱ	174.2 (3)
O4 ⁱ —Co1—O8	102.36 (11)	O8—C9—O7	119.8 (4)
O1—Co1—O3 ⁱ	157.87 (10)	O8—C9—C10	119.8 (3)
O10 ⁱⁱ —Co1—O3 ⁱ	87.29 (11)	O7—C9—C10	120.4 (4)
O4 ⁱ —Co1—O3 ⁱ	60.88 (9)	C15—C10—C11	119.5 (3)
O8—Co1—O3 ⁱ	85.42 (12)	C15—C10—C9	122.2 (3)
O1—Co1—O7	104.29 (10)	C11—C10—C9	118.4 (3)
O10 ⁱⁱ —Co1—O7	93.78 (9)	C12—C11—C10	121.5 (3)
O4 ⁱ —Co1—O7	152.88 (9)	C12—C11—H11	119.3
O8—Co1—O7	57.43 (10)	C10—C11—H11	119.3
O3 ⁱ —Co1—O7	97.09 (9)	C13—C12—C11	119.0 (3)
O1—Co1—C8 ⁱ	127.71 (11)	C13—C12—C16	120.9 (3)
O10 ⁱⁱ —Co1—C8 ⁱ	94.08 (11)	C11—C12—C16	120.1 (3)
O4 ⁱ —Co1—C8 ⁱ	30.72 (10)	C14—C13—C12	118.6 (3)
O8—Co1—C8 ⁱ	94.58 (11)	C14—C13—H13	120.7
O3 ⁱ —Co1—C8 ⁱ	30.16 (10)	C12—C13—H13	120.7
O7—Co1—C8 ⁱ	125.81 (10)	C13—C14—C15	123.0 (3)
O2—Co2—O15	173.94 (11)	C13—C14—N2	118.3 (4)
O2—Co2—O9 ⁱⁱ	97.87 (10)	C15—C14—N2	118.7 (4)
O15—Co2—O9 ⁱⁱ	88.15 (10)	C10—C15—C14	118.3 (3)
O2—Co2—O7	91.26 (11)	C10—C15—H15	120.8
O15—Co2—O7	89.54 (11)	C14—C15—H15	120.8
O9 ⁱⁱ —Co2—O7	89.27 (10)	O9—C16—O10	127.3 (3)
O2—Co2—O14	86.77 (11)	O9—C16—C12	117.3 (3)
O15—Co2—O14	92.79 (11)	O10—C16—C12	115.4 (3)
O9 ⁱⁱ —Co2—O14	87.47 (10)	O13—C17—N3	126.3 (4)
O7—Co2—O14	175.92 (11)	O13—C17—H17	116.9
O2—Co2—O13	87.22 (11)	N3—C17—H17	116.9
O15—Co2—O13	86.74 (10)	N3—C18—H18A	109.5
O9 ⁱⁱ —Co2—O13	174.03 (10)	N3—C18—H18B	109.5
O7—Co2—O13	93.76 (11)	H18A—C18—H18B	109.5
O14—Co2—O13	89.71 (11)	N3—C18—H18C	109.5
C1—O1—Co1	122.2 (2)	H18A—C18—H18C	109.5
C1—O2—Co2	149.4 (2)	H18B—C18—H18C	109.5
C8—O3—Co1 ⁱⁱⁱ	84.64 (19)	N3—C19—H19A	109.5
C8—O4—Co1 ⁱⁱⁱ	92.8 (2)	N3—C19—H19B	109.5
C9—O7—Co2	141.6 (2)	H19A—C19—H19B	109.5
C9—O7—Co1	84.1 (2)	N3—C19—H19C	109.5
Co2—O7—Co1	104.97 (11)	H19A—C19—H19C	109.5
C9—O8—Co1	98.6 (2)	H19B—C19—H19C	109.5

supplementary materials

C16—O9—Co2 ^{iv}	136.0 (2)	O14—C20—N4	126.0 (4)
C16—O10—Co1 ^{iv}	132.0 (2)	O14—C20—H20	117.0
C17—O13—Co2	128.5 (3)	N4—C20—H20	117.0
C20—O14—Co2	126.0 (3)	N4—C21—H21A	109.5
C23—O15—Co2	123.1 (2)	N4—C21—H21B	109.5
O6—N1—O5	123.1 (4)	H21A—C21—H21B	109.5
O6—N1—C6	118.2 (3)	N4—C21—H21C	109.5
O5—N1—C6	118.7 (3)	H21A—C21—H21C	109.5
O11—N2—O12	122.3 (4)	H21B—C21—H21C	109.5
O11—N2—C14	119.6 (4)	N4—C22—H22A	109.5
O12—N2—C14	118.1 (4)	N4—C22—H22B	109.5
C17—N3—C18	119.7 (4)	H22A—C22—H22B	109.5
C17—N3—C19	123.1 (4)	N4—C22—H22C	109.5
C18—N3—C19	117.2 (4)	H22A—C22—H22C	109.5
C20—N4—C21	119.5 (4)	H22B—C22—H22C	109.5
C20—N4—C22	123.7 (4)	O15—C23—N5	124.7 (4)
C21—N4—C22	116.8 (5)	O15—C23—H23	117.6
C23—N5—C25	121.9 (3)	N5—C23—H23	117.6
C23—N5—C24	120.6 (3)	N5—C24—H24A	109.5
C25—N5—C24	117.5 (3)	N5—C24—H24B	109.5
C26—N6—C27	120.0 (9)	H24A—C24—H24B	109.5
C26—N6—C28	128.5 (9)	N5—C24—H24C	109.5
C27—N6—C28	111.6 (9)	H24A—C24—H24C	109.5
O2—C1—O1	126.9 (3)	H24B—C24—H24C	109.5
O2—C1—C2	116.1 (3)	N5—C25—H25A	109.5
O1—C1—C2	116.9 (3)	N5—C25—H25B	109.5
C7—C2—C3	119.8 (3)	H25A—C25—H25B	109.5
C7—C2—C1	121.0 (3)	N5—C25—H25C	109.5
C3—C2—C1	119.2 (3)	H25A—C25—H25C	109.5
C4—C3—C2	120.5 (3)	H25B—C25—H25C	109.5
C4—C3—H3	119.7	N6—C26—O16	105.3 (9)
C2—C3—H3	119.7	N6—C26—H26	127.4
C5—C4—C3	120.2 (3)	O16—C26—H26	127.4
C5—C4—C8	121.5 (3)	N6—C27—H27A	109.5
C3—C4—C8	118.3 (3)	N6—C27—H27B	109.5
C4—C5—C6	118.5 (3)	H27A—C27—H27B	109.5
C4—C5—H5	120.7	N6—C27—H27C	109.5
C6—C5—H5	120.7	H27A—C27—H27C	109.5
C7—C6—C5	122.4 (3)	H27B—C27—H27C	109.5
C7—C6—N1	118.3 (3)	N6—C28—H28A	109.5
C5—C6—N1	119.3 (3)	N6—C28—H28B	109.5
C2—C7—C6	118.5 (3)	H28A—C28—H28B	109.5
C2—C7—H7	120.7	N6—C28—H28C	109.5
C6—C7—H7	120.7	H28A—C28—H28C	109.5
O3—C8—O4	121.7 (3)	H28B—C28—H28C	109.5
O10 ⁱⁱ —Co1—O1—C1	-72.3 (3)	C4—C5—C6—C7	2.9 (5)
O4 ⁱ —Co1—O1—C1	-173.5 (3)	C4—C5—C6—N1	-177.5 (3)
O8—Co1—O1—C1	82.4 (3)	O6—N1—C6—C7	-4.0 (5)

O3 ⁱ —Co1—O1—C1	-171.9 (3)	O5—N1—C6—C7	175.0 (4)
O7—Co1—O1—C1	23.4 (3)	O6—N1—C6—C5	176.3 (4)
C8 ⁱ —Co1—O1—C1	-172.8 (2)	O5—N1—C6—C5	-4.7 (5)
O9 ⁱⁱ —Co2—O2—C1	53.7 (5)	C3—C2—C7—C6	0.7 (5)
O7—Co2—O2—C1	-35.7 (5)	C1—C2—C7—C6	-179.5 (3)
O14—Co2—O2—C1	140.7 (5)	C5—C6—C7—C2	-2.8 (5)
O13—Co2—O2—C1	-129.4 (5)	N1—C6—C7—C2	177.5 (3)
O2—Co2—O7—C9	-59.1 (5)	Co1 ⁱⁱⁱ —O3—C8—O4	-0.7 (3)
O15—Co2—O7—C9	114.8 (5)	Co1 ⁱⁱⁱ —O3—C8—C4	177.4 (3)
O9 ⁱⁱ —Co2—O7—C9	-157.0 (5)	Co1 ⁱⁱⁱ —O4—C8—O3	0.7 (4)
O13—Co2—O7—C9	28.1 (5)	Co1 ⁱⁱⁱ —O4—C8—C4	-177.4 (3)
O2—Co2—O7—Co1	40.41 (11)	C5—C4—C8—O3	-176.8 (3)
O15—Co2—O7—Co1	-145.60 (11)	C3—C4—C8—O3	2.4 (5)
O9 ⁱⁱ —Co2—O7—Co1	-57.44 (10)	C5—C4—C8—O4	1.4 (5)
O13—Co2—O7—Co1	127.70 (10)	C3—C4—C8—O4	-179.5 (3)
O1—Co1—O7—C9	96.0 (2)	Co1—O8—C9—O7	3.2 (4)
O10 ⁱⁱ —Co1—O7—C9	-166.0 (2)	Co1—O8—C9—C10	-174.3 (3)
O4 ⁱ —Co1—O7—C9	-44.6 (3)	Co2—O7—C9—O8	103.9 (5)
O8—Co1—O7—C9	1.7 (2)	Co1—O7—C9—O8	-2.8 (3)
O3 ⁱ —Co1—O7—C9	-78.2 (2)	Co2—O7—C9—C10	-78.5 (5)
C8 ⁱ —Co1—O7—C9	-68.2 (2)	Co1—O7—C9—C10	174.8 (3)
O1—Co1—O7—Co2	-46.01 (12)	O8—C9—C10—C15	-171.4 (3)
O10 ⁱⁱ —Co1—O7—Co2	52.01 (12)	O7—C9—C10—C15	11.1 (5)
O4 ⁱ —Co1—O7—Co2	173.32 (15)	O8—C9—C10—C11	8.6 (5)
O8—Co1—O7—Co2	-140.32 (15)	O7—C9—C10—C11	-168.9 (3)
O3 ⁱ —Co1—O7—Co2	139.75 (11)	C15—C10—C11—C12	1.5 (5)
C8 ⁱ —Co1—O7—Co2	149.80 (11)	C9—C10—C11—C12	-178.5 (3)
O1—Co1—O8—C9	-101.7 (2)	C10—C11—C12—C13	0.3 (5)
O10 ⁱⁱ —Co1—O8—C9	22.7 (4)	C10—C11—C12—C16	177.3 (3)
O4 ⁱ —Co1—O8—C9	158.5 (2)	C11—C12—C13—C14	-1.5 (5)
O3 ⁱ —Co1—O8—C9	99.7 (2)	C16—C12—C13—C14	-178.5 (3)
O7—Co1—O8—C9	-1.7 (2)	C12—C13—C14—C15	0.9 (6)
C8 ⁱ —Co1—O8—C9	128.4 (2)	C12—C13—C14—N2	179.7 (3)
O2—Co2—O13—C17	-7.4 (3)	O11—N2—C14—C13	3.7 (7)
O15—Co2—O13—C17	172.2 (3)	O12—N2—C14—C13	-179.4 (5)
O7—Co2—O13—C17	-98.5 (3)	O11—N2—C14—C15	-177.5 (4)
O14—Co2—O13—C17	79.4 (3)	O12—N2—C14—C15	-0.6 (7)
O2—Co2—O14—C20	95.1 (3)	C11—C10—C15—C14	-2.0 (5)
O15—Co2—O14—C20	-78.8 (3)	C9—C10—C15—C14	178.0 (3)
O9 ⁱⁱ —Co2—O14—C20	-166.9 (3)	C13—C14—C15—C10	0.8 (6)
O13—Co2—O14—C20	7.9 (3)	N2—C14—C15—C10	-177.9 (4)
O9 ⁱⁱ —Co2—O15—C23	38.7 (2)	Co2 ^{iv} —O9—C16—O10	2.3 (6)
O7—Co2—O15—C23	128.0 (2)	Co2 ^{iv} —O9—C16—C12	-177.6 (2)
O14—Co2—O15—C23	-48.7 (2)	C13—C12—C16—O9	-160.5 (3)

supplementary materials

O13—Co2—O15—C23	-138.2 (2)	C11—C12—C16—O9	22.5 (5)
Co2—O2—C1—O1	8.8 (8)	C13—C12—C16—O10	19.6 (5)
Co2—O2—C1—C2	-171.4 (4)	C11—C12—C16—O10	-157.4 (3)
Co1—O1—C1—O2	-1.7 (5)	Co2—O13—C17—N3	-153.3 (3)
Co1—O1—C1—C2	178.5 (2)	C18—N3—C17—O13	0.2 (4)
O2—C1—C2—C7	171.4 (3)	C19—N3—C17—O13	178.7 (3)
O1—C1—C2—C7	-8.7 (5)	Co2—O14—C20—N4	-149.3 (3)
O2—C1—C2—C3	-8.8 (5)	C21—N4—C20—O14	0.2 (3)
O1—C1—C2—C3	171.1 (3)	C22—N4—C20—O14	179.4 (3)
C7—C2—C3—C4	1.3 (5)	Co2—O15—C23—N5	-161.2 (2)
C1—C2—C3—C4	-178.5 (3)	C25—N5—C23—O15	179.4 (3)
C2—C3—C4—C5	-1.3 (5)	C24—N5—C23—O15	0.2 (3)
C2—C3—C4—C8	179.6 (3)	C27—N6—C26—O16	-0.3 (3)
C3—C4—C5—C6	-0.7 (5)	C28—N6—C26—O16	179.0 (3)
C8—C4—C5—C6	178.4 (3)		

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1, y+1/2, -z+1/2$; (iii) $x+1, y, z$; (iv) $-x+1, y-1/2, -z+1/2$.

Fig. 1

