

## (R)-2-Methylpiperazine-1,4-dium diaquatetrachloridoferrate(II)

 Cong-Hu Peng<sup>a\*</sup> and Yun-Peng Li<sup>b</sup>

<sup>a</sup>Department of Chemical and Environmental Engineering, Anyang Institute of Technology, Anyang 455000, People's Republic of China, and <sup>b</sup>Anyang Administration of Work Safety, Henan Province 455000, People's Republic of China

Correspondence e-mail: ayitpch@yahoo.com.cn


Received 17 August 2010; accepted 3 September 2010

Key indicators: single-crystal X-ray study;  $T = 291$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.023;  $wR$  factor = 0.050; data-to-parameter ratio = 19.8.

In the title salt,  $(\text{C}_5\text{H}_{14}\text{N}_2)[\text{FeCl}_4(\text{H}_2\text{O})_2]$ , the  $\text{Fe}^{\text{II}}$  cation is coordinated by four  $\text{Cl}^-$  anions and two water molecules in a distorted octahedral geometry. The piperazine ring adopts a normal chair conformation. Intermolecular  $\text{N}-\text{H}\cdots\text{Cl}$ ,  $\text{N}-\text{H}\cdots(\text{Cl},\text{Cl})$  and  $\text{O}-\text{H}\cdots\text{Cl}$  hydrogen bonding is present in the crystal structure.

### Related literature

For hydrogen bonding in metal-chlorido complexes, see: Brammer *et al.* (2001); Bremner & Harrison (2003); Kefi & Nasr (2005). For the crystal structure of a related compound, piperazindium tetrachloridozincate(II), see: Sutherland & Harrison (2009).



### Experimental

#### Crystal data

 $(\text{C}_5\text{H}_{14}\text{N}_2)[\text{FeCl}_4(\text{H}_2\text{O})_2]$ 
 $M_r = 335.86$ 

 Monoclinic,  $P2_1$ 
 $a = 8.6013$  (17) Å

 $b = 6.4495$  (13) Å

 $c = 12.024$  (2) Å

 $\beta = 101.64$  (3)°

 $V = 653.3$  (2) Å<sup>3</sup>
 $Z = 2$ 

 Mo  $K\alpha$  radiation

 $\mu = 1.95$  mm<sup>-1</sup>
 $T = 291$  K

 $0.28 \times 0.24 \times 0.20$  mm

#### Data collection

Rigaku SCXmini diffractometer

Absorption correction: multi-scan

 (*CrystalClear*; Rigaku, 2005)

 $T_{\text{min}} = 0.8, T_{\text{max}} = 0.9$ 

6105 measured reflections

2558 independent reflections

 2456 reflections with  $I > 2\sigma(I)$ 
 $R_{\text{int}} = 0.025$ 

#### Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.023$ 
 $wR(F^2) = 0.050$ 
 $S = 1.08$ 

2558 reflections

129 parameters

1 restraint

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.19$  e Å<sup>-3</sup>
 $\Delta\rho_{\text{min}} = -0.24$  e Å<sup>-3</sup>

Absolute structure: Flack (1983),

1156 Friedel pairs

Flack parameter: 0.010 (14)

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1C}\cdots\text{Cl2}^{\text{i}}$	0.90	2.62	3.443 (2)	152
$\text{N1}-\text{H1C}\cdots\text{Cl4}^{\text{i}}$	0.90	2.81	3.379 (3)	122
$\text{N1}-\text{H1D}\cdots\text{Cl4}^{\text{ii}}$	0.90	2.28	3.169 (3)	167
$\text{N2}-\text{H2C}\cdots\text{Cl1}^{\text{iii}}$	0.90	2.26	3.145 (3)	168
$\text{N2}-\text{H2D}\cdots\text{Cl3}$	0.90	2.45	3.275 (2)	152
$\text{O1}-\text{H11}\cdots\text{Cl3}^{\text{iv}}$	0.82	2.33	3.147 (2)	173
$\text{O1}-\text{H12}\cdots\text{Cl3}^{\text{v}}$	0.89	2.24	3.127 (2)	176
$\text{O2}-\text{H21}\cdots\text{Cl2}^{\text{iii}}$	0.93	2.19	3.119 (2)	174
$\text{O2}-\text{H22}\cdots\text{Cl2}^{\text{vi}}$	0.86	2.31	3.1590 (18)	168

Symmetry codes: (i)  $x-1, y-1, z$ ; (ii)  $x-1, y, z$ ; (iii)  $x, y-1, z$ ; (iv)  $-x+2, y+\frac{1}{2}, -z+1$ ; (v)  $x, y+1, z$ ; (vi)  $-x+2, y-\frac{1}{2}, -z+2$ .

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This work was supported by a start-up grant from Anyang Institute of Technology, China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5019).

### References

- Brammer, L., Bruton, E. A. & Sherwood, P. (2001). *Cryst. Growth Des.* **1**, 277–290.
- Bremner, C. A. & Harrison, W. T. A. (2003). *Acta Cryst.* **E59**, m425–m426.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Kefi, R. & Nasr, C. B. (2005). *Z. Kristallogr. New Cryst. Struct.* **220**, 241.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sutherland, P. A. & Harrison, W. T. A. (2009). *Acta Cryst.* **E65**, m565.

**supplementary materials**

*Acta Cryst.* (2010). E66, m1224 [ doi:10.1107/S1600536810035506 ]

## (*R*)-2-Methylpiperazine-1,4-dium diaquatetrachloridoferrate(II)

C.-H. Peng and Y.-P. Li

### Comment

Recently much attention has been devoted to hydrogen bonding networks in molecular salts containing metal-chlorido complexes (Brammer *et al.*, 2001; Bremner & Harrison, 2003; Kefi & Nasr, 2005). The crystal structure of piperazinedium tetrachloridozincate(II) has been reported (Sutherland & Harrison, 2009). The construction of new members of this family is an important direction in the development of coordination chemistry. We report here the crystal structure of the title compound.

The crystal structure of the title compound (Fig. 1) contains the protonated piperazindium cations and trans-Fe(H<sub>2</sub>O)<sub>2</sub>Cl<sub>4</sub> octahedral anions. The piperazine ring adopts a chair conformation. An extensive network of N—H···Cl, N—H···(Cl,Cl) and O—H···Cl hydrogen bonds results in a structure with a three-dimensional hydrogen-bond network (Fig. 2).

### Experimental

(*R*)-2-Methylpiperazine (2 mmol, 0.2 g), FeCl<sub>3</sub> (2 mmol, 0.31 g), KI (1 mmol, 0.17), I<sub>2</sub> (0.5 mmol, 0.13 g) and 5% aqueous HCl (5 ml) were dissolved in 10 ml water, the solution was heated to 353 K (0.5 h), forming a clear solution. The reaction mixture was cooled slowly to room temperature, crystals of the title compound were formed after 6 d.

### Refinement

Water H atoms were located in a difference Fourier map and refined as riding their as found relative positions with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ . Other H atoms were placed in calculated positions with C—H = 0.9 or 0.98 and N—H = 0.90 Å, and refined using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$ .

### Figures




Fig. 1. The asymmetric unit of the title compound with atom labels. Displacement ellipsoids were drawn at the 30% probability level




Fig. 2. The packing viewed along the *a* axis. Hydrogen bonds are drawn as dashed lines

## (R)-2-Methylpiperazine-1,4-dium diaquatetrachloridoferrate(II)

### Crystal data

$(C_5H_{14}N_2)[FeCl_4(H_2O)_2]$	$F(000) = 344$
$M_r = 335.86$	$D_x = 1.707 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2yb	Cell parameters from 2456 reflections
$a = 8.6013 (17) \text{ \AA}$	$\theta = 3.2\text{--}26.0^\circ$
$b = 6.4495 (13) \text{ \AA}$	$\mu = 1.95 \text{ mm}^{-1}$
$c = 12.024 (2) \text{ \AA}$	$T = 291 \text{ K}$
$\beta = 101.64 (3)^\circ$	Block, yellow
$V = 653.3 (2) \text{ \AA}^3$	$0.28 \times 0.24 \times 0.20 \text{ mm}$
$Z = 2$	

### Data collection

Rigaku SCXmini diffractometer	2558 independent reflections
Radiation source: fine-focus sealed tube graphite	2456 reflections with $I > 2\sigma(I)$
Detector resolution: $13.6612 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.025$
$\omega$ scans	$\theta_{\text{max}} = 26.0^\circ$ , $\theta_{\text{min}} = 3.2^\circ$
Absorption correction: multi-scan ( <i>CrystalClear</i> ; Rigaku, 2005)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.8$ , $T_{\text{max}} = 0.9$	$k = -7 \rightarrow 7$
6105 measured reflections	$l = -14 \rightarrow 14$

### Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.023$	$w = 1/[\sigma^2(F_o^2) + (0.0172P)^2 + 0.0801P]$
$wR(F^2) = 0.050$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.08$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2558 reflections	$\Delta\rho_{\text{max}} = 0.19 \text{ e \AA}^{-3}$
129 parameters	$\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.116 (2)
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 1156 Friedel pairs
	Flack parameter: 0.010 (14)

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.92584 (4)	0.85265 (6)	0.73702 (3)	0.02261 (10)
Cl1	0.63527 (7)	0.89798 (9)	0.69296 (5)	0.03230 (17)
Cl2	0.95176 (8)	1.16607 (10)	0.86527 (5)	0.03236 (16)
Cl3	0.88976 (8)	0.53109 (10)	0.60858 (5)	0.03094 (16)
Cl4	1.21704 (8)	0.81475 (10)	0.77621 (7)	0.0463 (2)
N1	0.2717 (2)	0.3313 (4)	0.75683 (15)	0.0271 (5)
H1C	0.1922	0.2493	0.7690	0.033*
H1D	0.2396	0.4639	0.7587	0.033*
N2	0.5845 (2)	0.3786 (4)	0.71209 (16)	0.0309 (5)
H2C	0.6147	0.2453	0.7103	0.037*
H2D	0.6648	0.4588	0.6994	0.037*
O1	0.9382 (2)	1.0515 (3)	0.59861 (15)	0.0433 (5)
H11	0.9818	1.0347	0.5446	0.065*
H12	0.9188	1.1870	0.6013	0.065*
O2	0.91049 (19)	0.6462 (3)	0.87073 (13)	0.0303 (4)
H21	0.9234	0.5037	0.8634	0.045*
H22	0.9545	0.6697	0.9406	0.045*
C1	0.4148 (3)	0.2974 (4)	0.84969 (18)	0.0242 (5)
H1A	0.4447	0.1506	0.8508	0.029*
C2	0.5510 (3)	0.4269 (4)	0.82553 (19)	0.0271 (6)
H2A	0.5248	0.5727	0.8289	0.032*
H2B	0.6452	0.4003	0.8833	0.032*
C3	0.4415 (3)	0.4152 (5)	0.6213 (2)	0.0339 (6)
H3A	0.4649	0.3796	0.5480	0.041*
H3B	0.4127	0.5607	0.6198	0.041*
C4	0.3052 (3)	0.2856 (4)	0.64264 (19)	0.0317 (6)
H4A	0.2115	0.3146	0.5849	0.038*
H4B	0.3308	0.1398	0.6379	0.038*
C5	0.3760 (3)	0.3539 (5)	0.9633 (2)	0.0399 (6)
H5A	0.3347	0.4926	0.9600	0.060*
H5B	0.4705	0.3458	1.0213	0.060*
H5C	0.2981	0.2593	0.9807	0.060*

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Fe1	0.02514 (17)	0.01809 (17)	0.02560 (17)	-0.00031 (12)	0.00748 (12)	0.00170 (13)
Cl1	0.0251 (3)	0.0260 (4)	0.0452 (4)	0.0010 (2)	0.0056 (2)	-0.0016 (3)
Cl2	0.0461 (4)	0.0235 (3)	0.0255 (3)	0.0002 (3)	0.0024 (3)	-0.0033 (3)
Cl3	0.0468 (4)	0.0231 (3)	0.0269 (3)	0.0018 (3)	0.0171 (3)	-0.0002 (3)
Cl4	0.0270 (4)	0.0260 (5)	0.0866 (6)	0.0004 (3)	0.0131 (3)	0.0033 (4)
N1	0.0224 (10)	0.0257 (12)	0.0326 (11)	-0.0024 (9)	0.0039 (8)	0.0032 (10)
N2	0.0295 (11)	0.0285 (13)	0.0388 (11)	-0.0037 (10)	0.0163 (8)	-0.0018 (11)
O1	0.0727 (14)	0.0263 (10)	0.0418 (11)	0.0091 (10)	0.0376 (10)	0.0080 (9)
O2	0.0425 (10)	0.0245 (10)	0.0224 (8)	-0.0018 (8)	0.0033 (7)	0.0017 (8)
C1	0.0243 (12)	0.0214 (14)	0.0249 (12)	-0.0003 (10)	0.0005 (9)	0.0041 (10)
C2	0.0235 (12)	0.0272 (15)	0.0300 (13)	-0.0037 (11)	0.0044 (10)	-0.0014 (11)
C3	0.0438 (15)	0.0347 (16)	0.0247 (12)	-0.0036 (12)	0.0106 (10)	0.0005 (11)
C4	0.0350 (14)	0.0298 (15)	0.0273 (13)	-0.0028 (11)	-0.0007 (10)	-0.0031 (11)
C5	0.0432 (15)	0.0486 (17)	0.0298 (13)	-0.0074 (14)	0.0122 (11)	0.0019 (13)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Fe1—O2	2.1122 (17)	O2—H21	0.9315
Fe1—O1	2.1205 (18)	O2—H22	0.8622
Fe1—Cl1	2.4654 (9)	C1—C2	1.514 (3)
Fe1—Cl4	2.4655 (8)	C1—C5	1.515 (3)
Fe1—Cl2	2.5249 (9)	C1—H1A	0.9800
Fe1—Cl3	2.5669 (8)	C2—H2A	0.9700
N1—C4	1.488 (3)	C2—H2B	0.9700
N1—C1	1.502 (3)	C3—C4	1.503 (3)
N1—H1C	0.9000	C3—H3A	0.9700
N1—H1D	0.9000	C3—H3B	0.9700
N2—C2	1.483 (3)	C4—H4A	0.9700
N2—C3	1.490 (3)	C4—H4B	0.9700
N2—H2C	0.9000	C5—H5A	0.9600
N2—H2D	0.9000	C5—H5B	0.9600
O1—H11	0.8195	C5—H5C	0.9600
O1—H12	0.8919		
O2—Fe1—O1	177.97 (9)	H21—O2—H22	103.2
O2—Fe1—Cl1	91.26 (5)	N1—C1—C2	109.07 (19)
O1—Fe1—Cl1	88.22 (6)	N1—C1—C5	109.81 (19)
O2—Fe1—Cl4	90.52 (5)	C2—C1—C5	111.1 (2)
O1—Fe1—Cl4	90.00 (6)	N1—C1—H1A	108.9
Cl1—Fe1—Cl4	178.22 (3)	C2—C1—H1A	108.9
O2—Fe1—Cl2	92.94 (5)	C5—C1—H1A	108.9
O1—Fe1—Cl2	89.03 (6)	N2—C2—C1	111.2 (2)
Cl1—Fe1—Cl2	89.83 (3)	N2—C2—H2A	109.4
Cl4—Fe1—Cl2	90.09 (3)	C1—C2—H2A	109.4
O2—Fe1—Cl3	85.99 (5)	N2—C2—H2B	109.4

O1—Fe1—C13	92.03 (6)	C1—C2—H2B	109.4
C11—Fe1—C13	88.41 (3)	H2A—C2—H2B	108.0
C14—Fe1—C13	91.70 (3)	N2—C3—C4	110.1 (2)
C12—Fe1—C13	177.92 (3)	N2—C3—H3A	109.6
C4—N1—C1	112.03 (19)	C4—C3—H3A	109.6
C4—N1—H1C	109.2	N2—C3—H3B	109.6
C1—N1—H1C	109.2	C4—C3—H3B	109.6
C4—N1—H1D	109.2	H3A—C3—H3B	108.2
C1—N1—H1D	109.2	N1—C4—C3	110.5 (2)
H1C—N1—H1D	107.9	N1—C4—H4A	109.5
C2—N2—C3	110.8 (2)	C3—C4—H4A	109.5
C2—N2—H2C	109.5	N1—C4—H4B	109.5
C3—N2—H2C	109.5	C3—C4—H4B	109.5
C2—N2—H2D	109.5	H4A—C4—H4B	108.1
C3—N2—H2D	109.5	C1—C5—H5A	109.5
H2C—N2—H2D	108.1	C1—C5—H5B	109.5
Fe1—O1—H11	130.2	H5A—C5—H5B	109.5
Fe1—O1—H12	121.6	C1—C5—H5C	109.5
H11—O1—H12	106.1	H5A—C5—H5C	109.5
Fe1—O2—H21	121.5	H5B—C5—H5C	109.5
Fe1—O2—H22	123.2		
C4—N1—C1—C2	55.9 (3)	C5—C1—C2—N2	-177.4 (2)
C4—N1—C1—C5	177.8 (2)	C2—N2—C3—C4	-57.9 (3)
C3—N2—C2—C1	58.3 (3)	C1—N1—C4—C3	-56.9 (3)
N1—C1—C2—N2	-56.3 (3)	N2—C3—C4—N1	56.9 (3)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1C $\cdots$ C12 <sup>i</sup>	0.90	2.62	3.443 (2)	152
N1—H1C $\cdots$ C14 <sup>i</sup>	0.90	2.81	3.379 (3)	122
N1—H1D $\cdots$ C14 <sup>ii</sup>	0.90	2.28	3.169 (3)	167
N2—H2C $\cdots$ C11 <sup>iii</sup>	0.90	2.26	3.145 (3)	168
N2—H2D $\cdots$ C13	0.90	2.45	3.275 (2)	152
O1—H11 $\cdots$ C13 <sup>iv</sup>	0.82	2.33	3.147 (2)	173
O1—H12 $\cdots$ C13 <sup>v</sup>	0.89	2.24	3.127 (2)	176
O2—H21 $\cdots$ C12 <sup>iii</sup>	0.93	2.19	3.119 (2)	174
O2—H22 $\cdots$ C12 <sup>vi</sup>	0.86	2.31	3.1590 (18)	168

Symmetry codes: (i)  $x-1, y-1, z$ ; (ii)  $x-1, y, z$ ; (iii)  $x, y-1, z$ ; (iv)  $-x+2, y+1/2, -z+1$ ; (v)  $x, y+1, z$ ; (vi)  $-x+2, y-1/2, -z+2$ .

Fig. 1






Fig. 2

