

7-(1,3-Dioxolan-2-ylmethyl)-1,3-dimethyl-2,6-dioxo-2,3,6,7-tetrahydro-1H-purin-9-ium tetrachloridoferrate(III)

Ping Ping Wang,^a Jiang Gong^{b,c} and Shi-Feng Ni^{c*}

^aCollege of Pharmaceutical Sciences, Zhejiang University of Technology, Hangzhou 310014, People's Republic of China, ^bDepartment of Medicine, Tibet Nationalities Institute, Xianyang, Shaanxi 712082, People's Republic of China, and ^cKey Laboratory of Resource Biology and Biotechnology in Western China, Ministry of Education, College of Life Science, Northwest University, Xi'an 710069, People's Republic of China

Correspondence e-mail: nshyfe@sina.cn


Received 20 July 2010; accepted 28 August 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.013$ Å; disorder in main residue; R factor = 0.099; wR factor = 0.207; data-to-parameter ratio = 14.0.

The asymmetric unit of the title compound, $(\text{C}_{11}\text{H}_{15}\text{N}_4\text{O}_4)\text{[FeCl}_4\text{]}$, contains two independent protonated 7-(1,3-dioxolan-2-ylmethyl)-3,7-dihydro-1,3-dimethyl-1H-purine-2,6-dione (doxofyllinium) and two tetrahedral tetrachloridoferrate(III) anions. In the doxofyllinium, two disordered methylene C atoms are observed in each dioxolane ring with an occupancy ratio of 0.54 (4):0.46 (4). In the crystal, molecules are connected by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds and weak $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{Cl}$ interactions.

Related literature

For the biological activity of the drug doxofylline, see: Franzzone *et al.* (1981, 1989); Zhao & Li (2001). For bond distances and angles in other tetrachloridoferrate(III) compounds, see: Barbaro *et al.* (1992); Bottomley *et al.* (1984). For the synthesis of doxofylline, see: Li *et al.* (1995).



Experimental

Crystal data

$(\text{C}_{11}\text{H}_{15}\text{N}_4\text{O}_4)\text{[FeCl}_4\text{]}$
 $M_r = 464.92$
 Tetragonal, $P4_2$
 $a = 20.2947$ (4) Å

$c = 9.0692$ (4) Å
 $V = 3735.38$ (19) Å³
 $Z = 8$
 Mo $K\alpha$ radiation

$\mu = 1.40$ mm⁻¹
 $T = 293$ K

0.27 × 0.16 × 0.13 mm

Data collection

Bruker SMART APEX area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{\min} = 0.775$, $T_{\max} = 0.852$

19874 measured reflections
 6253 independent reflections
 5589 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.099$
 $wR(F^2) = 0.207$
 $S = 1.30$
 6253 reflections
 447 parameters
 13 restraints

H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.49$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.31$ e Å⁻³
 Absolute structure: Flack (1983),
 2724 Friedel pairs
 Flack parameter: 0.10 (4)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N4}-\text{H4}\cdots\text{O2}^{\text{i}}$	0.86	1.93	2.774 (9)	165
$\text{N8}-\text{H8}\cdots\text{O6}^{\text{ii}}$	0.86	1.91	2.754 (9)	166
$\text{C5}-\text{H5}\cdots\text{Cl5}^{\text{i}}$	0.93	2.80	3.650 (11)	153
$\text{C8}-\text{H8B}\cdots\text{O7}^{\text{iii}}$	0.97	2.29	3.114 (11)	142
$\text{C10}-\text{H10C}\cdots\text{O5}^{\text{iv}}$	0.95	2.46	3.098 (11)	124
$\text{C10}'-\text{H10B}\cdots\text{O5}^{\text{iv}}$	0.89	2.57	3.384 (11)	153
$\text{C10}''-\text{H10C}\cdots\text{O5}^{\text{iv}}$	0.97	2.46	3.384 (11)	159
$\text{C16}-\text{H16}\cdots\text{Cl1}^{\text{iii}}$	0.93	2.78	3.560 (11)	143
$\text{C19}-\text{H19A}\cdots\text{O4}^{\text{v}}$	0.97	2.48	3.227 (11)	134
$\text{C19}-\text{H19A}\cdots\text{O5}$	0.97	2.53	3.167 (11)	123
$\text{C22}'-\text{H22D}\cdots\text{O1}$	0.97	2.58	3.242 (11)	126

Symmetry codes: (i) $-y + 1, x, z + \frac{1}{2}$; (ii) $y, -x, z + \frac{1}{2}$; (iii) $-x + 1, -y, z$; (iv) $x, y, z + 1$; (v) $-x + 1, -y, z - 1$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

We are grateful for financial support by the Natural Science Foundation of Tibet (2009–10–12) and the Natural Science Foundation of the Key Laboratory of Resource Biology and Biotechnology in Western China (Northwest University), Ministry of Education (2009–11–12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2188).

References

- Barbaro, P., Bianchini, C., Fochi, M., Masi, D. & Mealli, C. (1992). *Acta Cryst.* **C48**, 625–627.
 Bottomley, G. A., Carter, A. M., Engelhardt, L. M., Lincoln, F. J. & Patric, J. M. (1984). *Aust. J. Chem.* **37**, 871–877.
 Bruker (2000). SMART, SADABS and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Franzzone, J. S., Cirillo, R. & Biffignandi, P. (1989). *Eur. J. Pharmacol.* **165**, 269–274.
 Franzzone, J. S., Reboani, C. & Fonzo, D. (1981). *Farmacol. Sci.* **36**, 201–219.
 Li, C. H., Luo, Z. & Li, Z. H. (1995). *Chin. J. Pharm.* **25**, 385–386.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Zhao, J. J. & Li, L. (2001). *J. Norman Bethune Univ. Med. Sci.* **27**, 646–676.

supplementary materials

Acta Cryst. (2010). E66, m1247 [doi:10.1107/S1600536810034720]

7-(1,3-Dioxolan-2-ylmethyl)-1,3-dimethyl-2,6-dioxo-2,3,6,7-tetrahydro-1*H*-purin-9-ium tetrachloridoferrate(III)

P. P. Wang, J. Gong and S.-F. Ni

Comment

Doxofylline [7-(1,3-dioxolan-2-ylmethyl)-1,3-dimethyl-3,7-dihydro-1*H*-purine-2,6-dione] is a xanthine drug with anti-asthmatic (Franzone *et al.*, 1989), antiinflammatory (Zhao *et al.*, 2001), and bronchodilating activities (Franzone *et al.*, 1981). Now we present here the structure of the title compound, (I).

In the title compound (Fig. 1), (I), the asymmetric unit contain two crystallographically independent molecules of doxofyllinium tetrachloroferrate(III). The furan rings of the doxofyllinium are disordered, and the five atoms of these rings aren't coplanar. The iron cation is tetra coordinated by chlorine anions, and it adopts a slightly distorted tetrahedral coordination with two angles smaller than the tetrahedral one, two almost equal to tetrahedral and two larger than tetrahedral (Table 1). Fe—Cl distances spanning the range 2.172 (3) Å–2.190 (4) Å, and Cl—Fe—Cl angles 107.41 (16)°–111.77 (15)°, are similar to those found in other tetrachloroferrate(III) (Bottomley *et al.*, 1984; Barbaro *et al.*, 1992).

In the crystal, doxofyllinium cations are linked by N8—H8ⁱ⋯O6ⁱ and N4—H4ⁱⁱ⋯O2ⁱⁱ hydrogen bonds (Table 2). The weak C—Hⁱⁱⁱ⋯O and C—H^{iv}⋯Cl interactions further link (I), reinforcing the structure (Table 2).

Experimental

Doxofylline was synthesized according to Li *et al.* (1995). Doxofylline, hydrochloric acid and trichloride were dissolved in sufficient ethanol by heating to 333 K, where a yellow solution resulted. Crystals of (I) were formed by gradual evaporation of ethanol over a period of one week at 293 K.

Refinement

All of the H atoms were placed in calculated positions and allowed to ride on their parent atoms at distances of 0.93 (C5—H5), 0.96 (methyl), 0.97 (methylene) and 0.98 Å (methine), with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5 U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. A view of the asymmetric unit of (I) with atom labels, showing 40% probability displacement ellipsoids. Parts of disorder furan rings are omitted for clarity.

supplementary materials

7-(1,3-Dioxolan-2-ylmethyl)-1,3-dimethyl-2,6-dioxo-2,3,6,7-tetrahydro- 1H-purin-9-ium tetrachloridoferrate(III)

Crystal data

(C ₁₁ H ₁₅ N ₄ O ₄)[FeCl ₄]	$D_x = 1.653 \text{ Mg m}^{-3}$
$M_r = 464.92$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Tetragonal, $P4_2$	Cell parameters from 8917 reflections
Hall symbol: P 4c	$\theta = 2.2\text{--}26.5^\circ$
$a = 20.2947 (4) \text{ \AA}$	$\mu = 1.40 \text{ mm}^{-1}$
$c = 9.0692 (4) \text{ \AA}$	$T = 293 \text{ K}$
$V = 3735.38 (19) \text{ \AA}^3$	Block, colourless
$Z = 8$	$0.27 \times 0.16 \times 0.13 \text{ mm}$
$F(000) = 1880.0$	

Data collection

Bruker SMART APEX area-detector diffractometer	6253 independent reflections
Radiation source: fine-focus sealed tube graphite	5589 reflections with $I > 2\sigma(I)$
φ and ω scan	$R_{\text{int}} = 0.046$
Absorption correction: multi-scan (SADABS; Bruker, 2000)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 1.0^\circ$
$T_{\text{min}} = 0.775$, $T_{\text{max}} = 0.852$	$h = -24 \rightarrow 23$
19874 measured reflections	$k = -24 \rightarrow 23$
	$l = -9 \rightarrow 10$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.099$	H-atom parameters constrained
$wR(F^2) = 0.207$	$w = 1/[\sigma^2(F_o^2) + (0.0665P)^2 + 7.3566P]$
$S = 1.30$	where $P = (F_o^2 + 2F_c^2)/3$
6253 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
447 parameters	$\Delta\rho_{\text{max}} = 0.49 \text{ e \AA}^{-3}$
13 restraints	$\Delta\rho_{\text{min}} = -0.31 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 2724 Friedel pairs
	Flack parameter: 0.10 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Fe1	0.68089 (8)	0.33071 (7)	0.16159 (18)	0.0569 (4)	
Fe2	0.16760 (8)	0.17898 (7)	0.22058 (17)	0.0586 (4)	
Cl1	0.69861 (18)	0.24505 (16)	0.2998 (4)	0.0869 (10)	
Cl2	0.59364 (17)	0.3131 (2)	0.0264 (4)	0.0898 (11)	
Cl3	0.76550 (17)	0.35346 (18)	0.0238 (4)	0.0913 (11)	
Cl4	0.6604 (2)	0.41656 (17)	0.3005 (4)	0.0913 (11)	
Cl5	0.17996 (16)	0.25893 (15)	0.3795 (4)	0.0735 (9)	
Cl6	0.1499 (2)	0.08671 (15)	0.3379 (4)	0.0886 (10)	
Cl7	0.08268 (18)	0.20197 (19)	0.0809 (5)	0.0966 (11)	
Cl8	0.25566 (19)	0.1697 (2)	0.0873 (5)	0.1016 (12)	
O1	0.5196 (4)	0.1673 (3)	0.2934 (8)	0.0592 (19)	
O2	0.4364 (3)	0.3755 (3)	0.2618 (9)	0.063 (2)	
O3	0.4982 (4)	0.1135 (4)	0.6244 (10)	0.086 (3)	
O4	0.5648 (4)	0.1321 (4)	0.8177 (8)	0.072 (2)	
O5	0.3355 (3)	0.0243 (4)	-0.1686 (8)	0.0587 (19)	
O6	0.1189 (4)	0.0815 (4)	-0.2061 (9)	0.068 (2)	
O7	0.3705 (4)	-0.0319 (3)	0.3520 (8)	0.063 (2)	
O8	0.3889 (4)	0.0414 (4)	0.1734 (9)	0.073 (2)	
N1	0.4750 (4)	0.2704 (4)	0.2853 (9)	0.0448 (19)	
N2	0.5058 (4)	0.3546 (3)	0.4494 (9)	0.044 (2)	
N3	0.5913 (3)	0.2156 (3)	0.5737 (8)	0.0370 (16)	
N4	0.5845 (4)	0.3180 (4)	0.6382 (9)	0.0450 (19)	
H4	0.5907	0.3539	0.6865	0.054*	
N5	0.2260 (4)	0.0565 (4)	-0.1796 (9)	0.0464 (19)	
N6	0.1468 (4)	0.0128 (4)	-0.0221 (10)	0.047 (2)	
N7	0.2938 (3)	-0.0580 (3)	0.1018 (8)	0.0343 (16)	
N8	0.1903 (4)	-0.0635 (3)	0.1613 (9)	0.0452 (19)	
H8	0.1551	-0.0744	0.2081	0.054*	
C1	0.5154 (4)	0.2206 (4)	0.3469 (11)	0.040 (2)	
C2	0.4696 (4)	0.3359 (5)	0.3274 (12)	0.046 (2)	
C3	0.5443 (4)	0.3099 (3)	0.5178 (10)	0.0317 (19)	
C4	0.5473 (4)	0.2455 (4)	0.4781 (10)	0.039 (2)	
C5	0.6115 (4)	0.2604 (4)	0.6661 (11)	0.040 (2)	
H5	0.6413	0.2526	0.7420	0.048*	
C6	0.4343 (6)	0.2508 (6)	0.1589 (13)	0.068 (3)	
H6A	0.4414	0.2050	0.1380	0.103*	
H6B	0.4463	0.2766	0.0744	0.103*	
H6C	0.3887	0.2581	0.1817	0.103*	

supplementary materials

C7	0.5043 (5)	0.4240 (5)	0.4975 (16)	0.068 (3)	
H7A	0.4743	0.4483	0.4364	0.102*	
H7B	0.5476	0.4426	0.4894	0.102*	
H7C	0.4899	0.4262	0.5983	0.102*	
C8	0.6086 (5)	0.1450 (4)	0.5793 (11)	0.049 (2)	
H8A	0.6530	0.1401	0.6176	0.059*	
H8B	0.6077	0.1268	0.4805	0.059*	
C9	0.5614 (6)	0.1078 (5)	0.6754 (12)	0.060 (3)	
H9	0.5739	0.0611	0.6762	0.072*	
C10	0.4622 (13)	0.1031 (15)	0.7592 (18)	0.068 (8)	0.46 (4)
H10A	0.4659	0.0580	0.7939	0.081*	0.46 (4)
H10B	0.4160	0.1147	0.7488	0.081*	0.46 (4)
C11	0.4989 (6)	0.1514 (14)	0.859 (3)	0.068 (8)	0.46 (4)
H11A	0.4895	0.1969	0.8345	0.081*	0.46 (4)
H11B	0.4901	0.1436	0.9627	0.081*	0.46 (4)
C10'	0.4559 (11)	0.1331 (16)	0.7455 (19)	0.094 (10)	0.54 (4)
H10C	0.4157	0.1072	0.7479	0.113*	0.54 (4)
H10D	0.4448	0.1795	0.7401	0.113*	0.54 (4)
C11'	0.4999 (7)	0.1183 (18)	0.878 (3)	0.094 (10)	0.54 (4)
H11C	0.4898	0.1469	0.9609	0.113*	0.54 (4)
H11D	0.4960	0.0727	0.9091	0.113*	0.54 (4)
C12	0.2815 (5)	0.0217 (4)	-0.1225 (11)	0.043 (2)	
C13	0.1621 (4)	0.0531 (4)	-0.1418 (12)	0.044 (2)	
C14	0.1953 (4)	-0.0211 (3)	0.0451 (10)	0.0309 (19)	
C15	0.2591 (4)	-0.0156 (4)	0.0084 (10)	0.035 (2)	
C16	0.2515 (5)	-0.0846 (4)	0.1871 (11)	0.042 (2)	
H16	0.2622	-0.1153	0.2593	0.050*	
C17	0.2421 (6)	0.1030 (6)	-0.3009 (13)	0.080 (4)	
H17A	0.2887	0.1019	-0.3196	0.119*	
H17B	0.2294	0.1468	-0.2731	0.119*	
H17C	0.2188	0.0902	-0.3885	0.119*	
C18	0.0772 (5)	0.0074 (6)	0.0225 (16)	0.071 (4)	
H18A	0.0507	0.0353	-0.0389	0.107*	
H18B	0.0726	0.0207	0.1235	0.107*	
H18C	0.0629	-0.0375	0.0119	0.107*	
C19	0.3647 (4)	-0.0690 (5)	0.1035 (11)	0.045 (2)	
H19A	0.3825	-0.0610	0.0058	0.054*	
H19B	0.3736	-0.1145	0.1292	0.054*	
C20	0.3981 (4)	-0.0246 (5)	0.2117 (13)	0.054 (3)	
H20	0.4453	-0.0347	0.2153	0.065*	
C22	0.367 (2)	0.0769 (14)	0.303 (2)	0.065 (11)	0.36 (4)
H22A	0.3207	0.0863	0.2988	0.078*	0.36 (4)
H22B	0.3915	0.1179	0.3141	0.078*	0.36 (4)
C21	0.383 (3)	0.0292 (8)	0.429 (3)	0.065 (11)	0.36 (4)
H21A	0.4289	0.0327	0.4606	0.078*	0.36 (4)
H21B	0.3543	0.0353	0.5127	0.078*	0.36 (4)
C22'	0.3901 (12)	0.0742 (11)	0.3145 (17)	0.078 (7)	0.64 (4)
H22C	0.3718	0.1183	0.3076	0.094*	0.64 (4)
H22D	0.4346	0.0769	0.3530	0.094*	0.64 (4)

C21'	0.3470 (14)	0.0299 (6)	0.411 (3)	0.078 (7)	0.64 (4)
H21C	0.3566	0.0348	0.5150	0.094*	0.64 (4)
H21D	0.3004	0.0365	0.3931	0.094*	0.64 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0695 (10)	0.0601 (10)	0.0411 (9)	-0.0070 (8)	0.0035 (8)	0.0009 (8)
Fe2	0.0745 (11)	0.0581 (10)	0.0432 (9)	0.0120 (8)	0.0031 (8)	0.0023 (8)
Cl1	0.105 (2)	0.078 (2)	0.078 (2)	0.0085 (18)	0.021 (2)	0.0265 (19)
Cl2	0.089 (2)	0.126 (3)	0.055 (2)	-0.026 (2)	-0.0054 (17)	-0.0144 (19)
Cl3	0.083 (2)	0.096 (2)	0.095 (3)	-0.0082 (18)	0.027 (2)	0.032 (2)
Cl4	0.128 (3)	0.082 (2)	0.064 (2)	0.003 (2)	-0.008 (2)	-0.0211 (18)
Cl5	0.087 (2)	0.0716 (18)	0.0619 (19)	-0.0058 (16)	0.0210 (17)	-0.0146 (15)
Cl6	0.136 (3)	0.0681 (19)	0.061 (2)	0.0021 (19)	-0.004 (2)	0.0153 (16)
Cl7	0.101 (3)	0.108 (3)	0.081 (2)	0.023 (2)	-0.027 (2)	0.010 (2)
Cl8	0.102 (3)	0.129 (3)	0.073 (2)	0.011 (2)	0.027 (2)	-0.018 (2)
O1	0.090 (5)	0.040 (4)	0.047 (4)	-0.004 (3)	-0.018 (4)	-0.019 (3)
O2	0.045 (4)	0.064 (4)	0.079 (6)	0.019 (3)	-0.012 (4)	0.022 (4)
O3	0.093 (6)	0.087 (6)	0.078 (7)	-0.033 (5)	-0.019 (5)	0.002 (5)
O4	0.086 (5)	0.093 (6)	0.036 (4)	0.012 (4)	-0.006 (4)	0.013 (4)
O5	0.035 (4)	0.097 (5)	0.044 (4)	-0.002 (3)	0.015 (3)	0.027 (4)
O6	0.065 (5)	0.068 (5)	0.071 (6)	0.020 (4)	-0.016 (4)	0.017 (4)
O7	0.098 (6)	0.055 (4)	0.034 (4)	-0.003 (4)	-0.015 (4)	0.011 (3)
O8	0.081 (5)	0.066 (5)	0.073 (6)	-0.024 (4)	0.002 (5)	0.009 (4)
N1	0.049 (4)	0.055 (5)	0.031 (4)	-0.003 (4)	-0.008 (4)	0.005 (4)
N2	0.050 (5)	0.024 (4)	0.056 (6)	0.008 (3)	-0.005 (4)	0.010 (3)
N3	0.048 (4)	0.041 (4)	0.022 (4)	0.009 (3)	-0.005 (3)	0.004 (3)
N4	0.061 (5)	0.036 (4)	0.038 (5)	0.003 (3)	-0.021 (4)	-0.014 (3)
N5	0.061 (5)	0.047 (4)	0.032 (4)	0.010 (4)	-0.010 (4)	0.009 (4)
N6	0.035 (4)	0.050 (5)	0.056 (5)	0.005 (3)	0.005 (4)	0.015 (4)
N7	0.037 (4)	0.039 (4)	0.027 (4)	0.003 (3)	0.002 (3)	0.012 (3)
N8	0.040 (4)	0.046 (4)	0.049 (5)	-0.010 (3)	0.013 (4)	0.000 (4)
C1	0.033 (5)	0.047 (5)	0.040 (6)	-0.003 (4)	0.004 (4)	0.010 (5)
C2	0.033 (5)	0.060 (6)	0.044 (6)	0.005 (4)	-0.009 (4)	0.014 (5)
C3	0.047 (5)	0.015 (4)	0.033 (5)	-0.012 (3)	0.009 (4)	-0.004 (3)
C4	0.038 (5)	0.052 (6)	0.026 (5)	-0.007 (4)	0.002 (4)	0.007 (4)
C5	0.052 (5)	0.035 (5)	0.033 (5)	0.004 (4)	-0.022 (4)	0.004 (4)
C6	0.088 (8)	0.071 (7)	0.046 (7)	0.008 (6)	-0.020 (6)	-0.001 (6)
C7	0.060 (7)	0.044 (6)	0.101 (10)	0.008 (5)	-0.027 (7)	0.001 (6)
C8	0.065 (6)	0.052 (6)	0.030 (5)	0.018 (5)	-0.010 (5)	0.003 (4)
C9	0.096 (9)	0.045 (6)	0.039 (6)	0.006 (5)	-0.021 (6)	0.010 (5)
C10	0.057 (13)	0.060 (14)	0.085 (17)	0.011 (9)	-0.009 (10)	0.043 (12)
C11	0.057 (13)	0.060 (14)	0.085 (17)	0.011 (9)	-0.009 (10)	0.043 (12)
C10'	0.083 (15)	0.061 (16)	0.14 (2)	0.025 (11)	0.014 (14)	0.054 (14)
C11'	0.083 (15)	0.061 (16)	0.14 (2)	0.025 (11)	0.014 (14)	0.054 (14)
C12	0.053 (6)	0.035 (5)	0.041 (6)	-0.004 (4)	-0.007 (5)	0.005 (4)
C13	0.026 (4)	0.046 (5)	0.060 (7)	0.017 (4)	-0.014 (5)	-0.009 (5)

supplementary materials

C14	0.037 (5)	0.019 (4)	0.037 (5)	-0.008 (3)	0.006 (4)	0.007 (3)
C15	0.048 (5)	0.025 (4)	0.032 (5)	0.003 (4)	0.002 (4)	0.002 (4)
C16	0.061 (6)	0.021 (4)	0.043 (6)	0.007 (4)	-0.008 (5)	0.018 (4)
C17	0.096 (9)	0.094 (9)	0.049 (8)	0.012 (7)	0.012 (7)	0.020 (7)
C18	0.038 (6)	0.072 (7)	0.103 (10)	0.003 (5)	0.003 (6)	0.015 (7)
C19	0.035 (5)	0.057 (6)	0.042 (6)	0.019 (4)	-0.008 (4)	0.007 (5)
C20	0.034 (5)	0.059 (6)	0.069 (8)	0.011 (4)	0.003 (5)	0.015 (6)
C22	0.06 (2)	0.051 (16)	0.08 (2)	-0.008 (12)	0.016 (14)	0.002 (14)
C21	0.06 (2)	0.051 (16)	0.08 (2)	-0.008 (12)	0.016 (14)	0.002 (14)
C22'	0.063 (12)	0.064 (11)	0.108 (15)	-0.001 (7)	-0.001 (9)	-0.009 (10)
C21'	0.063 (12)	0.064 (11)	0.108 (15)	-0.001 (7)	-0.001 (9)	-0.009 (10)

Geometric parameters (Å, °)

Fe1—C11	2.173 (3)	N8—H8	0.8600
Fe1—C13	2.173 (3)	C1—C4	1.446 (13)
Fe1—C12	2.183 (4)	C3—C4	1.357 (12)
Fe1—C14	2.190 (4)	C5—H5	0.9300
Fe2—C18	2.165 (4)	C6—H6A	0.9600
Fe2—C16	2.183 (3)	C6—H6B	0.9600
Fe2—C15	2.185 (3)	C6—H6C	0.9600
Fe2—C17	2.189 (4)	C7—H7A	0.9600
O1—C1	1.189 (10)	C7—H7B	0.9600
O2—C2	1.206 (11)	C7—H7C	0.9600
O3—C9	1.368 (13)	C8—C9	1.499 (14)
O3—C10	1.441 (10)	C8—H8A	0.9700
O3—C10'	1.450 (10)	C8—H8B	0.9700
O4—C9	1.383 (13)	C9—H9	0.9800
O4—C11	1.442 (10)	C10—C11	1.528 (10)
O4—C11'	1.454 (10)	C10—H10A	0.9700
O5—C12	1.173 (10)	C10—H10B	0.9700
O6—C13	1.201 (10)	C11—H11A	0.9700
O7—C20	1.398 (13)	C11—H11B	0.9700
O7—C21'	1.444 (9)	C10'—C11'	1.529 (10)
O7—C21	1.447 (10)	C10'—H10C	0.9700
O8—C20	1.396 (12)	C10'—H10D	0.9700
O8—C22'	1.443 (10)	C11'—H11C	0.9700
O8—C22	1.446 (10)	C11'—H11D	0.9700
N1—C2	1.386 (12)	C12—C15	1.479 (12)
N1—C1	1.417 (11)	C14—C15	1.342 (11)
N1—C6	1.467 (13)	C16—H16	0.9300
N2—C3	1.349 (11)	C17—H17A	0.9600
N2—C2	1.382 (12)	C17—H17B	0.9600
N2—C7	1.473 (12)	C17—H17C	0.9600
N3—C5	1.302 (11)	C18—H18A	0.9600
N3—C4	1.384 (11)	C18—H18B	0.9600
N3—C8	1.477 (11)	C18—H18C	0.9600
N4—C5	1.316 (10)	C19—C20	1.494 (14)
N4—C3	1.372 (11)	C19—H19A	0.9700

N4—H4	0.8600	C19—H19B	0.9700
N5—C13	1.344 (11)	C20—H20	0.9800
N5—C12	1.427 (11)	C22—C21	1.530 (10)
N5—C17	1.486 (13)	C22—H22A	0.9700
N6—C14	1.346 (11)	C22—H22B	0.9700
N6—C13	1.394 (13)	C21—H21A	0.9700
N6—C18	1.473 (12)	C21—H21B	0.9700
N7—C16	1.274 (11)	C22'—C21'	1.528 (10)
N7—C15	1.399 (10)	C22'—H22C	0.9700
N7—C19	1.456 (10)	C22'—H22D	0.9700
N8—C16	1.335 (11)	C21'—H21C	0.9700
N8—C14	1.365 (11)	C21'—H21D	0.9700
C11—Fe1—C13	111.77 (15)	O3—C10—C11	99.2 (16)
C11—Fe1—C12	109.10 (16)	O3—C10—H10A	111.9
C13—Fe1—C12	110.66 (17)	C11—C10—H10A	111.9
C11—Fe1—C14	109.65 (16)	O3—C10—H10B	111.9
C13—Fe1—C14	108.16 (16)	C11—C10—H10B	111.9
C12—Fe1—C14	107.41 (16)	H10A—C10—H10B	109.6
C18—Fe2—C16	109.43 (17)	O4—C11—C10	97.2 (16)
C18—Fe2—C15	109.77 (15)	O4—C11—H11A	112.3
C16—Fe2—C15	109.54 (15)	C10—C11—H11A	112.3
C18—Fe2—C17	110.21 (18)	O4—C11—H11B	112.3
C16—Fe2—C17	109.58 (17)	C10—C11—H11B	112.3
C15—Fe2—C17	108.28 (15)	H11A—C11—H11B	109.9
C9—O3—C10	100.2 (14)	O3—C10'—C11'	101.4 (17)
C9—O3—C10'	108.8 (12)	O3—C10'—H10C	111.5
C10—O3—C10'	25.3 (10)	C11'—C10'—H10C	111.5
C9—O4—C11	107.1 (13)	O3—C10'—H10D	111.5
C9—O4—C11'	103.8 (15)	C11'—C10'—H10D	111.5
C11—O4—C11'	27.7 (10)	H10C—C10'—H10D	109.3
C20—O7—C21'	112.1 (10)	O4—C11'—C10'	101.1 (15)
C20—O7—C21	106.0 (16)	O4—C11'—H11C	111.6
C21'—O7—C21	30.3 (14)	C10'—C11'—H11C	111.6
C20—O8—C22'	102.7 (12)	O4—C11'—H11D	111.6
C20—O8—C22	108.4 (14)	C10'—C11'—H11D	111.6
C22'—O8—C22	18.9 (19)	H11C—C11'—H11D	109.4
C2—N1—C1	128.4 (8)	O5—C12—N5	125.8 (9)
C2—N1—C6	115.5 (8)	O5—C12—C15	126.6 (8)
C1—N1—C6	116.1 (8)	N5—C12—C15	107.5 (8)
C3—N2—C2	119.4 (7)	O6—C13—N5	123.8 (10)
C3—N2—C7	121.2 (8)	O6—C13—N6	119.9 (8)
C2—N2—C7	119.3 (8)	N5—C13—N6	116.3 (7)
C5—N3—C4	107.5 (7)	C15—C14—N6	123.4 (7)
C5—N3—C8	125.4 (7)	C15—C14—N8	108.4 (7)
C4—N3—C8	126.9 (7)	N6—C14—N8	128.2 (7)
C5—N4—C3	107.2 (7)	C14—C15—N7	106.5 (7)
C5—N4—H4	126.4	C14—C15—C12	122.6 (8)
C3—N4—H4	126.4	N7—C15—C12	130.2 (8)
C13—N5—C12	130.1 (8)	N7—C16—N8	112.6 (8)

supplementary materials

C13—N5—C17	115.8 (8)	N7—C16—H16	123.7
C12—N5—C17	114.1 (8)	N8—C16—H16	123.7
C14—N6—C13	119.3 (7)	N5—C17—H17A	109.5
C14—N6—C18	122.5 (8)	N5—C17—H17B	109.5
C13—N6—C18	118.1 (8)	H17A—C17—H17B	109.5
C16—N7—C15	106.8 (7)	N5—C17—H17C	109.5
C16—N7—C19	126.5 (7)	H17A—C17—H17C	109.5
C15—N7—C19	126.7 (7)	H17B—C17—H17C	109.5
C16—N8—C14	105.6 (7)	N6—C18—H18A	109.5
C16—N8—H8	127.2	N6—C18—H18B	109.5
C14—N8—H8	127.2	H18A—C18—H18B	109.5
O1—C1—N1	122.0 (9)	N6—C18—H18C	109.5
O1—C1—C4	128.4 (9)	H18A—C18—H18C	109.5
N1—C1—C4	109.5 (8)	H18B—C18—H18C	109.5
O2—C2—N2	120.6 (9)	N7—C19—C20	111.3 (8)
O2—C2—N1	123.2 (9)	N7—C19—H19A	109.4
N2—C2—N1	116.2 (8)	C20—C19—H19A	109.4
N2—C3—C4	123.5 (8)	N7—C19—H19B	109.4
N2—C3—N4	129.1 (7)	C20—C19—H19B	109.4
C4—C3—N4	107.4 (7)	H19A—C19—H19B	108.0
C3—C4—N3	106.5 (8)	O8—C20—O7	105.9 (8)
C3—C4—C1	122.3 (8)	O8—C20—C19	110.7 (9)
N3—C4—C1	130.6 (8)	O7—C20—C19	110.6 (8)
N3—C5—N4	111.3 (8)	O8—C20—H20	109.8
N3—C5—H5	124.3	O7—C20—H20	109.8
N4—C5—H5	124.3	C19—C20—H20	109.8
N1—C6—H6A	109.5	O8—C22—C21	103.2 (18)
N1—C6—H6B	109.5	O8—C22—H22A	111.1
H6A—C6—H6B	109.5	C21—C22—H22A	111.1
N1—C6—H6C	109.5	O8—C22—H22B	111.1
H6A—C6—H6C	109.5	C21—C22—H22B	111.1
H6B—C6—H6C	109.5	H22A—C22—H22B	109.1
N2—C7—H7A	109.5	O7—C21—C22	98.3 (18)
N2—C7—H7B	109.5	O7—C21—H21A	112.1
H7A—C7—H7B	109.5	C22—C21—H21A	112.1
N2—C7—H7C	109.5	O7—C21—H21B	112.1
H7A—C7—H7C	109.5	C22—C21—H21B	112.1
H7B—C7—H7C	109.5	H21A—C21—H21B	109.7
N3—C8—C9	110.9 (8)	O8—C22'—C21'	103.0 (13)
N3—C8—H8A	109.5	O8—C22'—H22C	111.2
C9—C8—H8A	109.5	C21'—C22'—H22C	111.2
N3—C8—H8B	109.5	O8—C22'—H22D	111.2
C9—C8—H8B	109.5	C21'—C22'—H22D	111.2
H8A—C8—H8B	108.0	H22C—C22'—H22D	109.1
O3—C9—O4	109.3 (10)	O7—C21'—C22'	96.4 (14)
O3—C9—C8	111.1 (8)	O7—C21'—H21C	112.5
O4—C9—C8	109.3 (9)	C22'—C21'—H21C	112.5
O3—C9—H9	109.0	O7—C21'—H21D	112.5
O4—C9—H9	109.0	C22'—C21'—H21D	112.5

C8—C9—H9

109.0

H21C—C21'—H21D

110.0

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N4—H4···O2 ⁱ	0.86	1.93	2.774 (9)	165
N8—H8···O6 ⁱⁱ	0.86	1.91	2.754 (9)	166
C5—H5···C15 ⁱ	0.93	2.80	3.650 (11)	153
C8—H8B···O7 ⁱⁱⁱ	0.97	2.29	3.114 (11)	142
C10—H10C···O5 ^{iv}	0.95	2.46	3.098 (11)	124
C10'—H10B···O5 ^{iv}	0.89	2.57	3.384 (11)	153
C10'—H10C···O5 ^{iv}	0.97	2.46	3.384 (11)	159
C16—H16···C11 ⁱⁱⁱ	0.93	2.78	3.560 (11)	143
C19—H19A···O4 ^v	0.97	2.48	3.227 (11)	134
C19—H19A···O5	0.97	2.53	3.167 (11)	123
C22'—H22D···O1	0.97	2.58	3.242 (11)	126

Symmetry codes: (i) $-y+1, x, z+1/2$; (ii) $y, -x, z+1/2$; (iii) $-x+1, -y, z$; (iv) $x, y, z+1$; (v) $-x+1, -y, z-1$.

Fig. 1

