

## 8-Hydroxy-2-methylquinolinium dibromido(2-methylquinolin-8-olato- $\kappa^2 N,O$ )zincate(II) methanol monosolvate

Ezzatollah Najafi,<sup>a</sup> Mostafa M. Amini<sup>a</sup> and Seik Weng Ng<sup>b\*</sup>

<sup>a</sup>Department of Chemistry, General Campus, Shahid Beheshti University, Tehran 1983963113, Iran, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 12 September 2010; accepted 14 September 2010

Key indicators: single-crystal X-ray study;  $T = 100\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.025;  $wR$  factor = 0.057; data-to-parameter ratio = 18.3.

The anion of the title salt,  $(\text{C}_{10}\text{H}_{10}\text{NO})[\text{ZnBr}_2(\text{C}_{10}\text{H}_8\text{NO})]\cdot\text{CH}_3\text{OH}$ , has its metal atom  $N,O$ -chelated by the deprotonated 2-methyl-8-hydroxyquinoline ligand. The hydroxy unit of the cation is a hydrogen-bond donor to the alkoxide O atom of the tetrahedrally coordinated anion, whereas the ammonium cation is a hydrogen-bond donor to the methanolic O atom. In the crystal, adjacent ion pairs and solvent molecules are linked by a methanol–halogen O–H···Br hydrogen bond, generating a chain running along the  $a$  axis.

### Related literature

For the isostructural chloro analog, see: Sattarzadeh *et al.* (2009).



### Experimental

#### Crystal data

$(\text{C}_{10}\text{H}_{10}\text{NO})[\text{ZnBr}_2(\text{C}_{10}\text{H}_8\text{NO})]\cdot\text{CH}_3\text{OH}$	$\beta = 105.815 (1)^\circ$
$M_r = 575.60$	$V = 2132.2 (3)\text{ \AA}^3$
Monoclinic, $P2_1/n$	$Z = 4$
$a = 9.9704 (8)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 13.9954 (11)\text{ \AA}$	$\mu = 4.93\text{ mm}^{-1}$
$c = 15.8815 (12)\text{ \AA}$	$T = 100\text{ K}$
	$0.35 \times 0.30 \times 0.25\text{ mm}$

#### Data collection

Bruker SMART APEX diffractometer	19908 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	4889 independent reflections
$T_{\min} = 0.278$ , $T_{\max} = 0.372$	4044 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.040$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$	267 parameters
$wR(F^2) = 0.057$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\max} = 0.49\text{ e \AA}^{-3}$
4889 reflections	$\Delta\rho_{\min} = -0.37\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2···O1	0.84	1.71	2.546 (2)	172
O3—H3···Br1 <sup>i</sup>	0.84	2.48	3.2941 (17)	163
N2—H2n···O3	0.86	1.91	2.739 (3)	162

Symmetry code: (i)  $x - 1, y, z$ .

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank Shahid Beheshti University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5354).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sattarzadeh, E., Mohammadnezhad, G., Amini, M. M. & Ng, S. W. (2009). *Acta Cryst. E65*, m553.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## **supplementary materials**

*Acta Cryst.* (2010). E66, m1276 [doi:10.1107/S1600536810036706]

## 8-Hydroxy-2-methylquinolinium dibromido(2-methylquinolin-8-olato- $\kappa^2N,O$ )zincate(II) methanol monosolvate

E. Najafi, M. M. Amini and S. W. Ng

### Comment

An earlier study reported  $C_{10}H_{10}NO^+\cdot ZnCl_2(C_{10}H_8NO)\cdot CH_3OH$ , which feature cations, tetrahedral anions and solvent molecules linked by  $N\cdots O$ ,  $O\cdots O$  and  $O\cdots Cl$  hydrogen bonds into a linear chain (Sattarzadeh *et al.*, 2009). The present bromide analog (Scheme I, Fig. 1) is isostructural, the two compounds displaying matching cell dimensions. The hydroxy unit of the cation is hydrogen-bond donor to the alkoxide O atom of the tetrahedrally coordinated anion whereas the ammonium unit is hydrogen-bond donor to the methanolic O atom. Adjacent ion-pairs and solvent molecules are linked by a  $O\text{--}H_{\text{methanol}}\cdots Br$  hydrogen bond to generate a linear chain running along the  $a$ -axis of the monoclinic unit cell.

### Experimental

Zinc bromide (0.10 g, 0.75 mmol) and 2-methyl-8-hydroxyquinoline (0.24 g, 1.5 mmol) were loaded into a convection tube; the tube was filled with dry methanol and kept at 333 K. Crystals were collected from the side arm after several days.

### Refinement

Hydrogen atoms were placed in calculated positions ( $C\text{--}H$  0.95–0.98,  $N\text{--}H$  0.86 and  $O\text{--}H$  0.84 Å) and were included in the refinement in the riding model approximation, with  $U(H)$  set to  $1.2\text{--}1.5U_{\text{eq}}(C,N,O)$ .

### Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of the title compound at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

## 8-Hydroxy-2-methylquinolinium dibromido(2-methylquinolin-8-olato- $\kappa^2N,O$ )zincate(II) methanol monosolvate

### Crystal data

$(C_{10}H_{10}NO)[ZnBr_2(C_{10}H_8NO)]\cdot CH_4O$

$F(000) = 1144$

$M_r = 575.60$

$D_x = 1.793 \text{ Mg m}^{-3}$

Monoclinic,  $P2_1/n$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Hall symbol: -P 2yn

Cell parameters from 6099 reflections

# supplementary materials

---

$a = 9.9704 (8)$ Å	$\theta = 2.6\text{--}28.2^\circ$
$b = 13.9954 (11)$ Å	$\mu = 4.93 \text{ mm}^{-1}$
$c = 15.8815 (12)$ Å	$T = 100$ K
$\beta = 105.815 (1)^\circ$	Prism, yellow
$V = 2132.2 (3)$ Å <sup>3</sup>	$0.35 \times 0.30 \times 0.25$ mm
$Z = 4$	

## Data collection

Bruker SMART APEX diffractometer	4889 independent reflections
Radiation source: fine-focus sealed tube graphite	4044 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.040$
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 27.5^\circ$ , $\theta_{\text{min}} = 2.0^\circ$
$T_{\text{min}} = 0.278$ , $T_{\text{max}} = 0.372$	$h = -12\text{--}12$
19908 measured reflections	$k = -18\text{--}18$
	$l = -19\text{--}20$

## Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.025$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.057$	H-atom parameters constrained
$S = 1.03$	$w = 1/[\sigma^2(F_o^2) + (0.0224P)^2 + 0.8543P]$ where $P = (F_o^2 + 2F_c^2)/3$
4889 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
267 parameters	$\Delta\rho_{\text{max}} = 0.49 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.37 \text{ e \AA}^{-3}$

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	1.00294 (2)	0.228905 (17)	0.184698 (15)	0.01708 (6)
Br2	1.12097 (3)	0.497149 (17)	0.227454 (16)	0.02018 (7)
Zn1	0.99449 (3)	0.369708 (19)	0.266658 (17)	0.01403 (7)
O1	0.79813 (16)	0.40171 (12)	0.25997 (10)	0.0166 (4)
O2	0.58392 (16)	0.36051 (12)	0.13590 (10)	0.0155 (3)
H2	0.6590	0.3718	0.1741	0.023*
O3	0.30839 (18)	0.27619 (14)	0.15184 (11)	0.0255 (4)
H3	0.2240	0.2660	0.1482	0.038*
N1	1.01652 (19)	0.36488 (13)	0.39831 (12)	0.0120 (4)
N2	0.35285 (19)	0.35026 (13)	0.00209 (12)	0.0129 (4)
H2N	0.3540	0.3348	0.0547	0.015*
C1	0.8930 (2)	0.38963 (15)	0.41453 (15)	0.0124 (5)

C2	0.7784 (2)	0.40932 (16)	0.33955 (15)	0.0141 (5)
C3	0.6537 (2)	0.43641 (17)	0.35416 (16)	0.0168 (5)
H3a	0.5762	0.4505	0.3058	0.020*
C4	0.6399 (3)	0.44345 (17)	0.43988 (16)	0.0181 (5)
H4	0.5525	0.4619	0.4480	0.022*
C5	0.7483 (3)	0.42455 (17)	0.51175 (16)	0.0185 (5)
H5	0.7362	0.4295	0.5689	0.022*
C6	0.8786 (2)	0.39750 (16)	0.50003 (15)	0.0143 (5)
C7	0.9995 (3)	0.37848 (16)	0.56932 (15)	0.0169 (5)
H7	0.9952	0.3826	0.6283	0.020*
C8	1.1213 (3)	0.35443 (16)	0.55191 (15)	0.0157 (5)
H8	1.2016	0.3415	0.5987	0.019*
C9	1.1288 (2)	0.34858 (15)	0.46464 (15)	0.0140 (5)
C10	1.2609 (2)	0.32358 (17)	0.44237 (16)	0.0177 (5)
H10A	1.2913	0.3781	0.4135	0.027*
H10B	1.2447	0.2683	0.4030	0.027*
H10C	1.3332	0.3079	0.4961	0.027*
C11	0.4758 (2)	0.37589 (15)	-0.01425 (15)	0.0124 (5)
C12	0.5986 (2)	0.38162 (15)	0.05651 (15)	0.0127 (5)
C13	0.7208 (2)	0.40884 (16)	0.03862 (15)	0.0144 (5)
H13	0.8044	0.4129	0.0847	0.017*
C14	0.7223 (2)	0.43073 (16)	-0.04781 (16)	0.0157 (5)
H14	0.8076	0.4494	-0.0588	0.019*
C15	0.6052 (2)	0.42585 (16)	-0.11591 (15)	0.0157 (5)
H15	0.6090	0.4413	-0.1735	0.019*
C16	0.4782 (2)	0.39768 (15)	-0.10042 (15)	0.0133 (5)
C17	0.3494 (2)	0.39320 (16)	-0.16628 (16)	0.0164 (5)
H17	0.3470	0.4067	-0.2253	0.020*
C18	0.2294 (3)	0.36989 (16)	-0.14609 (16)	0.0178 (5)
H18	0.1438	0.3684	-0.1908	0.021*
C19	0.2312 (2)	0.34802 (16)	-0.05948 (15)	0.0152 (5)
C20	0.1026 (2)	0.32331 (18)	-0.03312 (17)	0.0212 (5)
H20A	0.1008	0.3598	0.0192	0.032*
H20B	0.1025	0.2548	-0.0204	0.032*
H20C	0.0202	0.3392	-0.0809	0.032*
C21	0.3867 (3)	0.27543 (19)	0.24135 (16)	0.0218 (5)
H21A	0.4864	0.2742	0.2452	0.033*
H21B	0.3623	0.2186	0.2701	0.033*
H21C	0.3652	0.3329	0.2704	0.033*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.01686 (12)	0.02067 (12)	0.01393 (12)	-0.00406 (9)	0.00458 (9)	-0.00180 (9)
Br2	0.02141 (13)	0.02021 (12)	0.02170 (14)	-0.00056 (10)	0.01060 (11)	0.00474 (10)
Zn1	0.01209 (14)	0.02082 (14)	0.00922 (14)	-0.00032 (11)	0.00298 (10)	0.00126 (11)
O1	0.0102 (8)	0.0304 (9)	0.0080 (8)	0.0014 (7)	0.0003 (6)	0.0007 (7)
O2	0.0124 (8)	0.0244 (9)	0.0078 (8)	-0.0009 (7)	-0.0005 (6)	0.0012 (7)

## supplementary materials

---

O3	0.0159 (9)	0.0454 (12)	0.0159 (9)	-0.0032 (9)	0.0056 (7)	0.0036 (8)
N1	0.0126 (9)	0.0126 (9)	0.0103 (9)	-0.0021 (8)	0.0024 (8)	0.0006 (7)
N2	0.0143 (10)	0.0131 (9)	0.0103 (10)	0.0007 (8)	0.0019 (8)	0.0002 (7)
C1	0.0154 (12)	0.0108 (10)	0.0119 (11)	-0.0042 (9)	0.0051 (9)	-0.0005 (8)
C2	0.0157 (12)	0.0144 (11)	0.0119 (12)	-0.0035 (9)	0.0032 (9)	0.0000 (9)
C3	0.0130 (11)	0.0203 (12)	0.0154 (12)	-0.0033 (9)	0.0011 (10)	0.0006 (9)
C4	0.0150 (12)	0.0197 (12)	0.0217 (13)	-0.0031 (10)	0.0084 (10)	-0.0042 (10)
C5	0.0270 (14)	0.0165 (12)	0.0152 (12)	-0.0053 (10)	0.0113 (11)	-0.0031 (10)
C6	0.0189 (12)	0.0124 (11)	0.0116 (11)	-0.0042 (9)	0.0043 (10)	0.0006 (9)
C7	0.0262 (13)	0.0152 (12)	0.0090 (12)	-0.0049 (10)	0.0045 (10)	-0.0006 (9)
C8	0.0197 (12)	0.0131 (11)	0.0120 (12)	-0.0016 (9)	0.0003 (10)	0.0007 (9)
C9	0.0177 (12)	0.0097 (11)	0.0144 (12)	-0.0032 (9)	0.0039 (10)	0.0010 (9)
C10	0.0158 (12)	0.0202 (12)	0.0153 (13)	0.0008 (10)	0.0013 (10)	0.0005 (10)
C11	0.0160 (12)	0.0095 (10)	0.0118 (11)	0.0033 (9)	0.0041 (9)	-0.0007 (8)
C12	0.0153 (11)	0.0120 (11)	0.0100 (11)	0.0022 (9)	0.0018 (9)	-0.0016 (8)
C13	0.0114 (11)	0.0165 (11)	0.0140 (12)	0.0022 (9)	0.0012 (9)	-0.0018 (9)
C14	0.0142 (12)	0.0155 (11)	0.0199 (13)	0.0023 (9)	0.0090 (10)	-0.0004 (9)
C15	0.0212 (13)	0.0161 (11)	0.0112 (12)	0.0023 (10)	0.0065 (10)	-0.0001 (9)
C16	0.0168 (12)	0.0108 (10)	0.0115 (11)	0.0049 (9)	0.0021 (9)	-0.0008 (9)
C17	0.0217 (13)	0.0139 (11)	0.0134 (12)	0.0052 (9)	0.0042 (10)	-0.0012 (9)
C18	0.0175 (12)	0.0186 (12)	0.0144 (12)	0.0021 (10)	-0.0009 (10)	-0.0022 (9)
C19	0.0152 (12)	0.0114 (11)	0.0175 (13)	0.0007 (9)	0.0017 (10)	-0.0005 (9)
C20	0.0145 (12)	0.0254 (13)	0.0220 (14)	-0.0006 (10)	0.0023 (11)	0.0035 (11)
C21	0.0207 (13)	0.0275 (14)	0.0160 (13)	-0.0011 (11)	0.0032 (10)	0.0022 (10)

*Geometric parameters (Å, °)*

Br1—Zn1	2.3758 (4)	C8—H8	0.9500
Br2—Zn1	2.3637 (4)	C9—C10	1.496 (3)
Zn1—O1	1.9828 (16)	C10—H10A	0.9800
Zn1—N1	2.0431 (19)	C10—H10B	0.9800
O1—C2	1.335 (3)	C10—H10C	0.9800
O2—C12	1.341 (3)	C11—C16	1.409 (3)
O2—H2	0.8400	C11—C12	1.421 (3)
O3—C21	1.423 (3)	C12—C13	1.378 (3)
O3—H3	0.8400	C13—C14	1.411 (3)
N1—C9	1.331 (3)	C13—H13	0.9500
N1—C1	1.370 (3)	C14—C15	1.359 (3)
N2—C19	1.335 (3)	C14—H14	0.9500
N2—C11	1.368 (3)	C15—C16	1.411 (3)
N2—H2N	0.8600	C15—H15	0.9500
C1—C6	1.408 (3)	C16—C17	1.419 (3)
C1—C2	1.434 (3)	C17—C18	1.360 (3)
C2—C3	1.379 (3)	C17—H17	0.9500
C3—C4	1.409 (3)	C18—C19	1.404 (3)
C3—H3a	0.9500	C18—H18	0.9500
C4—C5	1.367 (3)	C19—C20	1.494 (3)
C4—H4	0.9500	C20—H20A	0.9800
C5—C6	1.414 (3)	C20—H20B	0.9800

C5—H5	0.9500	C20—H20C	0.9800
C6—C7	1.418 (3)	C21—H21A	0.9800
C7—C8	1.359 (3)	C21—H21B	0.9800
C7—H7	0.9500	C21—H21C	0.9800
C8—C9	1.411 (3)		
O1—Zn1—N1	83.77 (7)	H10A—C10—H10B	109.5
O1—Zn1—Br2	113.91 (5)	C9—C10—H10C	109.5
N1—Zn1—Br2	112.22 (5)	H10A—C10—H10C	109.5
O1—Zn1—Br1	109.87 (5)	H10B—C10—H10C	109.5
N1—Zn1—Br1	121.55 (5)	N2—C11—C16	119.6 (2)
Br2—Zn1—Br1	112.362 (14)	N2—C11—C12	119.2 (2)
C2—O1—Zn1	111.41 (13)	C16—C11—C12	121.1 (2)
C12—O2—H2	109.5	O2—C12—C13	125.7 (2)
C21—O3—H3	109.5	O2—C12—C11	116.1 (2)
C9—N1—C1	119.98 (19)	C13—C12—C11	118.2 (2)
C9—N1—Zn1	130.41 (16)	C12—C13—C14	120.4 (2)
C1—N1—Zn1	109.43 (14)	C12—C13—H13	119.8
C19—N2—C11	123.3 (2)	C14—C13—H13	119.8
C19—N2—H2N	118.3	C15—C14—C13	121.9 (2)
C11—N2—H2N	118.3	C15—C14—H14	119.0
N1—C1—C6	122.3 (2)	C13—C14—H14	119.0
N1—C1—C2	116.5 (2)	C14—C15—C16	119.5 (2)
C6—C1—C2	121.2 (2)	C14—C15—H15	120.3
O1—C2—C3	123.6 (2)	C16—C15—H15	120.3
O1—C2—C1	118.8 (2)	C11—C16—C15	118.9 (2)
C3—C2—C1	117.6 (2)	C11—C16—C17	117.1 (2)
C2—C3—C4	120.8 (2)	C15—C16—C17	124.0 (2)
C2—C3—H3a	119.6	C18—C17—C16	120.9 (2)
C4—C3—H3a	119.6	C18—C17—H17	119.5
C5—C4—C3	122.0 (2)	C16—C17—H17	119.5
C5—C4—H4	119.0	C17—C18—C19	120.4 (2)
C3—C4—H4	119.0	C17—C18—H18	119.8
C4—C5—C6	119.2 (2)	C19—C18—H18	119.8
C4—C5—H5	120.4	N2—C19—C18	118.6 (2)
C6—C5—H5	120.4	N2—C19—C20	118.6 (2)
C1—C6—C5	119.1 (2)	C18—C19—C20	122.8 (2)
C1—C6—C7	116.5 (2)	C19—C20—H20A	109.5
C5—C6—C7	124.4 (2)	C19—C20—H20B	109.5
C8—C7—C6	120.4 (2)	H20A—C20—H20B	109.5
C8—C7—H7	119.8	C19—C20—H20C	109.5
C6—C7—H7	119.8	H20A—C20—H20C	109.5
C7—C8—C9	120.3 (2)	H20B—C20—H20C	109.5
C7—C8—H8	119.9	O3—C21—H21A	109.5
C9—C8—H8	119.9	O3—C21—H21B	109.5
N1—C9—C8	120.6 (2)	H21A—C21—H21B	109.5
N1—C9—C10	117.2 (2)	O3—C21—H21C	109.5
C8—C9—C10	122.1 (2)	H21A—C21—H21C	109.5
C9—C10—H10A	109.5	H21B—C21—H21C	109.5
C9—C10—H10B	109.5		

## supplementary materials

---

N1—Zn1—O1—C2	3.23 (14)	C6—C7—C8—C9	0.3 (3)
Br2—Zn1—O1—C2	−108.35 (14)	C1—N1—C9—C8	1.4 (3)
Br1—Zn1—O1—C2	124.58 (13)	Zn1—N1—C9—C8	176.10 (15)
O1—Zn1—N1—C9	−178.0 (2)	C1—N1—C9—C10	−178.93 (19)
Br2—Zn1—N1—C9	−64.7 (2)	Zn1—N1—C9—C10	−4.3 (3)
Br1—Zn1—N1—C9	72.5 (2)	C7—C8—C9—N1	−1.1 (3)
O1—Zn1—N1—C1	−2.90 (14)	C7—C8—C9—C10	179.2 (2)
Br2—Zn1—N1—C1	110.42 (13)	C19—N2—C11—C16	−2.7 (3)
Br1—Zn1—N1—C1	−112.43 (13)	C19—N2—C11—C12	176.3 (2)
C9—N1—C1—C6	−0.9 (3)	N2—C11—C12—O2	0.2 (3)
Zn1—N1—C1—C6	−176.60 (17)	C16—C11—C12—O2	179.15 (19)
C9—N1—C1—C2	177.82 (19)	N2—C11—C12—C13	−179.13 (19)
Zn1—N1—C1—C2	2.1 (2)	C16—C11—C12—C13	−0.2 (3)
Zn1—O1—C2—C3	176.09 (18)	O2—C12—C13—C14	−179.0 (2)
Zn1—O1—C2—C1	−3.1 (2)	C11—C12—C13—C14	0.3 (3)
N1—C1—C2—O1	0.6 (3)	C12—C13—C14—C15	−0.1 (3)
C6—C1—C2—O1	179.3 (2)	C13—C14—C15—C16	−0.3 (3)
N1—C1—C2—C3	−178.6 (2)	N2—C11—C16—C15	178.7 (2)
C6—C1—C2—C3	0.1 (3)	C12—C11—C16—C15	−0.2 (3)
O1—C2—C3—C4	−179.7 (2)	N2—C11—C16—C17	1.1 (3)
C1—C2—C3—C4	−0.6 (3)	C12—C11—C16—C17	−177.8 (2)
C2—C3—C4—C5	0.4 (4)	C14—C15—C16—C11	0.4 (3)
C3—C4—C5—C6	0.3 (4)	C14—C15—C16—C17	177.9 (2)
N1—C1—C6—C5	179.2 (2)	C11—C16—C17—C18	0.8 (3)
C2—C1—C6—C5	0.6 (3)	C15—C16—C17—C18	−176.8 (2)
N1—C1—C6—C7	0.1 (3)	C16—C17—C18—C19	−1.2 (3)
C2—C1—C6—C7	−178.6 (2)	C11—N2—C19—C18	2.2 (3)
C4—C5—C6—C1	−0.7 (3)	C11—N2—C19—C20	−177.2 (2)
C4—C5—C6—C7	178.4 (2)	C17—C18—C19—N2	−0.3 (3)
C1—C6—C7—C8	0.2 (3)	C17—C18—C19—C20	179.1 (2)
C5—C6—C7—C8	−178.9 (2)		

### Hydrogen-bond geometry ( $\text{\AA}$ , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O2—H2 $\cdots$ O1	0.84	1.71	2.546 (2)	172
O3—H3 $\cdots$ Br1 <sup>i</sup>	0.84	2.48	3.2941 (17)	163
N2—H2n $\cdots$ O3	0.86	1.91	2.739 (3)	162

Symmetry codes: (i)  $x-1, y, z$ .

Fig. 1

