

catena-Poly[[aquabis(pyridine- κN)-copper(II)]- μ -2,2'-(*p*-phenylenedioxy)-diacetato- $\kappa^2 O:O'$]

Xiu-Mei Zhang^a and Ya-Feng Li^{b*}

^aDepartment of X-ray, First Hospital, Jilin University, Changchun 130021, People's Republic of China, and ^bSchool of Chemical Engineering, Changchun University of Technology, Changchun 130012, People's Republic of China
Correspondence e-mail: fly012345@sohu.com


Received 6 September 2010; accepted 12 September 2010

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.009$ Å;
 R factor = 0.071; wR factor = 0.123; data-to-parameter ratio = 12.4.

In the title compound, $[Cu(C_{10}H_8O_6)(C_5H_5N)_2(H_2O)]_n$, the Cu atom is five-coordinated by two O atoms from two carboxylate groups of two different 2,2'-(*p*-phenylenedioxy)diacetate ligands, two N atoms from two pyridine molecules and one water O atom. The geometry is square-pyramidal with the water O atom occupying the apical position. The carboxylate group bridges adjacent Cu atoms, forming an infinite zigzag chain extending parallel to [001]. The chains are linked into layers by O—H···O hydrogen bonds. The Cu and water O atoms lie on special positions of site symmetry 2.

Related literature

For the isotopic zinc analog, see: Hong *et al.* (2005).



Experimental

Crystal data

$[Cu(C_{10}H_8O_6)(C_5H_5N)_2(H_2O)]$

$M_r = 463.93$

Monoclinic, $C2/c$	$Z = 4$
$a = 15.363$ (4) Å	Mo $K\alpha$ radiation
$b = 6.0888$ (12) Å	$\mu = 1.14$ mm $^{-1}$
$c = 21.896$ (6) Å	$T = 298$ K
$\beta = 103.67$ (3) $^\circ$	$0.12 \times 0.11 \times 0.09$ mm
$V = 1990.2$ (8) Å 3	

Data collection

Rigaku R-AXIS RAPID	7227 measured reflections
diffractometer	1737 independent reflections
Absorption correction: multi-scan	1096 reflections with $I > 2\sigma(I)$
(ABSCOR; Higashi, 1995)	$R_{\text{int}} = 0.119$
$T_{\min} = 0.875$, $T_{\max} = 0.907$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.071$	H atoms treated by a mixture of
$wR(F^2) = 0.123$	independent and constrained
$S = 1.08$	refinement
1737 reflections	$\Delta\rho_{\max} = 0.32$ e Å $^{-3}$
140 parameters	$\Delta\rho_{\min} = -0.38$ e Å $^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1W—H1A···O2 ⁱ	0.81 (6)	1.87 (6)	2.677 (5)	174 (7)
Symmetry code: (i) $-x, y + 1, -z + \frac{1}{2}$.				

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2000); software used to prepare material for publication: *SHELXL97*.

This project was sponsored by the Scientific Research Foundation for Returned Overseas Chinese Scholars, Chinese Education Ministry (20071108) and the Scientific Research Foundation for the Returned Overseas Team, Chinese Education Ministry.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5029).

References

- Brandenburg, K. (2000). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Hong, X.-L., Li, Y.-Z. & Bai, J.-F. (2005). *Acta Cryst. E* **61**, m1863–m1865.
Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
Rigaku/MSC (2002). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, m1283 [doi:10.1107/S1600536810036469]

catena-Poly[[aquabis(pyridine- κN)copper(II)]- μ -2,2'-(*p*-phenylenedioxy)diacetato- $\kappa^2 O:O'$]

X.-M. Zhang and Y.-F. Li

Comment

The metal-organic framework, which is formed by carboxylate ligand as the strut and transition metal as the node, has recently attracted more attentions owing to the potential applications of catalyst and H-storage. In this paper, flexible ligand-BDOA as the strut bridges the Cu cations to result in the formation of infinite zigzag chain of $[Cu(py)_2(H_2O)BDOA]_n$.

The asymmetrical unit of $[Cu(py)_2(H_2O)BDOA]_n$ (py=pyridine, BDOA=benzene-1,4-dioxyacetate) which is isostructural with of $[Zn(py)_2(H_2O)BDOA]_n$ (Hong, *et al.*, 2005), is composed of one half of Cu cation, one half of BDOA, one half of water molecule and one pyridine molecule (Fig.1). Five-coordinated Cu cation lies in the basal position of pyramid constructed by two O atoms from two carboxyl groups of two different BDOA, two nitrogen atoms of two pyridines and one water oxygen atom situated at the apical position. The bond distances of Cu—N, Cu—O and Cu—Ow are 2.109 (23) Å, 1.965 (35) Å and 2.192 (6) Å, respectively. The bond angles of O—Cu—O and N—Cu—O are 179.85 (14) $^\circ$ and 167.68 (18) $^\circ$, respectively. Cu and water oxygen lie at 2-fold axis and BDOA at the inversion center.

The monodentate μ_2 -BDOA bridges the adjacent Cu cations to form the infinite zigzag chain along (001) direction. The H-bonds of Ow—H \cdots O (free oxygen of carboxyl) link the adjacent chains to two-dimensional layer (*bc* planar), which is packed by the van der Waals force (Fig.2).

Experimental

(I) was synthesized under hydrothermal condition. In a typically route, H_2BDOA (0.22 g, 1 mmol) was dissolved in 10 ml deionized water under stirring, and then pyridine (1.6 ml, 20 mmol) and $Cu(Ac)_2 \cdot 3H_2O$ (0.235 g, 1 mmol) were added to a blue solution. After continuously stirred for 1 h, the solution with the molar ratio of H_2BDOA : 20py: $Cu(Ac)_2 \cdot 3H_2O$: 555 H_2O was transferred into 23 ml autoclave and heated at 438 K for 5 days. After naturally cooling to room temperature, blue block product was collected by filtration as a single phase.

Refinement

Water H atoms were located in a difference Fourier map and were refined with O—H = 0.82 (2) Å, H \cdots H = 1.37 (2) Å and $U_{iso}(H) = 1.2U_{eq}(O)$. The remaining H-atoms were placed in calculated positions (C—H (phenyl and pyridine ring) = 0.93 Å, C—H (methylene) = 0.97 Å) and were included in the refinement in the riding-model approximation, with U(H) = 1.2U_{eq}(C).

supplementary materials

Figures




Fig. 1. The unit cell of (I), showing the atomic labelling scheme and displacement ellipsoids at the 50% probability level. [Symmetry codes: (i) $-x, y, 0.5 - z$; (ii) $-x, 1 - y, -z$.]




Fig. 2. The ball-stick plot of (I), displaying the zigzag chain along (001) direction composed of bridging the Cu cation with monodentate μ_2 -BDOA. Cu is shown in the cyan, O in red, N in blue and C in grey.

catena-Poly[[aquabis(pyridine-κN)copper(II)]-μ-2,2'-(*p*-phenylenedioxy)diacetato-κ²O:O']

Crystal data

[Cu(C ₁₀ H ₈ O ₆)(C ₅ H ₅ N) ₂ (H ₂ O)]	$F(000) = 956$
$M_r = 463.93$	$D_x = 1.548 \text{ Mg m}^{-3}$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -C 2yc	Cell parameters from 2000 reflections
$a = 15.363 (4) \text{ \AA}$	$\theta = 3.6\text{--}25^\circ$
$b = 6.0888 (12) \text{ \AA}$	$\mu = 1.14 \text{ mm}^{-1}$
$c = 21.896 (6) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 103.67 (3)^\circ$	Block, blue
$V = 1990.2 (8) \text{ \AA}^3$	$0.12 \times 0.11 \times 0.09 \text{ mm}$
$Z = 4$	

Data collection

Rigaku R-AXIS RAPID diffractometer	1737 independent reflections
Radiation source: fine-focus sealed tube graphite	1096 reflections with $I > 2\sigma(I)$
Detector resolution: 10.00 pixels mm ⁻¹	$R_{\text{int}} = 0.119$
ω scans	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 3.6^\circ$
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	$h = -18 \rightarrow 18$
$T_{\text{min}} = 0.875, T_{\text{max}} = 0.907$	$k = -7 \rightarrow 6$
7227 measured reflections	$l = -25 \rightarrow 25$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.071$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.123$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.0254P)^2 + 4.0919P]$ where $P = (F_o^2 + 2F_c^2)/3$
1737 reflections	$(\Delta/\sigma)_{\max} < 0.001$
140 parameters	$\Delta\rho_{\max} = 0.32 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.38 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.0000	1.02668 (15)	0.2500	0.0480 (4)
O1	-0.0864 (2)	1.0271 (7)	0.16798 (16)	0.0584 (10)
O2	-0.0691 (3)	0.6662 (7)	0.15674 (17)	0.0679 (12)
O3	-0.1479 (3)	0.6794 (7)	0.03254 (17)	0.0656 (12)
C1	-0.0948 (4)	0.8473 (11)	0.1365 (2)	0.0506 (15)
C2	-0.1406 (4)	0.8781 (10)	0.0675 (2)	0.0581 (16)
H2A	-0.2001	0.9377	0.0643	0.070*
H2B	-0.1070	0.9840	0.0493	0.070*
C3	-0.0715 (4)	0.5975 (10)	0.0191 (2)	0.0528 (15)
C4	0.0125 (4)	0.6894 (10)	0.0344 (3)	0.0618 (17)
H4	0.0218	0.8182	0.0579	0.074*
C5	-0.0824 (4)	0.4048 (10)	-0.0158 (3)	0.0605 (17)
H5	-0.1385	0.3387	-0.0266	0.073*
N1	0.0988 (3)	0.9911 (8)	0.20420 (18)	0.0503 (12)
C6	0.1565 (4)	0.8241 (10)	0.2156 (3)	0.0571 (16)
H6	0.1512	0.7228	0.2463	0.068*
C7	0.2226 (4)	0.7939 (11)	0.1847 (3)	0.0681 (18)
H7	0.2605	0.6731	0.1937	0.082*
C8	0.2327 (4)	0.9417 (13)	0.1405 (3)	0.0713 (19)
H8	0.2787	0.9270	0.1198	0.086*
C9	0.1732 (5)	1.1140 (12)	0.1271 (3)	0.075 (2)
H9	0.1771	1.2156	0.0962	0.090*
C10	0.1086 (4)	1.1333 (10)	0.1598 (3)	0.0659 (18)

supplementary materials

H10	0.0693	1.2514	0.1508	0.079*
O1W	0.0000	1.3867 (10)	0.2500	0.085 (2)
H1A	0.023 (5)	1.463 (10)	0.280 (3)	0.102*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0449 (6)	0.0454 (6)	0.0508 (6)	0.000	0.0055 (4)	0.000
O1	0.047 (2)	0.067 (3)	0.055 (2)	0.003 (2)	0.0017 (18)	-0.007 (2)
O2	0.082 (3)	0.057 (3)	0.059 (3)	0.000 (2)	0.006 (2)	0.007 (2)
O3	0.052 (3)	0.080 (3)	0.062 (2)	-0.005 (2)	0.009 (2)	-0.016 (2)
C1	0.040 (4)	0.069 (5)	0.042 (3)	-0.005 (3)	0.008 (3)	0.001 (3)
C2	0.055 (4)	0.064 (4)	0.052 (3)	0.004 (3)	0.005 (3)	-0.001 (3)
C3	0.043 (4)	0.070 (4)	0.042 (3)	-0.001 (3)	0.004 (3)	-0.002 (3)
C4	0.057 (5)	0.065 (4)	0.060 (4)	-0.010 (3)	0.008 (3)	-0.015 (3)
C5	0.042 (4)	0.072 (4)	0.067 (4)	-0.012 (3)	0.013 (3)	-0.006 (3)
N1	0.042 (3)	0.057 (3)	0.048 (3)	-0.005 (3)	0.004 (2)	0.001 (2)
C6	0.051 (4)	0.062 (4)	0.057 (4)	0.005 (3)	0.010 (3)	0.011 (3)
C7	0.054 (5)	0.073 (5)	0.081 (5)	0.005 (4)	0.024 (4)	0.006 (4)
C8	0.049 (4)	0.103 (6)	0.066 (4)	-0.026 (4)	0.020 (3)	-0.020 (4)
C9	0.063 (5)	0.093 (5)	0.070 (4)	-0.018 (4)	0.018 (4)	0.017 (4)
C10	0.053 (4)	0.069 (4)	0.070 (4)	-0.004 (3)	0.004 (3)	0.015 (4)
O1W	0.114 (6)	0.041 (4)	0.078 (5)	0.000	-0.021 (4)	0.000

Geometric parameters (\AA , $^\circ$)

Cu1—O1	1.964 (3)	C4—H4	0.9300
Cu1—O1 ⁱ	1.964 (3)	C5—C4 ⁱⁱ	1.362 (8)
Cu1—N1	2.019 (4)	C5—H5	0.9300
Cu1—N1 ⁱ	2.019 (4)	N1—C6	1.333 (7)
Cu1—O1W	2.192 (6)	N1—C10	1.338 (7)
O1—C1	1.284 (6)	C6—C7	1.359 (8)
O2—C1	1.219 (6)	C6—H6	0.9300
O3—C3	1.370 (6)	C7—C8	1.357 (8)
O3—C2	1.422 (6)	C7—H7	0.9300
C1—C2	1.520 (7)	C8—C9	1.377 (9)
C2—H2A	0.9700	C8—H8	0.9300
C2—H2B	0.9700	C9—C10	1.357 (8)
C3—C4	1.373 (7)	C9—H9	0.9300
C3—C5	1.388 (8)	C10—H10	0.9300
C4—C5 ⁱⁱ	1.362 (8)	O1W—H1A	0.81 (6)
O1—Cu1—O1 ⁱ	179.8 (3)	C5 ⁱⁱ —C4—H4	119.4
O1—Cu1—N1	88.32 (15)	C3—C4—H4	119.4
O1 ⁱ —Cu1—N1	91.70 (15)	C4 ⁱⁱ —C5—C3	121.3 (6)
O1—Cu1—N1 ⁱ	91.70 (15)	C4 ⁱⁱ —C5—H5	119.4
O1 ⁱ —Cu1—N1 ⁱ	88.32 (15)	C3—C5—H5	119.4
N1—Cu1—N1 ⁱ	167.7 (3)	C6—N1—C10	116.5 (5)

O1—Cu1—O1W	89.92 (13)	C6—N1—Cu1	122.2 (4)
O1 ⁱ —Cu1—O1W	89.92 (13)	C10—N1—Cu1	121.3 (4)
N1—Cu1—O1W	96.16 (14)	N1—C6—C7	123.5 (6)
N1 ⁱ —Cu1—O1W	96.16 (14)	N1—C6—H6	118.3
C1—O1—Cu1	116.8 (4)	C7—C6—H6	118.3
C3—O3—C2	117.5 (5)	C8—C7—C6	119.4 (6)
O2—C1—O1	126.4 (5)	C8—C7—H7	120.3
O2—C1—C2	120.4 (5)	C6—C7—H7	120.3
O1—C1—C2	113.1 (5)	C7—C8—C9	118.3 (6)
O3—C2—C1	113.0 (5)	C7—C8—H8	120.8
O3—C2—H2A	109.0	C9—C8—H8	120.8
C1—C2—H2A	109.0	C10—C9—C8	119.0 (6)
O3—C2—H2B	109.0	C10—C9—H9	120.5
C1—C2—H2B	109.0	C8—C9—H9	120.5
H2A—C2—H2B	107.8	N1—C10—C9	123.3 (6)
O3—C3—C4	127.1 (6)	N1—C10—H10	118.4
O3—C3—C5	115.3 (5)	C9—C10—H10	118.4
C4—C3—C5	117.6 (6)	Cu1—O1W—H1A	125 (5)
C5 ⁱⁱ —C4—C3	121.1 (6)		
N1—Cu1—O1—C1	66.8 (4)	O1 ⁱ —Cu1—N1—C6	56.9 (4)
N1 ⁱ —Cu1—O1—C1	-100.9 (4)	N1 ⁱ —Cu1—N1—C6	-33.0 (4)
O1W—Cu1—O1—C1	162.9 (4)	O1W—Cu1—N1—C6	147.0 (4)
Cu1—O1—C1—O2	15.7 (8)	O1—Cu1—N1—C10	55.6 (4)
Cu1—O1—C1—C2	-163.2 (3)	O1 ⁱ —Cu1—N1—C10	-124.3 (4)
C3—O3—C2—C1	-73.6 (6)	N1 ⁱ —Cu1—N1—C10	145.9 (4)
O2—C1—C2—O3	-0.3 (8)	O1W—Cu1—N1—C10	-34.1 (4)
O1—C1—C2—O3	178.6 (5)	C10—N1—C6—C7	-0.1 (8)
C2—O3—C3—C4	-1.8 (8)	Cu1—N1—C6—C7	178.8 (4)
C2—O3—C3—C5	-179.3 (4)	N1—C6—C7—C8	1.3 (10)
O3—C3—C4—C5 ⁱⁱ	-177.2 (5)	C6—C7—C8—C9	-2.2 (9)
C5—C3—C4—C5 ⁱⁱ	0.3 (9)	C7—C8—C9—C10	2.0 (9)
O3—C3—C5—C4 ⁱⁱ	177.5 (5)	C6—N1—C10—C9	-0.1 (8)
C4—C3—C5—C4 ⁱⁱ	-0.3 (9)	Cu1—N1—C10—C9	-179.0 (5)
O1—Cu1—N1—C6	-123.2 (4)	C8—C9—C10—N1	-0.9 (10)

Symmetry codes: (i) $-x, y, -z+1/2$; (ii) $-x, -y+1, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1W—H1A \cdots O2 ⁱⁱⁱ	0.81 (6)	1.87 (6)	2.677 (5)	174 (7)

Symmetry codes: (iii) $-x, y+1, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

