

Bis[2-(4-benzoyloxy-2-hydroxybenzoyl)-1-phenylethenolato]diethanolzinc(II)

Kai Dong,^a Juan Sun,^a Ban-Feng Ruan^{a*} and Hai-Bin Gong^{a,b*}

^aState Key Laboratory of Pharmaceutical Biotechnology, Nanjing University, Nanjing 210093, People's Republic of China, and ^bXuzhou Central Hospital, Xuzhou Cardiovascular Disease Institute, Xuzhou 221009, Jiangsu, People's Republic of China

Correspondence e-mail: bf_ruan@163.com, adler_20008@yahoo.com.cn


Received 2 September 2010; accepted 8 September 2010

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.050; wR factor = 0.142; data-to-parameter ratio = 15.3.

The mononuclear title complex, $[\text{Zn}(\text{C}_{22}\text{H}_{15}\text{O}_5)_2(\text{C}_2\text{H}_5\text{OH})_2]$, contains a Zn^{II} atom (site symmetry $\bar{1}$) surrounded by six O atoms of the keto groups of two substituted 1,3-diketonate ligands and of two ethanol molecules, resulting in a distorted octahedral coordination environment. The molecular configuration is stabilized by an intramolecular hydrogen bond between the phenolic hydroxy group and the adjacent keto group. The hydroxy group acts likewise as an acceptor of an intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond with the hydroxy group of the ethanol molecule as the donor. The hydrogen-bonding scheme leads to the formation of supramolecular layers parallel to (010).

Related literature

For the role of zinc in enzymes and in bioinorganic chemistry, see: Bertini *et al.* (1994); Lipscomb & Strater (1996); Vallee & Auld (1993); Zhu *et al.* (2003).



Experimental

Crystal data

$[\text{Zn}(\text{C}_{22}\text{H}_{15}\text{O}_5)_2(\text{C}_2\text{H}_5\text{O})_2]$	$\gamma = 90.989 (5)^\circ$
$M_r = 876.19$	$V = 1056.6 (10) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 1$
$a = 7.170 (5) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.399 (5) \text{ \AA}$	$\mu = 0.65 \text{ mm}^{-1}$
$c = 16.457 (5) \text{ \AA}$	$T = 298 \text{ K}$
$\alpha = 106.590 (5)^\circ$	$0.30 \times 0.20 \times 0.20 \text{ mm}$
$\beta = 95.596 (5)^\circ$	

Data collection

Bruker SMART CCD area-detector diffractometer	6185 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2000)	4255 independent reflections
$T_{\text{min}} = 0.830$, $T_{\text{max}} = 0.882$	2898 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$	14 restraints
$wR(F^2) = 0.142$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.80 \text{ e \AA}^{-3}$
4255 reflections	$\Delta\rho_{\text{min}} = -0.49 \text{ e \AA}^{-3}$
279 parameters	

Table 1

Selected bond lengths (Å).

Zn1—O5 ⁱ	2.002 (2)	Zn1—O6 ⁱ	2.203 (3)
Zn1—O4 ⁱ	2.034 (2)		

Symmetry code: (i) $-x, -y + 1, -z + 1$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O3}-\text{H3}\cdots\text{O4}$	0.82	1.74	2.470 (3)	147
$\text{O6}-\text{H10A}\cdots\text{O3}^{\text{ii}}$	0.86	2.04	2.824 (4)	151

Symmetry code: (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The work was financed by a grant (project 30772627) from the National Natural Science Foundation of China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2402).

References

- Bertini, I., Gray, H. B., Lippard, S. J. & Valentine, J. S. (1994). *Bioinorganic Chemistry*. Mills Valley, CA, USA: University Science Books.
- Bruker (2000). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Lipscomb, W. N. & Strater, N. (1996). *Chem. Rev.* **96**, 2375–2434.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Vallee, B. L. & Auld, D. S. (1993). *Acc. Chem. Res.* **26**, 543–551.
- Zhu, H.-L., Meng, F.-J., Wang, Z.-D. & Yang, F. (2003). *Z. Kristallogr. New Cryst. Struct.* **219**, 321–322.

supplementary materials

Acta Cryst. (2010). E66, m1290 [doi:10.1107/S1600536810036202]

Bis[2-(4-benzoyloxy-2-hydroxybenzoyl)-1-phenylethenolato]diethanolzinc(II)

K. Dong, J. Sun, B.-F. Ruan and H.-B. Gong

Comment

Zinc, the second most abundant transition metal in biology, functions as the active site of hydrolytic enzymes, such as carboxypeptidase and carbonic anhydrase, where it is in a hard donor coordination environment of nitrogen and oxygen (Lipscomb & Strater, 1996; Bertini *et al.*, 1994). Zinc has long been recognized as an important cofactor in biological molecules, either as a structural template in protein folding or as a Lewis acid catalyst that can readily adopt 4-, 5- or 6-coordination (Vallee & Auld, 1993). Studies of the structures and properties of the metal coordination sites in zinc enzymes recently were in focus in the field of bioinorganic chemistry (Zhu *et al.*, 2003). A couple of chemical systems have been developed to reproduce the zinc coordination core. On our continuation of the research in this field, the crystal structure of a mononuclear zinc(II) compound, (I), is reported here.

In the structure of (I), the Zn^{II} atom is situated on an inversion center and adopts a slightly distorted octahedral coordination (Fig. 1). It is surrounded by six O atoms from the keto groups of two 1,3-diketonate ligands and by two ethanol molecules. The Zn^{II} atom and the four O atoms from the diketone ligands form the equatorial plane with similar Zn—O bond lengths of 2.002 (2) and 2.034 (2) Å. The two ethanol O atoms are in the axial positions with considerably longer Zn—O bond lengths of 2.203 (3) Å (Table 1). The two trans-angles at the zinc(II) center are 180° by symmetry; all angles around Zn1 are close to 90°, ranging from 85.93 (11) to 94.07 (11)°. In the crystal structure, intramolecular and intermolecular O—H...O hydrogen bonding leads to a layered assembly parallel to (010) (Table 2; Fig. 2).

Experimental

To a ethanol solution (15 ml) of 3-hydroxy-4-(3-oxo-3-phenylpropanoyl)phenyl benzoic acid (0.36 g, 0.001 mol), anhydrous zinc acetate (0.092 g, 0.0005 mol) was added at room temperature. After addition, the reaction mixture was refluxed for 30 min with stirring and then was filtered. Single crystals of the title compound were obtained by slow evaporation of the filtrate.

Refinement

All hydrogen atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H = 0.93, 0.96(—CH₃), 0.97(—CH₂), 0.86(—OH) Å, $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C or —OH}), 1.5 U_{\text{eq}}(\text{—CH}_3)$, respectively.

Figures




Fig. 1. The structure and atom-numbering scheme of the title compound (I), showing 30% probability displacement ellipsoids. [Symmetry code: (A) $-x, -y+1, -z+1$.]




Fig. 2. The crystal packing of the title compound viewed along the *b* axis. Hydrogen bonding interactions are indicated by dashed lines.

Bis[2-(4-benzoyloxy-2-hydroxybenzoyl)-1-phenylethenolato]diethanolzinc(II)

Crystal data

[Zn(C₂₂H₁₅O₅)₂(C₂H₆O)₂]

M_r = 876.19

Triclinic, *P* $\bar{1}$

Hall symbol: -P 1

a = 7.170 (5) Å

b = 9.399 (5) Å

c = 16.457 (5) Å

α = 106.590 (5)°

β = 95.596 (5)°

γ = 90.989 (5)°

V = 1056.6 (10) Å³

Z = 1

F(000) = 456

D_x = 1.377 Mg m⁻³

Mo *K*α radiation, λ = 0.71069 Å

Cell parameters from 2800 reflections

θ = 2.5–26.2°

μ = 0.65 mm⁻¹

T = 298 K

Block, colourless

0.30 × 0.20 × 0.20 mm

Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2000)

T_{min} = 0.830, *T_{max}* = 0.882

6185 measured reflections

4255 independent reflections

2898 reflections with *I* > 2σ(*I*)

R_{int} = 0.017

θ_{\max} = 26.5°, θ_{\min} = 2.3°

h = -9→8

k = -11→11

l = -18→20

Refinement

Refinement on *F*²

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.142$

S = 1.02

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0597P)^2 + 0.7172P]$

where $P = (F_o^2 + 2F_c^2)/3$

4255 reflections	$(\Delta/\sigma)_{\max} < 0.001$
279 parameters	$\Delta\rho_{\max} = 0.80 \text{ e } \text{\AA}^{-3}$
14 restraints	$\Delta\rho_{\min} = -0.49 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.0307 (7)	0.0321 (6)	-0.1364 (3)	0.0787 (14)
H1	1.1025	-0.0287	-0.1751	0.094*
C2	1.0779 (7)	0.1805 (6)	-0.1042 (3)	0.0828 (14)
H2	1.1814	0.2205	-0.1214	0.099*
C3	0.9730 (6)	0.2707 (5)	-0.0466 (3)	0.0670 (11)
H3A	1.0040	0.3720	-0.0260	0.080*
C4	0.8221 (5)	0.2120 (4)	-0.0193 (2)	0.0465 (8)
C5	0.7732 (5)	0.0614 (4)	-0.0524 (2)	0.0572 (10)
H5	0.6708	0.0209	-0.0347	0.069*
C6	0.8769 (7)	-0.0277 (5)	-0.1115 (3)	0.0723 (12)
H6	0.8431	-0.1283	-0.1345	0.087*
C7	0.7210 (5)	0.3140 (5)	0.0470 (2)	0.0522 (9)
C8	0.4913 (5)	0.3187 (4)	0.1426 (2)	0.0533 (9)
C9	0.3464 (5)	0.4014 (5)	0.1234 (2)	0.0625 (11)
H9	0.3287	0.4179	0.0701	0.075*
C10	0.2282 (5)	0.4591 (4)	0.1841 (2)	0.0560 (10)
H10	0.1290	0.5137	0.1707	0.067*
C11	0.2513 (4)	0.4390 (4)	0.2655 (2)	0.0417 (8)
C12	0.4069 (5)	0.3582 (4)	0.2832 (2)	0.0462 (8)
C13	0.5241 (5)	0.2980 (4)	0.2214 (2)	0.0539 (9)
H13	0.6247	0.2436	0.2336	0.065*
C14	0.1192 (4)	0.4955 (4)	0.3303 (2)	0.0427 (8)
C15	-0.0393 (5)	0.5691 (4)	0.3129 (2)	0.0475 (8)
H15	-0.0592	0.5804	0.2584	0.057*
C16	-0.1725 (4)	0.6280 (4)	0.3689 (2)	0.0409 (7)
C17	-0.3308 (4)	0.7128 (4)	0.3413 (2)	0.0425 (8)
C18	-0.3685 (5)	0.7198 (5)	0.2587 (2)	0.0648 (11)
H18	-0.2969	0.6672	0.2169	0.078*
C19	-0.5106 (6)	0.8032 (6)	0.2374 (3)	0.0817 (14)

supplementary materials

H19	-0.5329	0.8071	0.1815	0.098*
C20	-0.6184 (6)	0.8797 (6)	0.2966 (3)	0.0816 (14)
H20	-0.7108	0.9394	0.2824	0.098*
C21	-0.5890 (7)	0.8675 (6)	0.3773 (3)	0.0971 (18)
H21	-0.6660	0.9158	0.4179	0.117*
C22	-0.4472 (6)	0.7849 (5)	0.3996 (3)	0.0749 (13)
H22	-0.4299	0.7778	0.4551	0.090*
C23	0.1763 (9)	0.8386 (7)	0.5795 (5)	0.132 (2)
H23A	0.1946	0.8477	0.5236	0.159*
H23B	0.2850	0.8879	0.6176	0.159*
C24	0.0214 (9)	0.9216 (7)	0.6062 (5)	0.129 (2)
H24A	-0.0890	0.8780	0.5689	0.193*
H24B	0.0426	1.0218	0.6046	0.193*
H24C	0.0044	0.9217	0.6634	0.193*
O1	0.7452 (5)	0.4469 (3)	0.0716 (2)	0.0838 (10)
O2	0.5993 (4)	0.2389 (3)	0.07885 (17)	0.0653 (7)
O3	0.4461 (4)	0.3341 (4)	0.36033 (16)	0.0670 (8)
H3	0.3672	0.3722	0.3917	0.100*
O4	0.1634 (3)	0.4709 (3)	0.40277 (15)	0.0600 (7)
O5	-0.1712 (3)	0.6190 (3)	0.44437 (14)	0.0547 (7)
O6	0.1818 (4)	0.6942 (3)	0.5742 (2)	0.0773 (9)
H10A	0.2735	0.6729	0.6057	0.093*
Zn1	0.0000	0.5000	0.5000	0.0548 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.084 (3)	0.097 (4)	0.057 (3)	0.036 (3)	0.033 (2)	0.014 (3)
C2	0.083 (3)	0.093 (4)	0.089 (3)	0.017 (3)	0.055 (3)	0.035 (3)
C3	0.068 (3)	0.065 (3)	0.077 (3)	0.008 (2)	0.033 (2)	0.026 (2)
C4	0.0436 (19)	0.058 (2)	0.0402 (18)	0.0108 (17)	0.0117 (15)	0.0155 (17)
C5	0.051 (2)	0.063 (3)	0.057 (2)	0.0030 (19)	0.0140 (18)	0.0138 (19)
C6	0.078 (3)	0.066 (3)	0.063 (3)	0.013 (2)	0.014 (2)	0.002 (2)
C7	0.045 (2)	0.061 (3)	0.053 (2)	0.0063 (18)	0.0159 (16)	0.0166 (19)
C8	0.049 (2)	0.057 (2)	0.055 (2)	0.0120 (18)	0.0262 (17)	0.0105 (18)
C9	0.061 (2)	0.087 (3)	0.049 (2)	0.019 (2)	0.0232 (18)	0.027 (2)
C10	0.049 (2)	0.076 (3)	0.052 (2)	0.0238 (19)	0.0175 (16)	0.0276 (19)
C11	0.0332 (17)	0.050 (2)	0.0443 (18)	0.0100 (14)	0.0108 (13)	0.0142 (15)
C12	0.0352 (17)	0.059 (2)	0.0461 (19)	0.0112 (16)	0.0106 (14)	0.0148 (17)
C13	0.0413 (19)	0.062 (2)	0.060 (2)	0.0182 (17)	0.0158 (16)	0.0157 (19)
C14	0.0331 (17)	0.056 (2)	0.0419 (18)	0.0105 (15)	0.0102 (13)	0.0158 (16)
C15	0.0419 (19)	0.066 (2)	0.0399 (18)	0.0205 (17)	0.0128 (14)	0.0207 (17)
C16	0.0343 (17)	0.051 (2)	0.0398 (18)	0.0085 (15)	0.0068 (13)	0.0150 (15)
C17	0.0362 (17)	0.050 (2)	0.0434 (18)	0.0107 (15)	0.0061 (14)	0.0152 (15)
C18	0.058 (2)	0.096 (3)	0.053 (2)	0.033 (2)	0.0175 (18)	0.037 (2)
C19	0.075 (3)	0.121 (4)	0.066 (3)	0.046 (3)	0.010 (2)	0.050 (3)
C20	0.068 (3)	0.105 (4)	0.082 (3)	0.049 (3)	0.008 (2)	0.041 (3)
C21	0.088 (3)	0.139 (5)	0.073 (3)	0.078 (3)	0.027 (3)	0.034 (3)

C22	0.072 (3)	0.111 (4)	0.051 (2)	0.054 (3)	0.019 (2)	0.030 (2)
C23	0.112 (4)	0.086 (2)	0.197 (5)	0.020 (3)	-0.007 (4)	0.044 (3)
C24	0.114 (4)	0.110 (4)	0.147 (4)	0.040 (3)	0.014 (3)	0.009 (3)
O1	0.092 (2)	0.0581 (19)	0.098 (2)	0.0027 (17)	0.0523 (19)	0.0045 (17)
O2	0.0656 (17)	0.0624 (17)	0.0721 (18)	0.0123 (14)	0.0423 (14)	0.0140 (14)
O3	0.0497 (15)	0.107 (2)	0.0525 (15)	0.0400 (15)	0.0158 (12)	0.0316 (16)
O4	0.0456 (14)	0.100 (2)	0.0461 (14)	0.0360 (14)	0.0165 (11)	0.0350 (14)
O5	0.0496 (14)	0.0804 (18)	0.0429 (13)	0.0331 (13)	0.0165 (11)	0.0264 (13)
O6	0.0566 (17)	0.0793 (18)	0.092 (2)	0.0247 (15)	-0.0049 (15)	0.0215 (17)
Zn1	0.0475 (4)	0.0847 (5)	0.0425 (3)	0.0331 (3)	0.0159 (3)	0.0296 (3)

Geometric parameters (Å, °)

C1—C2	1.366 (7)	C15—H15	0.9300
C1—C6	1.378 (6)	C16—O5	1.269 (4)
C1—H1	0.9300	C16—C17	1.506 (4)
C2—C3	1.373 (6)	C17—C22	1.372 (5)
C2—H2	0.9300	C17—C18	1.379 (5)
C3—C4	1.375 (5)	C18—C19	1.375 (5)
C3—H3A	0.9300	C18—H18	0.9300
C4—C5	1.389 (5)	C19—C20	1.352 (6)
C4—C7	1.489 (5)	C19—H19	0.9300
C5—C6	1.377 (5)	C20—C21	1.363 (6)
C5—H5	0.9300	C20—H20	0.9300
C6—H6	0.9300	C21—C22	1.374 (5)
C7—O1	1.202 (4)	C21—H21	0.9300
C7—O2	1.346 (4)	C22—H22	0.9300
C8—C13	1.365 (5)	C23—O6	1.337 (6)
C8—C9	1.374 (5)	C23—C24	1.402 (6)
C8—O2	1.411 (4)	C23—H23A	0.9700
C9—C10	1.372 (5)	C23—H23B	0.9700
C9—H9	0.9300	C24—H24A	0.9600
C10—C11	1.400 (5)	C24—H24B	0.9600
C10—H10	0.9300	C24—H24C	0.9600
C11—C12	1.414 (4)	O3—H3	0.8200
C11—C14	1.486 (4)	O4—Zn1	2.034 (2)
C12—O3	1.356 (4)	O5—Zn1	2.002 (2)
C12—C13	1.383 (5)	O6—Zn1	2.203 (3)
C13—H13	0.9300	O6—H10A	0.8600
C14—O4	1.289 (4)	Zn1—O5 ⁱ	2.002 (2)
C14—C15	1.389 (4)	Zn1—O4 ⁱ	2.034 (2)
C15—C16	1.402 (4)	Zn1—O6 ⁱ	2.203 (3)
C2—C1—C6	120.2 (4)	C18—C17—C16	123.2 (3)
C2—C1—H1	119.9	C19—C18—C17	120.9 (4)
C6—C1—H1	119.9	C19—C18—H18	119.5
C1—C2—C3	120.2 (4)	C17—C18—H18	119.5
C1—C2—H2	119.9	C20—C19—C18	121.0 (4)
C3—C2—H2	119.9	C20—C19—H19	119.5

supplementary materials

C2—C3—C4	120.4 (4)	C18—C19—H19	119.5
C2—C3—H3A	119.8	C19—C20—C21	118.7 (4)
C4—C3—H3A	119.8	C19—C20—H20	120.7
C3—C4—C5	119.4 (3)	C21—C20—H20	120.7
C3—C4—C7	117.8 (3)	C20—C21—C22	121.0 (4)
C5—C4—C7	122.8 (3)	C20—C21—H21	119.5
C6—C5—C4	119.8 (4)	C22—C21—H21	119.5
C6—C5—H5	120.1	C17—C22—C21	120.9 (4)
C4—C5—H5	120.1	C17—C22—H22	119.6
C5—C6—C1	119.9 (4)	C21—C22—H22	119.6
C5—C6—H6	120.0	O6—C23—C24	121.8 (6)
C1—C6—H6	120.0	O6—C23—H23A	106.9
O1—C7—O2	122.9 (3)	C24—C23—H23A	106.9
O1—C7—C4	125.4 (3)	O6—C23—H23B	106.9
O2—C7—C4	111.7 (3)	C24—C23—H23B	106.9
C13—C8—C9	121.4 (3)	H23A—C23—H23B	106.7
C13—C8—O2	117.3 (3)	C23—C24—H24A	109.5
C9—C8—O2	121.0 (3)	C23—C24—H24B	109.5
C10—C9—C8	118.9 (3)	H24A—C24—H24B	109.5
C10—C9—H9	120.6	C23—C24—H24C	109.5
C8—C9—H9	120.6	H24A—C24—H24C	109.5
C9—C10—C11	122.5 (3)	H24B—C24—H24C	109.5
C9—C10—H10	118.7	C7—O2—C8	119.2 (3)
C11—C10—H10	118.7	C12—O3—H3	109.5
C10—C11—C12	116.4 (3)	C14—O4—Zn1	126.7 (2)
C10—C11—C14	122.9 (3)	C16—O5—Zn1	126.6 (2)
C12—C11—C14	120.7 (3)	C23—O6—Zn1	132.2 (4)
O3—C12—C13	117.2 (3)	C23—O6—H10A	114.0
O3—C12—C11	121.8 (3)	Zn1—O6—H10A	113.8
C13—C12—C11	121.0 (3)	O5—Zn1—O5 ⁱ	180.00 (11)
C8—C13—C12	119.7 (3)	O5—Zn1—O4	89.37 (9)
C8—C13—H13	120.1	O5 ⁱ —Zn1—O4	90.63 (9)
C12—C13—H13	120.1	O5—Zn1—O4 ⁱ	90.63 (9)
O4—C14—C15	123.3 (3)	O5 ⁱ —Zn1—O4 ⁱ	89.37 (9)
O4—C14—C11	115.0 (3)	O4—Zn1—O4 ⁱ	179.999 (1)
C15—C14—C11	121.7 (3)	O5—Zn1—O6	94.07 (11)
C14—C15—C16	126.8 (3)	O5 ⁱ —Zn1—O6	85.93 (11)
C14—C15—H15	116.6	O4—Zn1—O6	89.52 (12)
C16—C15—H15	116.6	O4 ⁱ —Zn1—O6	90.48 (12)
O5—C16—C15	125.3 (3)	O5—Zn1—O6 ⁱ	85.93 (11)
O5—C16—C17	114.8 (3)	O5 ⁱ —Zn1—O6 ⁱ	94.07 (11)
C15—C16—C17	119.9 (3)	O4—Zn1—O6 ⁱ	90.48 (12)
C22—C17—C18	117.4 (3)	O4 ⁱ —Zn1—O6 ⁱ	89.52 (12)
C22—C17—C16	119.4 (3)	O6—Zn1—O6 ⁱ	180.0

Symmetry codes: (i) $-x, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O3—H3···O4	0.82	1.74	2.470 (3)	147
O6—H10A···O3 ⁱⁱ	0.86	2.04	2.824 (4)	151

Symmetry codes: (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

