

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(2E)-2-Benzylidene-5,6-dimethoxyindan-1-oneMohamed Ashraf Ali,^a Rusli Ismail,^a Soo Choon Tan,^a Chin Sing Yeap^{b‡} and Hoong-Kun Fun^{b*§}^aInstitute for Research in Molecular Medicine, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia

Correspondence e-mail: hkfun@usm.my


Received 3 September 2010; accepted 6 September 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.043; wR factor = 0.130; data-to-parameter ratio = 31.3.

The molecular structure of the title compound, $\text{C}_{18}\text{H}_{16}\text{O}_3$, is roughly planar; the maximum deviation of the indanone ring system is 0.027 (1) Å and it makes a dihedral angle of 2.69 (3)° with the phenyl ring. The torsion angles between the two methoxy groups and the indanone ring are -14.67 (11) and -1.11 (12)°. In the crystal, molecules are connected into a ribbon along the a axis via weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds. Weak intermolecular $\text{C}-\text{H}\cdots\pi$ and $\pi-\pi$ [centroid-centroid distance = 3.7086 (6) Å] interactions are also observed.

Related literature

For general background to and the biological activity of chalcone derivatives, see: Boumendjel *et al.* (2009); D'Archivio *et al.* (2008); Dicarolo *et al.* (1999); Echeverria *et al.* (2009); Heidenreich *et al.* (2008); Katsori & Hadjipavlou-Latina (2009); Miranda *et al.* (1999); Nowakowska (2007); Shah *et al.* (2008); Syed *et al.* (2008). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



‡ Thomson Reuters ResearcherID: A-5523-2009.

§ Thomson Reuters ResearcherID: A-3561-2009.

Experimental

Crystal data

$\text{C}_{18}\text{H}_{16}\text{O}_3$
 $M_r = 280.31$
 Monoclinic, $P2_1/c$
 $a = 6.0209$ (6) Å
 $b = 14.8550$ (14) Å
 $c = 15.2292$ (15) Å
 $\beta = 90.603$ (2)°
 $V = 1362.0$ (2) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 100$ K
 $0.44 \times 0.29 \times 0.16$ mm

Data collection

Bruker APEXII DUO CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.960$, $T_{\max} = 0.985$
 31475 measured reflections
 6003 independent reflections
 4902 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.130$
 $S = 1.09$
 6003 reflections
 192 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.60$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.58$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of C2–C7 benzene ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C17}-\text{H17A}\cdots\text{O3}^{\text{i}}$	0.96	2.53	3.4107 (11)	152
$\text{C18}-\text{H18B}\cdots\text{O2}^{\text{ii}}$	0.96	2.58	3.5320 (11)	173
$\text{C16}-\text{H16A}\cdots\text{Cg1}^{\text{iii}}$	0.93	2.99	3.7224 (9)	137

Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $x + 1, y, z$; (iii) $x, -y + \frac{1}{2}, z - \frac{3}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

The authors wish to express their thanks to Universiti Sains Malaysia (USM) for providing research facilities. HKF thanks USM for the Research University Grant No. 1001/PFIZIK/811160 and CSY thanks USM for the award of a USM Fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2598).

References

- Boumendjel, A., Ronot, X. & Boutonnat, J. (2009). *Curr. Drug Targets*, **10**, 363–371.
 Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
 D'Archivio, M., Santangelo, C., Scazzocchio, B., Vari, R., Filesi, C., Masella, R. & Giovannini, C. (2008). *Int. J. Mol. Sci.* **9**, 213–228.
 Dicarolo, G., Mascolo, N., Izzo, A. A. & Capasso, F. (1999). *Life Sci.* **65**, 337–353.
 Echeverria, C., Santibanez, J. S., Donoso-Tauda, O., Escobar, C. A. & Ramirez-Tagle, R. (2009). *Int. J. Mol. Sci.* **10**, 221–231.
 Heidenreich, A., Aus, G., Bolla, M., Joniau, S., Matveev, V. B., Schmid, H. P. & Zattoni, F. (2008). *Eur. Urol.* **53**, 68–80.

- Katsori, A. M. & Hadjipavlou-Latina, D. (2009). *Curr. Med. Chem.* **16**, 1062–1081.
- Miranda, C. L., Stevens, J. F., Helmrich, A., Henderson, M. C., Rodriguez, R. J., Yang, Y. H., Deinzer, M. L., Barnes, D. W. & Buhler, D. R. (1999). *Food Chem. Toxicol.* **37**, 271–285.
- Nowakowska, Z. (2007). *Eur. J. Med. Chem.* **42**, 125–137.
- Shah, A., Khan, A. M., Qureshi, R., Ansari, F. L., Nazar, M. F. & Shah, S. S. (2008). *Int. J. Mol. Sci.* **9**, 1424–1434.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Syed, D. N., Suh, Y., Afag, F. & Mukhtar, H. (2008). *Cancer Lett.* **265**, 167–176.

supplementary materials

Acta Cryst. (2010). E66, o2531-o2532 [doi:10.1107/S1600536810035695]

(2*E*)-2-Benzylidene-5,6-dimethoxyindan-1-one

M. A. Ali, R. Ismail, S. C. Tan, C. S. Yeap and H.-K. Fun

Comment

Chalcones have been reported to possess antiinflammatory, antimicrobial, antioxidant and anticancer properties (Echeverria *et al.*, 2009; Nowakowska, 2007; Miranda *et al.*, 1999; Shah *et al.*, 2008; Boumendjel *et al.*, 2009; Katsori & Hadjipavlou-Latina, 2009). Chalcones are one of the major classes of natural products with widespread distribution in spices, tea, beer, fruits and vegetables. They have been recently subjects of great interest for their pharmacological activities (Dicarlo *et al.*, 1999). Prostate cancer is one of the most commonly diagnosed cancers in men and the second leading cause of cancer deaths in the European Union and United States of America (Heidenreich *et al.*, 2008). Many antitumor drugs have been developed for prostate cancer patients, but their intolerable systemic toxicity often limits their clinical use. Chemoprevention is one of the most promising approaches in prostate cancer research, in which natural or synthetic agents are used to prevent this malignant disease (Heidenreich *et al.*, 2008; Syed *et al.*, 2008; D'Archivio *et al.*, 2008).

The molecular structure of the title compound is essentially coplanar (Fig. 1). The maximum deviation of the indanone group is 0.027 (1) Å and it makes dihedral angle of 2.69 (3)° with the phenyl ring [C11–C16]. The torsion angles of the two methoxy groups are [C17–O2–C4–C5] 165.99 (7) and [C18–O3–C5–C4] 179.19 (7)°.

In the crystal structure, intermolecular C17—H17A···O3 hydrogen bonds (Table 1) link the molecules into dimers (Fig. 2). These dimers are interconnected into ribbons propagating along the [100] direction *via* intermolecular C18—H18B···O2 hydrogen bonds (Fig. 2, Table 1). Weak intermolecular C—H··· π (Table 1) and π – π interactions are also observed. [Cg1···Cg2^{iv} of 3.7086 (6) Å; (iv) 1 - x, 1 - y, -z. Cg1 and Cg2 are the centroids of C2–C7 and C11–C16 benzene ring.]

Experimental

A mixture of 5,6-dimethoxyindan-1-one (0.001 mmol) and benzaldehyde (0.001 mmol) were dissolved in methanol (10 ml) and 30% sodium hydroxide solution (5 ml) was added and stirred for 5 h. After completion of the reaction as evident from TLC, the mixture was poured into crushed ice then neutralized with concentrated HCl. The precipitated solid was filtered, washed with water and recrystallized from ethanol to reveal the title compound as light yellow crystals.

Refinement

All H atoms were positioned geometrically (C—H = 0.93–0.97 Å) and refined using a riding model. A rotating-group model were applied for the methyl groups.

Figures




Fig. 1. The molecular structure of the title compound with atom labels and 50% probability ellipsoids for non-H atoms.




Fig. 2. The crystal packing of title compound, showing chains along the [100] direction. Intermolecular hydrogen bonds are shown as dashed lines.

(2E)-2-Benzylidene-5,6-dimethoxyindan-1-one

Crystal data

$C_{18}H_{16}O_3$

$M_r = 280.31$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 6.0209$ (6) Å

$b = 14.8550$ (14) Å

$c = 15.2292$ (15) Å

$\beta = 90.603$ (2)°

$V = 1362.0$ (2) Å³

$Z = 4$

$F(000) = 592$

$D_x = 1.367$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7296 reflections

$\theta = 2.7\text{--}34.9^\circ$

$\mu = 0.09$ mm⁻¹

$T = 100$ K

Yellow, colourless

$0.44 \times 0.29 \times 0.16$ mm

Data collection

Bruker APEXII DUO CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2009)

$T_{\min} = 0.960$, $T_{\max} = 0.985$

31475 measured reflections

6003 independent reflections

4902 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.048$

$\theta_{\max} = 35.0^\circ$, $\theta_{\min} = 1.9^\circ$

$h = -9 \rightarrow 9$

$k = -23 \rightarrow 23$

$l = -23 \rightarrow 24$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.043$

$wR(F^2) = 0.130$

$S = 1.09$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0698P)^2 + 0.2363P]$

6003 reflections
192 parameters
0 restraints

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.60 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.58 \text{ e } \text{\AA}^{-3}$$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.07043 (10)	0.71482 (4)	0.09037 (4)	0.01685 (13)
O2	0.07512 (10)	0.55593 (4)	0.41523 (4)	0.01527 (12)
O3	0.42675 (10)	0.46101 (4)	0.40716 (4)	0.01623 (13)
C1	0.22764 (13)	0.66575 (5)	0.11035 (5)	0.01189 (13)
C2	0.26214 (13)	0.61731 (5)	0.19344 (5)	0.01122 (13)
C3	0.12486 (13)	0.61613 (5)	0.26768 (5)	0.01208 (13)
H3A	-0.0055	0.6497	0.2689	0.014*
C4	0.18842 (13)	0.56395 (5)	0.33862 (5)	0.01188 (13)
C5	0.38768 (13)	0.51154 (5)	0.33496 (5)	0.01213 (13)
C6	0.52425 (13)	0.51522 (5)	0.26166 (5)	0.01263 (14)
H6A	0.6561	0.4827	0.2602	0.015*
C7	0.45911 (12)	0.56873 (5)	0.19043 (5)	0.01121 (13)
C8	0.57760 (13)	0.58179 (5)	0.10412 (5)	0.01231 (13)
H8A	0.5966	0.5249	0.0738	0.015*
H8B	0.7219	0.6095	0.1131	0.015*
C9	0.42250 (13)	0.64368 (5)	0.05349 (5)	0.01186 (13)
C10	0.43407 (13)	0.67809 (5)	-0.02828 (5)	0.01274 (14)
H10A	0.3154	0.7150	-0.0441	0.015*
C11	0.60191 (13)	0.66724 (5)	-0.09638 (5)	0.01229 (13)
C12	0.79718 (14)	0.61621 (6)	-0.08618 (5)	0.01469 (14)
H12A	0.8259	0.5868	-0.0334	0.018*
C13	0.94801 (14)	0.60945 (6)	-0.15463 (6)	0.01690 (15)
H13A	1.0760	0.5751	-0.1472	0.020*
C14	0.90917 (15)	0.65353 (6)	-0.23413 (6)	0.01682 (15)
H14A	1.0106	0.6487	-0.2795	0.020*
C15	0.71743 (15)	0.70489 (6)	-0.24506 (6)	0.01576 (15)

supplementary materials

H15A	0.6910	0.7349	-0.2977	0.019*
C16	0.56526 (14)	0.71135 (5)	-0.17716 (5)	0.01416 (14)
H16A	0.4370	0.7454	-0.1853	0.017*
C17	-0.09525 (14)	0.62140 (6)	0.43137 (6)	0.01619 (15)
H17A	-0.1545	0.6122	0.4890	0.024*
H17B	-0.2118	0.6149	0.3883	0.024*
H17C	-0.0333	0.6808	0.4276	0.024*
C18	0.62128 (14)	0.40534 (6)	0.40883 (6)	0.01640 (15)
H18A	0.6235	0.3701	0.4617	0.025*
H18B	0.7514	0.4426	0.4071	0.025*
H18C	0.6192	0.3660	0.3588	0.025*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0151 (3)	0.0202 (3)	0.0153 (3)	0.0057 (2)	0.0018 (2)	0.0026 (2)
O2	0.0144 (3)	0.0191 (3)	0.0123 (3)	0.0035 (2)	0.0052 (2)	0.0029 (2)
O3	0.0157 (3)	0.0197 (3)	0.0133 (3)	0.0048 (2)	0.0024 (2)	0.0056 (2)
C1	0.0121 (3)	0.0125 (3)	0.0111 (3)	0.0000 (2)	0.0014 (2)	-0.0003 (2)
C2	0.0117 (3)	0.0116 (3)	0.0104 (3)	-0.0001 (2)	0.0013 (2)	0.0003 (2)
C3	0.0116 (3)	0.0128 (3)	0.0119 (3)	0.0004 (2)	0.0017 (2)	0.0001 (2)
C4	0.0109 (3)	0.0137 (3)	0.0111 (3)	-0.0003 (2)	0.0024 (2)	0.0000 (2)
C5	0.0122 (3)	0.0126 (3)	0.0116 (3)	0.0001 (2)	0.0005 (2)	0.0014 (2)
C6	0.0116 (3)	0.0143 (3)	0.0120 (3)	0.0013 (2)	0.0012 (2)	0.0009 (2)
C7	0.0112 (3)	0.0114 (3)	0.0110 (3)	-0.0005 (2)	0.0013 (2)	-0.0001 (2)
C8	0.0118 (3)	0.0140 (3)	0.0111 (3)	0.0013 (2)	0.0021 (2)	0.0007 (2)
C9	0.0118 (3)	0.0124 (3)	0.0115 (3)	0.0005 (2)	0.0019 (2)	0.0000 (2)
C10	0.0134 (3)	0.0130 (3)	0.0118 (3)	0.0010 (2)	0.0015 (2)	0.0003 (2)
C11	0.0136 (3)	0.0120 (3)	0.0113 (3)	-0.0004 (2)	0.0015 (2)	0.0002 (2)
C12	0.0143 (3)	0.0165 (3)	0.0133 (3)	0.0018 (3)	0.0019 (3)	0.0014 (3)
C13	0.0148 (3)	0.0193 (4)	0.0167 (4)	0.0021 (3)	0.0038 (3)	-0.0005 (3)
C14	0.0175 (4)	0.0185 (4)	0.0146 (3)	-0.0029 (3)	0.0057 (3)	-0.0015 (3)
C15	0.0186 (4)	0.0164 (3)	0.0123 (3)	-0.0030 (3)	0.0024 (3)	0.0015 (3)
C16	0.0154 (3)	0.0145 (3)	0.0126 (3)	0.0003 (2)	0.0012 (3)	0.0013 (2)
C17	0.0141 (3)	0.0189 (4)	0.0156 (3)	0.0024 (3)	0.0043 (3)	-0.0007 (3)
C18	0.0153 (3)	0.0169 (3)	0.0171 (4)	0.0032 (3)	-0.0010 (3)	0.0032 (3)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.2303 (10)	C10—C11	1.4645 (11)
O2—C4	1.3628 (10)	C10—H10A	0.9300
O2—C17	1.4366 (10)	C11—C12	1.4061 (11)
O3—C5	1.3499 (10)	C11—C16	1.4092 (11)
O3—C18	1.4338 (10)	C12—C13	1.3933 (12)
C1—C2	1.4687 (11)	C12—H12A	0.9300
C1—C9	1.5017 (11)	C13—C14	1.3941 (12)
C2—C7	1.3894 (11)	C13—H13A	0.9300
C2—C3	1.4076 (11)	C14—C15	1.3923 (13)
C3—C4	1.3805 (11)	C14—H14A	0.9300

C3—H3A	0.9300	C15—C16	1.3920 (12)
C4—C5	1.4318 (11)	C15—H15A	0.9300
C5—C6	1.3945 (11)	C16—H16A	0.9300
C6—C7	1.3977 (11)	C17—H17A	0.9600
C6—H6A	0.9300	C17—H17B	0.9600
C7—C8	1.5146 (11)	C17—H17C	0.9600
C8—C9	1.5154 (11)	C18—H18A	0.9600
C8—H8A	0.9700	C18—H18B	0.9600
C8—H8B	0.9700	C18—H18C	0.9600
C9—C10	1.3487 (11)		
C4—O2—C17	116.88 (6)	C9—C10—H10A	114.5
C5—O3—C18	118.07 (7)	C11—C10—H10A	114.5
O1—C1—C2	127.21 (7)	C12—C11—C16	118.04 (7)
O1—C1—C9	126.20 (7)	C12—C11—C10	124.30 (7)
C2—C1—C9	106.59 (6)	C16—C11—C10	117.65 (7)
C7—C2—C3	121.92 (7)	C13—C12—C11	120.47 (8)
C7—C2—C1	109.82 (7)	C13—C12—H12A	119.8
C3—C2—C1	128.26 (7)	C11—C12—H12A	119.8
C4—C3—C2	118.38 (7)	C12—C13—C14	120.74 (8)
C4—C3—H3A	120.8	C12—C13—H13A	119.6
C2—C3—H3A	120.8	C14—C13—H13A	119.6
O2—C4—C3	125.58 (7)	C15—C14—C13	119.50 (8)
O2—C4—C5	114.40 (7)	C15—C14—H14A	120.2
C3—C4—C5	120.02 (7)	C13—C14—H14A	120.2
O3—C5—C6	125.08 (7)	C16—C15—C14	120.02 (8)
O3—C5—C4	114.17 (7)	C16—C15—H15A	120.0
C6—C5—C4	120.75 (7)	C14—C15—H15A	120.0
C5—C6—C7	118.72 (7)	C15—C16—C11	121.21 (8)
C5—C6—H6A	120.6	C15—C16—H16A	119.4
C7—C6—H6A	120.6	C11—C16—H16A	119.4
C2—C7—C6	120.16 (7)	O2—C17—H17A	109.5
C2—C7—C8	111.85 (7)	O2—C17—H17B	109.5
C6—C7—C8	127.99 (7)	H17A—C17—H17B	109.5
C7—C8—C9	103.06 (6)	O2—C17—H17C	109.5
C7—C8—H8A	111.2	H17A—C17—H17C	109.5
C9—C8—H8A	111.2	H17B—C17—H17C	109.5
C7—C8—H8B	111.2	O3—C18—H18A	109.5
C9—C8—H8B	111.2	O3—C18—H18B	109.5
H8A—C8—H8B	109.1	H18A—C18—H18B	109.5
C10—C9—C1	119.86 (7)	O3—C18—H18C	109.5
C10—C9—C8	131.47 (7)	H18A—C18—H18C	109.5
C1—C9—C8	108.67 (6)	H18B—C18—H18C	109.5
C9—C10—C11	130.91 (7)		
O1—C1—C2—C7	179.73 (8)	C5—C6—C7—C2	0.24 (11)
C9—C1—C2—C7	-0.12 (8)	C5—C6—C7—C8	-178.92 (7)
O1—C1—C2—C3	0.09 (13)	C2—C7—C8—C9	-1.30 (8)
C9—C1—C2—C3	-179.76 (7)	C6—C7—C8—C9	177.91 (8)
C7—C2—C3—C4	-0.86 (11)	O1—C1—C9—C10	-0.86 (13)

supplementary materials

C1—C2—C3—C4	178.74 (7)	C2—C1—C9—C10	179.00 (7)
C17—O2—C4—C3	-14.67 (11)	O1—C1—C9—C8	179.44 (8)
C17—O2—C4—C5	165.99 (7)	C2—C1—C9—C8	-0.71 (8)
C2—C3—C4—O2	179.62 (7)	C7—C8—C9—C10	-178.48 (8)
C2—C3—C4—C5	-1.07 (11)	C7—C8—C9—C1	1.18 (8)
C18—O3—C5—C6	-1.11 (12)	C1—C9—C10—C11	-179.66 (8)
C18—O3—C5—C4	179.19 (7)	C8—C9—C10—C11	-0.03 (15)
O2—C4—C5—O3	1.72 (10)	C9—C10—C11—C12	-1.40 (14)
C3—C4—C5—O3	-177.66 (7)	C9—C10—C11—C16	178.97 (8)
O2—C4—C5—C6	-178.00 (7)	C16—C11—C12—C13	-0.44 (12)
C3—C4—C5—C6	2.62 (12)	C10—C11—C12—C13	179.93 (8)
O3—C5—C6—C7	178.15 (7)	C11—C12—C13—C14	0.46 (13)
C4—C5—C6—C7	-2.16 (12)	C12—C13—C14—C15	0.04 (13)
C3—C2—C7—C6	1.31 (11)	C13—C14—C15—C16	-0.54 (13)
C1—C2—C7—C6	-178.36 (7)	C14—C15—C16—C11	0.55 (12)
C3—C2—C7—C8	-179.41 (7)	C12—C11—C16—C15	-0.06 (12)
C1—C2—C7—C8	0.92 (9)	C10—C11—C16—C15	179.60 (7)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of C2—C7 benzene ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C17—H17A \cdots O3 ⁱ	0.96	2.53	3.4107 (11)	152
C18—H18B \cdots O2 ⁱⁱ	0.96	2.58	3.5320 (11)	173
C16—H16A \cdots Cg1 ⁱⁱⁱ	0.93	2.99	3.7224 (9)	137

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $x+1, y, z$; (iii) $x, -y+1/2, z-3/2$.

Fig. 1




Fig. 2

