

(2E)-1-(3-Bromophenyl)-3-(6-methoxy-2-naphthyl)prop-2-en-1-one

William T. A. Harrison,^{a*} A. N. Mayekar,^{b,c}
H. S. Yathirajan^b and B. Narayana^d

^aDepartment of Chemistry, University of Aberdeen, Meston Walk, Aberdeen AB24 3UE, Scotland, ^bDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, ^cSeQuent Scientific Limited, New Mangalore 575 011, India, and ^dDepartment of Chemistry, Mangalore University, Mangalagangotri 574 199, India

Correspondence e-mail: w.harrison@abdn.ac.uk


Received 9 August 2010; accepted 31 August 2010

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; R factor = 0.082; wR factor = 0.163; data-to-parameter ratio = 17.0.

In the title compound, $\text{C}_{20}\text{H}_{15}\text{BrO}_2$, the prop-2-en-1-one fragment is substantially twisted [$\text{C}-\text{C}-\text{O} = 23.0$ (11°)]. The dihedral angle between the benzene and naphthalene rings is 44.28 (13°). The only possible directional interactions in the crystal are weak $\text{C}-\text{H}\cdots\pi$ contacts, which generate (001) sheets.

Related literature

For related structures, see: Yathirajan *et al.* (2007*a,b*); Jasinski *et al.* (2009). For background to the non-linear optical properties of chalcones, see: Sarojini *et al.* (2006). For reference structural data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{15}\text{BrO}_2$
 $M_r = 367.23$
Orthorhombic, *Pbca*
 $a = 14.0955$ (14) Å
 $b = 6.1295$ (6) Å
 $c = 36.119$ (4) Å

$V = 3120.6$ (5) Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 2.64$ mm⁻¹
 $T = 120$ K
 $0.11 \times 0.09 \times 0.03$ mm

Data collection

Nonius KappaCCD diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2003)
 $T_{\min} = 0.760$, $T_{\max} = 0.925$
28579 measured reflections
3545 independent reflections
1719 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.228$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.082$
 $wR(F^2) = 0.163$
 $S = 1.05$
3545 reflections
209 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.61$ e Å⁻³
 $\Delta\rho_{\min} = -0.63$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg2 is the centroid of the C3–C8 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C4}-\text{H4}\cdots\text{Cg2}^{\text{i}}$	0.95	2.70	3.432 (6)	134
$\text{C7}-\text{H7}\cdots\text{Cg2}^{\text{ii}}$	0.95	2.80	3.520 (6)	134

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{3}{2}, z$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997), *SCALEPACK* and *SORTAV* (Blessing, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

ANM thanks the University of Mysore for providing research facilities. HSY thanks the University of Mysore for sanctioning sabbatical leave.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2699).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
- Bruker (2003). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Jasinski, J. P., Butcher, R. J., Mayekar, A. N., Yathirajan, H. S. & Narayana, B. (2009). *J. Chem. Crystallogr.* **39**, 157–162.
- Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sarojini, B. K., Narayana, B., Ashalatha, B. V., Indira, J. & Lobo, K. G. (2006). *J. Cryst. Growth*, **295**, 54–59.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Yathirajan, H. S., Mayekar, A. N., Narayana, B., Sarojini, B. K. & Bolte, M. (2007*a*). *Acta Cryst.* **E63**, o540–o541.
- Yathirajan, H. S., Mayekar, A. N., Sarojini, B. K., Narayana, B. & Bolte, M. (2007*b*). *Acta Cryst.* **E63**, o1012–o1013.

supplementary materials

Acta Cryst. (2010). E66, o2552 [doi:10.1107/S1600536810035117]

(2*E*)-1-(3-Bromophenyl)-3-(6-methoxy-2-naphthyl)prop-2-en-1-one

W. T. A. Harrison, A. N. Mayekar, H. S. Yathirajan and B. Narayana

Comment

The title compound, (I), (Fig. 1), was prepared as part of our ongoing studies (Yathirajan *et al.*, 2007*a,b*; Jasinski *et al.*, 2009) of substituted phenyl/naphthyl chalcone derivatives as possible candidates for non-linear optical materials (Sarojini *et al.*, 2006). However, (I) crystallizes in a centrosymmetric space group, thus its second-harmonic generation (SHG) response must be zero.

The prop-2-en-1-one (enone) fragment in (I) is substantially twisted, as indicated by the C11—C12—C13—O2 torsion angle of 23.0 (11)°. The dihedral angle between the aromatic ring systems is 44.28 (13)°. Equivalent data for related structures are as follows: (2*E*)-1-(2,4-dichlorophenyl)-3-(6-methoxy-2-naphthyl)prop-2-en-1-one (Yathirajan *et al.*, 2007*a*): -10.9 (2) and 44.94 (4)°; (2*E*)-3-(6-methoxy-2-naphthyl)-1-phenylprop-2-en-1-one (Yathirajan *et al.*, 2007*b*): -15.9 (4) and 14.9 (8)°; (2*E*)-1-(2-hydroxyphenyl)-3-(6-methoxy-2-naphthyl)prop-2-en-1-one (Jasinski *et al.*, 2009): -14.9 (2) and 31.7 (3)°. Otherwise, the bond lengths for (I) fall within their expected ranges (Allen *et al.*, 1987).

In the crystal of (I), the only possible directional interactions between molecules are weak C—H··· π contacts in which the C3—C8 ring of the naphthyl moiety provides both the C—H donor groups and the aromatic acceptor surface (Table 1, Fig. 2). Together, these generate (001) sheets.

Experimental

To a thoroughly stirred solution of 6-methoxy-2-naphthaldehyde (1.86 g, 0.01 mol) and 3-bromoacetophenone (1.99 g, 0.01 mol) in 25 ml methanol, 5 ml of 40% KOH solution was added. The reaction mixture was stirred overnight and the solid separated was collected by filtration. The product obtained was recrystallized from methanol. Colourless slabs of (I) were grown by the slow evaporation of the ethylacetate solution (m.p. 427–429 K).

Refinement

The crystal studied was a weak scatterer, which may correlate with the high R_{int} value. The hydrogen atoms were geometrically placed (C—H = 0.95–0.98 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$. A rotating rigid-group model was applied to the methyl group.

Figures




Fig. 1. View of the molecular structure of (I) showing 50% displacement ellipsoids (arbitrary spheres for the H atoms).




Fig. 2. Partial packing diagram for (I) showing the possible weak C—H... π contacts. All H atoms except H4 and H7 omitted for clarity. Symmetry codes: (i) $1/2-x, y-1/2, z$; (ii) $1-x, 1/2+y, 1/2-z$.

(2E)-1-(3-Bromophenyl)-3-(6-methoxy-2-naphthyl)prop-2-en-1-one

Crystal data

$C_{20}H_{15}BrO_2$

$M_r = 367.23$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 14.0955 (14) \text{ \AA}$

$b = 6.1295 (6) \text{ \AA}$

$c = 36.119 (4) \text{ \AA}$

$V = 3120.6 (5) \text{ \AA}^3$

$Z = 8$

$F(000) = 1488$

$D_x = 1.563 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 55128 reflections

$\theta = 2.9\text{--}27.5^\circ$

$\mu = 2.64 \text{ mm}^{-1}$

$T = 120 \text{ K}$

Slab, colourless

$0.11 \times 0.09 \times 0.03 \text{ mm}$

Data collection

Nonius KappaCCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

ω and ϕ scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2003)

$T_{\min} = 0.760, T_{\max} = 0.925$

28579 measured reflections

3545 independent reflections

1719 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.228$

$\theta_{\max} = 27.5^\circ, \theta_{\min} = 3.1^\circ$

$h = -18 \rightarrow 18$

$k = -7 \rightarrow 7$

$l = -46 \rightarrow 46$

Refinement

Refinement on F^2

Least-squares matrix: full

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

$$R[F^2 > 2\sigma(F^2)] = 0.082$$

$$wR(F^2) = 0.163$$

$$S = 1.05$$

3545 reflections

209 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0343P)^2 + 12.2862P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.61 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.63 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3881 (4)	0.3963 (12)	0.14070 (18)	0.0269 (16)
C2	0.3536 (5)	0.2944 (11)	0.17160 (18)	0.0260 (16)
H2	0.3224	0.1576	0.1695	0.031*
C3	0.3647 (4)	0.3940 (11)	0.20683 (18)	0.0244 (15)
C4	0.3332 (4)	0.2939 (11)	0.23966 (18)	0.0247 (16)
H4	0.3057	0.1526	0.2384	0.030*
C5	0.3411 (4)	0.3933 (11)	0.27329 (17)	0.0257 (15)
H5	0.3192	0.3202	0.2949	0.031*
C6	0.3820 (4)	0.6074 (12)	0.27654 (18)	0.0226 (15)
C7	0.4144 (4)	0.7064 (11)	0.24441 (19)	0.0266 (17)
H7	0.4418	0.8477	0.2460	0.032*
C8	0.4083 (4)	0.6051 (12)	0.20954 (18)	0.0249 (16)
C9	0.4425 (5)	0.7045 (12)	0.1768 (2)	0.0316 (18)
H9	0.4724	0.8432	0.1782	0.038*
C10	0.4332 (4)	0.6034 (12)	0.14329 (19)	0.0290 (16)
H10	0.4569	0.6718	0.1216	0.035*
C11	0.3820 (4)	0.7262 (12)	0.31156 (18)	0.0262 (18)
H11	0.4035	0.8730	0.3107	0.031*
C12	0.3550 (5)	0.6518 (11)	0.34459 (18)	0.0290 (17)
H12	0.3356	0.5039	0.3468	0.035*
C13	0.3544 (5)	0.7926 (12)	0.3780 (2)	0.0305 (17)
C14	0.3642 (4)	0.6878 (12)	0.4150 (2)	0.0276 (17)
C15	0.3486 (5)	0.8158 (12)	0.44671 (19)	0.0312 (17)
H15	0.3274	0.9622	0.4442	0.037*

supplementary materials

C16	0.3638 (5)	0.7308 (12)	0.4810 (2)	0.0313 (17)
C17	0.3927 (5)	0.5156 (12)	0.48559 (19)	0.0290 (18)
H17	0.4014	0.4555	0.5096	0.035*
C18	0.4085 (5)	0.3918 (13)	0.4543 (2)	0.0363 (18)
H18	0.4305	0.2461	0.4571	0.044*
C19	0.3934 (5)	0.4728 (12)	0.4191 (2)	0.035 (2)
H19	0.4029	0.3827	0.3981	0.042*
C20	0.3351 (5)	0.1184 (12)	0.10028 (19)	0.0400 (19)
H20A	0.3318	0.0822	0.0739	0.060*
H20B	0.2707	0.1298	0.1103	0.060*
H20C	0.3697	0.0036	0.1135	0.060*
O1	0.3827 (3)	0.3202 (8)	0.10487 (13)	0.0351 (13)
O2	0.3495 (4)	0.9934 (9)	0.37483 (13)	0.0361 (12)
Br1	0.34948 (6)	0.90763 (13)	0.52413 (2)	0.0403 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.024 (4)	0.033 (4)	0.024 (4)	0.007 (4)	-0.003 (3)	-0.002 (4)
C2	0.027 (4)	0.023 (4)	0.027 (4)	-0.005 (4)	-0.011 (4)	-0.001 (3)
C3	0.022 (4)	0.022 (4)	0.029 (4)	0.002 (4)	0.000 (3)	-0.003 (4)
C4	0.021 (4)	0.019 (4)	0.034 (4)	0.002 (3)	0.004 (3)	0.006 (3)
C5	0.022 (3)	0.031 (4)	0.024 (4)	-0.002 (4)	0.003 (3)	0.008 (4)
C6	0.014 (3)	0.023 (4)	0.030 (4)	-0.001 (3)	-0.002 (3)	-0.001 (4)
C7	0.013 (3)	0.029 (4)	0.038 (5)	-0.001 (3)	0.002 (3)	0.004 (4)
C8	0.018 (3)	0.031 (4)	0.026 (4)	0.005 (4)	-0.001 (3)	0.000 (4)
C9	0.027 (4)	0.034 (4)	0.034 (5)	0.002 (4)	0.003 (4)	0.003 (4)
C10	0.022 (4)	0.032 (4)	0.033 (4)	0.000 (4)	0.003 (3)	0.009 (4)
C11	0.019 (4)	0.028 (4)	0.031 (4)	0.001 (3)	0.000 (3)	-0.009 (4)
C12	0.024 (4)	0.031 (4)	0.032 (4)	-0.006 (4)	0.000 (4)	-0.009 (3)
C13	0.021 (4)	0.032 (5)	0.038 (4)	-0.001 (4)	0.011 (4)	-0.001 (4)
C14	0.016 (4)	0.033 (4)	0.034 (4)	-0.002 (3)	0.002 (3)	-0.004 (4)
C15	0.025 (4)	0.030 (4)	0.039 (4)	-0.004 (4)	0.002 (4)	-0.007 (4)
C16	0.028 (4)	0.034 (4)	0.032 (4)	0.003 (3)	0.003 (4)	-0.004 (4)
C17	0.027 (4)	0.034 (4)	0.026 (4)	-0.003 (3)	-0.001 (3)	0.002 (3)
C18	0.030 (4)	0.038 (5)	0.041 (5)	0.000 (4)	-0.005 (4)	-0.003 (5)
C19	0.023 (4)	0.040 (5)	0.042 (5)	-0.001 (3)	-0.011 (4)	-0.007 (4)
C20	0.059 (5)	0.033 (4)	0.029 (4)	-0.007 (4)	-0.005 (4)	-0.004 (4)
O1	0.045 (3)	0.035 (3)	0.026 (3)	-0.002 (2)	0.001 (2)	-0.001 (3)
O2	0.034 (3)	0.044 (3)	0.031 (3)	0.006 (3)	0.001 (3)	-0.002 (3)
Br1	0.0471 (4)	0.0440 (5)	0.0298 (4)	0.0048 (5)	0.0016 (4)	-0.0062 (4)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.369 (9)	C11—H11	0.9500
C1—O1	1.378 (8)	C12—C13	1.483 (9)
C1—C10	1.422 (9)	C12—H12	0.9500
C2—C3	1.420 (9)	C13—O2	1.238 (8)
C2—H2	0.9500	C13—C14	1.489 (10)

C3—C4	1.407 (9)	C14—C19	1.389 (9)
C3—C8	1.436 (9)	C14—C15	1.406 (9)
C4—C5	1.364 (9)	C15—C16	1.361 (9)
C4—H4	0.9500	C15—H15	0.9500
C5—C6	1.439 (9)	C16—C17	1.391 (10)
C5—H5	0.9500	C16—Br1	1.909 (7)
C6—C7	1.387 (9)	C17—C18	1.379 (9)
C6—C11	1.459 (9)	C17—H17	0.9500
C7—C8	1.407 (9)	C18—C19	1.380 (10)
C7—H7	0.9500	C18—H18	0.9500
C8—C9	1.414 (9)	C19—H19	0.9500
C9—C10	1.366 (9)	C20—O1	1.418 (8)
C9—H9	0.9500	C20—H20A	0.9800
C10—H10	0.9500	C20—H20B	0.9800
C11—C12	1.333 (9)	C20—H20C	0.9800
C2—C1—O1	126.3 (6)	C6—C11—H11	116.4
C2—C1—C10	120.8 (6)	C11—C12—C13	122.0 (7)
O1—C1—C10	112.9 (6)	C11—C12—H12	119.0
C1—C2—C3	119.7 (6)	C13—C12—H12	119.0
C1—C2—H2	120.1	O2—C13—C12	120.3 (7)
C3—C2—H2	120.1	O2—C13—C14	121.1 (7)
C4—C3—C2	122.2 (6)	C12—C13—C14	118.6 (6)
C4—C3—C8	118.1 (6)	C19—C14—C15	119.2 (7)
C2—C3—C8	119.7 (6)	C19—C14—C13	122.2 (7)
C5—C4—C3	122.0 (6)	C15—C14—C13	118.5 (6)
C5—C4—H4	119.0	C16—C15—C14	120.2 (7)
C3—C4—H4	119.0	C16—C15—H15	119.9
C4—C5—C6	120.9 (6)	C14—C15—H15	119.9
C4—C5—H5	119.6	C15—C16—C17	121.2 (7)
C6—C5—H5	119.6	C15—C16—Br1	120.6 (5)
C7—C6—C5	117.6 (6)	C17—C16—Br1	118.2 (5)
C7—C6—C11	120.5 (6)	C18—C17—C16	118.1 (7)
C5—C6—C11	121.7 (6)	C18—C17—H17	120.9
C6—C7—C8	122.4 (6)	C16—C17—H17	120.9
C6—C7—H7	118.8	C17—C18—C19	122.1 (7)
C8—C7—H7	118.8	C17—C18—H18	119.0
C7—C8—C9	122.5 (7)	C19—C18—H18	119.0
C7—C8—C3	119.0 (6)	C18—C19—C14	119.1 (8)
C9—C8—C3	118.5 (6)	C18—C19—H19	120.4
C10—C9—C8	120.8 (7)	C14—C19—H19	120.4
C10—C9—H9	119.6	O1—C20—H20A	109.5
C8—C9—H9	119.6	O1—C20—H20B	109.5
C9—C10—C1	120.4 (7)	H20A—C20—H20B	109.5
C9—C10—H10	119.8	O1—C20—H20C	109.5
C1—C10—H10	119.8	H20A—C20—H20C	109.5
C12—C11—C6	127.2 (7)	H20B—C20—H20C	109.5
C12—C11—H11	116.4	C1—O1—C20	115.6 (6)
O1—C1—C2—C3	-180.0 (6)	C7—C6—C11—C12	-178.5 (7)

supplementary materials

C10—C1—C2—C3	-1.4 (9)	C5—C6—C11—C12	7.2 (10)
C1—C2—C3—C4	-178.0 (6)	C6—C11—C12—C13	-177.3 (6)
C1—C2—C3—C8	2.7 (9)	C11—C12—C13—O2	23.0 (11)
C2—C3—C4—C5	-177.7 (6)	C11—C12—C13—C14	-154.4 (6)
C8—C3—C4—C5	1.6 (9)	O2—C13—C14—C19	-163.5 (7)
C3—C4—C5—C6	0.2 (10)	C12—C13—C14—C19	13.8 (10)
C4—C5—C6—C7	-1.1 (9)	O2—C13—C14—C15	12.6 (10)
C4—C5—C6—C11	173.4 (6)	C12—C13—C14—C15	-170.1 (6)
C5—C6—C7—C8	0.1 (9)	C19—C14—C15—C16	1.2 (10)
C11—C6—C7—C8	-174.5 (6)	C13—C14—C15—C16	-175.0 (6)
C6—C7—C8—C9	-179.1 (6)	C14—C15—C16—C17	-1.5 (10)
C6—C7—C8—C3	1.7 (10)	C14—C15—C16—Br1	176.6 (5)
C4—C3—C8—C7	-2.5 (9)	C15—C16—C17—C18	1.9 (10)
C2—C3—C8—C7	176.8 (6)	Br1—C16—C17—C18	-176.2 (5)
C4—C3—C8—C9	178.3 (6)	C16—C17—C18—C19	-2.2 (10)
C2—C3—C8—C9	-2.4 (9)	C17—C18—C19—C14	2.0 (10)
C7—C8—C9—C10	-178.3 (6)	C15—C14—C19—C18	-1.5 (10)
C3—C8—C9—C10	0.9 (10)	C13—C14—C19—C18	174.6 (6)
C8—C9—C10—C1	0.4 (10)	C2—C1—O1—C20	1.2 (9)
C2—C1—C10—C9	-0.2 (10)	C10—C1—O1—C20	-177.5 (6)
O1—C1—C10—C9	178.6 (6)		

Hydrogen-bond geometry (\AA , $^\circ$)

Cg2 is the centroid of the C3—C8 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4 \cdots Cg2 ⁱ	0.95	2.70	3.432 (6)	134
C7—H7 \cdots Cg2 ⁱⁱ	0.95	2.80	3.520 (6)	134

Symmetry codes: (i) $-x+1/2, y-3/2, z$; (ii) $-x+1, y+1/2, -z+1/2$.

Fig. 1




Fig. 2

