

## 3-Benzoyl-1,1-dibenzylthiourea

N. Gunasekaran,<sup>a</sup> R. Karvembu,<sup>a‡</sup> Seik Weng Ng<sup>b</sup> and Edward R. T. Tiekink<sup>b\*</sup>

<sup>a</sup>Department of Chemistry, National Institute of Technology, Tiruchirappalli 620 015, India, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: edward.tiekink@gmail.com


Received 16 August 2010; accepted 8 September 2010

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.058;  $wR$  factor = 0.157; data-to-parameter ratio = 18.0.

Two independent thiourea molecules comprise the asymmetric unit of the title compound,  $\text{C}_{22}\text{H}_{20}\text{N}_2\text{OS}$ . The central  $\text{N}-\text{C}(=\text{S})\text{N}(\text{H})\text{C}(=\text{O})$  atoms in each molecule are virtually superimposable and each is twisted [ $\text{C}-\text{N}-\text{C}-\text{S}$  torsion angles =  $121.3$  (3) and  $-62.3$  (4) $^\circ$ ]. The molecules differ only in terms of the relative orientations of the benzyl benzene rings [major difference between the  $\text{C}-\text{N}-\text{C}$  torsion angles of  $-146.6$  (3) and  $-132.9$  (3) $^\circ$ ]. The presence of  $\text{N}-\text{H}\cdots\text{S}$  hydrogen bonding leads to the formation of supra-molecular chains along the  $a$  axis. These are consolidated in the crystal packing by  $\text{C}-\text{H}\cdots\text{O}$  interactions. The crystal was found to be a combined non-merohedral and racemic twin (twin law  $\bar{1}00/0\bar{1}0/001$ ), with the fractional contribution of the minor components being approximately 9 and 28%.

### Related literature

For our studies of thiourea and its derivatives, see: Gunasekaran *et al.* (2010). For the biological activity of thiourea derivatives, see: Venkatachalam *et al.* (2004); Yuan *et al.* (2001); Zhou *et al.* (2004). For additional geometric analysis, see: Spek (2009).



<sup>‡</sup> Additional correspondence author, e-mail: kar@nitt.edu.

### Experimental

#### Crystal data

$\text{C}_{22}\text{H}_{20}\text{N}_2\text{OS}$   
 $M_r = 360.46$   
 Monoclinic,  $P2_1$   
 $a = 7.7338$  (5) Å  
 $b = 24.3478$  (16) Å  
 $c = 9.8593$  (6) Å  
 $\beta = 90.074$  (1) $^\circ$   
 $V = 1856.5$  (2) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.19$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.30 \times 0.20 \times 0.10$  mm

#### Data collection

Bruker SMART APEX diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.946$ ,  $T_{\max} = 0.982$   
 17536 measured reflections  
 8469 independent reflections  
 7807 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.055$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.058$   
 $wR(F^2) = 0.157$   
 $S = 1.03$   
 8469 reflections  
 470 parameters  
 1 restraint  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 1.27$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.36$  e Å<sup>-3</sup>  
 Absolute structure: Flack (1983),  
 4101 Friedel pairs  
 Flack parameter: 0.25 (8)

Table 1

Hydrogen-bond geometry (Å,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H2}\cdots\text{S2}$	0.86	2.54	3.334 (3)	154
$\text{N4}-\text{H4}\cdots\text{S1}^i$	0.86	2.54	3.334 (3)	154
$\text{C13}-\text{H13}\cdots\text{O2}^{ii}$	0.95	2.57	3.193 (5)	124
$\text{C14}-\text{H14}\cdots\text{O2}^{ii}$	0.95	2.60	3.207 (5)	122
$\text{C25}-\text{H25}\cdots\text{O1}^{iii}$	0.95	2.55	3.228 (4)	129

Symmetry codes: (i)  $x - 1, y, z$ ; (ii)  $-x + 3, y + \frac{1}{2}, -z$ ; (iii)  $-x + 2, y - \frac{1}{2}, -z$ .

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997), DIAMOND (Brandenburg, 2006) and Qmol (Gans & Shalloway, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

NG thanks the NITT for a Fellowship. The authors also thank the University of Malaya for support of the crystallographic facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5120).

### References

- Brandenburg, K. (2006). DIAMOND. Crystal Impact GbR, Bonn, Germany.  
 Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
 Gans, J. & Shalloway, D. (2001). *J. Mol. Graph. Model.* **19**, 557–559.  
 Gunasekaran, N., Karvembu, R., Ng, S. W. & Tiekink, E. R. T. (2010). *Acta Cryst.* **E66**, o2113.  
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

Venkatachalam, T. K., Mao, C. & Uckun, F. M. (2004). *Bioorg. Med. Chem.* **12**, 4275–4284.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

Yuan, Y. F., Wang, J. T., Gimeno, M. C., Laguna, A. & Jones, P. G. (2001). *Inorg. Chim. Acta*, **324**, 309–317.  
Zhou, W. Q., Li, B. L., Zhu, L. M., Ding, J. G., Yong, Z., Lu, L. & Yang, X. J. (2004). *J. Mol. Struct.* **690**, 145–150.

**supplementary materials**

*Acta Cryst.* (2010). E66, o2572-o2573 [ doi:10.1107/S1600536810036226 ]

### 3-Benzoyl-1,1-dibenzylthiourea

N. Gunasekaran, R. Karvembu, S. W. Ng and E. R. T. Tiekink

#### Comment

The title compound, (I), was investigated in continuation of studies (Gunasekaran *et al.*, 2010) of thiourea and its derivatives, which are useful as anti-tumour, anti-fungal, anti-bacterial, insecticidal, herbicidal, pesticidal agents, and plant-growth regulators (Venkatachalam *et al.*, 2004; Yuan *et al.*, 2001; Zhou *et al.*, 2004).

Two independent molecules comprise the asymmetric unit of (I). The central N–C(=S)N(H)C(=O) atoms of the first independent molecule, Fig. 1, are virtually super-imposable upon those of the second, Fig. 2. The C7–N2–C8–S1 and C29–N4–C30–N3 torsion angles of 121.3 (3) and –62.3 (4) °, respectively, indicate significant twists in the central part of each molecule. The major differences between the molecules relate to the orientations of the benzene rings as indicated in Fig. 3. The major conformational difference is quantified in the C8–N1–C9–C10 and C30–N3–C38–C39 torsion angles –146.6 (3) and –132.9 (3) °, respectively. The r.m.s. deviations for bond distances and angles are 0.0105 Å and 0.651 °, respectively (Spek, 2009).

The most notable feature in the crystal packing is the formation of supramolecular chains along the *a* axis mediated by N–H⋯S hydrogen bonding, Fig. 4 and Table 1. The chains pack in the *ac* plane and stack along the *b* axis with the primary connections along this axis being of the type C–H⋯O, Fig. 5 and Table 1.

#### Experimental

A solution of benzoyl chloride (0.7029 g, 5 mmol) in acetone (50 ml) was added drop wise to a suspension of potassium thiocyanate (0.4859 g, 5 mmol) in anhydrous acetone (50 ml). The reaction mixture was heated under reflux for 45 minutes and then cooled to room temperature. A solution of dibenzyl amine (0.9864 g, 5 mmol) in acetone (30 ml) was added and the resulting mixture was stirred for 2 h. Hydrochloric acid (0.1 N, 300 ml) was added and the resulting white solid was filtered, washed with water and dried *in vacuo*. Single crystals were grown at room temperature from its ethyl acetate solution by the diffusion of diethyl ether vapour. Yield 78%; *M. Pt.* 403 K; FT–IR (KBr)  $\nu(\text{N–H})$  3239,  $\nu(\text{C=O})$  1690,  $\nu(\text{C=S})$  1314  $\text{cm}^{-1}$ .

#### Refinement

Carbon-bound H-atoms were placed in calculated positions (N–H = 0.86 Å and C–H = 0.95 to 0.99 Å) and were included in the refinement in the riding model approximation, with  $U_{\text{iso}}(\text{H})$  set to  $1.2U_{\text{equiv}}(\text{N, C})$ . The maximum and minimum residual electron density peaks of 1.27 and 0.36  $\text{e} \text{Å}^{-3}$ , respectively, were located 1.65 Å and 0.89 Å from the H38*a* and S2 atoms, respectively. As the structure is a non-merohedral twin that belongs to a non-centric space group, the non-merohedral twinning and racemic twinning were treated in combination. The twin law –1 0 0 / 0 –1 0 / 0 0 1 was used as the monoclinic unit cell emulated an orthorhombic unit cell. The Flack parameter refined to 0.25 (8).

## Figures




Fig. 1. The molecular structure of the first independent molecule in (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 2. The molecular structure of the second independent molecule in (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 3. Overlay diagram of the first independent molecule (shown in red) and the second independent molecule (shown in blue).




Fig. 4. Linear supramolecular chain along the *a* axis in (I) mediated by N–H···S hydrogen bonding, shown as blue dashed lines.




Fig. 5. Unit-cell contents shown in projection down the *c* axis in (I). The N–H···S hydrogen bonding and C–H···O contacts are shown as blue and orange dashed lines, respectively.

## 3-Benzoyl-1,1-dibenzylthiourea

### Crystal data

$C_{22}H_{20}N_2OS$

$M_r = 360.46$

Monoclinic,  $P2_1$

Hall symbol: P 2yb

$a = 7.7338$  (5) Å

$b = 24.3478$  (16) Å

$F(000) = 760$

$D_x = 1.290$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 6510 reflections

$\theta = 2.2$ – $28.1^\circ$

$\mu = 0.19$  mm<sup>-1</sup>

$c = 9.8593$  (6) Å  
 $\beta = 90.074$  (1)°  
 $V = 1856.5$  (2) Å<sup>3</sup>  
 $Z = 4$

$T = 100$  K  
 Block, colourless  
 $0.30 \times 0.20 \times 0.10$  mm

*Data collection*

Bruker SMART APEX diffractometer  
 Radiation source: fine-focus sealed tube  
 graphite  
 $\omega$  scans  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.946$ ,  $T_{\max} = 0.982$   
 17536 measured reflections

8469 independent reflections  
 7807 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.055$   
 $\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 0.8^\circ$   
 $h = -10 \rightarrow 10$   
 $k = -31 \rightarrow 31$   
 $l = -12 \rightarrow 11$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.058$   
 $wR(F^2) = 0.157$   
 $S = 1.03$   
 8469 reflections  
 470 parameters  
 1 restraint  
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
 Hydrogen site location: inferred from neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0934P)^2 + 0.2767P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 1.27 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.36 \text{ e \AA}^{-3}$   
 Absolute structure: Flack (1983), 4101 Friedel pairs  
 Flack parameter: 0.25 (8)

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	1.52015 (10)	0.49995 (3)	0.21954 (8)	0.01637 (17)
S2	1.02727 (11)	0.46914 (3)	0.10081 (9)	0.01665 (17)
O1	1.3589 (3)	0.66362 (10)	0.2194 (3)	0.0228 (6)
O2	0.8629 (3)	0.30856 (10)	0.0894 (3)	0.0239 (6)
N1	1.5083 (4)	0.58006 (11)	0.0339 (3)	0.0157 (6)
N2	1.2782 (4)	0.57501 (10)	0.1861 (3)	0.0140 (5)
H2	1.1911	0.5531	0.1843	0.017*
N3	1.0001 (4)	0.38816 (12)	0.2845 (3)	0.0179 (6)
N4	0.7766 (4)	0.39659 (11)	0.1265 (3)	0.0162 (6)
H4	0.6899	0.4187	0.1277	0.019*
C1	1.0820 (4)	0.63713 (13)	0.3027 (3)	0.0154 (6)
C2	0.9587 (5)	0.59635 (14)	0.3240 (4)	0.0204 (7)
H2A	0.9822	0.5597	0.2969	0.024*

## supplementary materials

---

C3	0.8020 (5)	0.60887 (16)	0.3846 (4)	0.0242 (8)
H3	0.7183	0.5809	0.3986	0.029*
C4	0.7672 (5)	0.66233 (15)	0.4247 (4)	0.0239 (8)
H4A	0.6601	0.6708	0.4668	0.029*
C5	0.8880 (5)	0.70339 (15)	0.4038 (4)	0.0240 (8)
H5	0.8635	0.7400	0.4310	0.029*
C6	1.0446 (5)	0.69113 (14)	0.3431 (4)	0.0208 (7)
H6	1.1273	0.7194	0.3287	0.025*
C7	1.2514 (4)	0.62775 (13)	0.2336 (3)	0.0151 (6)
C8	1.4372 (4)	0.55502 (13)	0.1409 (3)	0.0139 (6)
C9	1.6810 (4)	0.56209 (14)	-0.0142 (4)	0.0195 (7)
H9A	1.6654	0.5336	-0.0849	0.023*
H9B	1.7441	0.5451	0.0624	0.023*
C10	1.7895 (4)	0.60781 (13)	-0.0715 (4)	0.0176 (7)
C11	1.8660 (5)	0.60174 (15)	-0.1997 (4)	0.0214 (7)
H11	1.8424	0.5700	-0.2527	0.026*
C12	1.9766 (5)	0.64222 (16)	-0.2494 (4)	0.0234 (8)
H12	2.0286	0.6380	-0.3360	0.028*
C13	2.0104 (5)	0.68838 (16)	-0.1727 (4)	0.0252 (8)
H13	2.0856	0.7159	-0.2069	0.030*
C14	1.9355 (5)	0.69483 (16)	-0.0462 (4)	0.0272 (8)
H14	1.9600	0.7267	0.0059	0.033*
C15	1.8248 (5)	0.65500 (15)	0.0050 (4)	0.0242 (8)
H15	1.7731	0.6598	0.0916	0.029*
C16	1.4092 (5)	0.61491 (14)	-0.0616 (3)	0.0178 (7)
H16A	1.4751	0.6489	-0.0810	0.021*
H16B	1.2978	0.6255	-0.0197	0.021*
C17	1.3751 (4)	0.58415 (14)	-0.1933 (3)	0.0163 (6)
C18	1.4287 (4)	0.60578 (15)	-0.3151 (4)	0.0201 (7)
H18	1.4929	0.6390	-0.3166	0.024*
C19	1.3891 (5)	0.57905 (16)	-0.4367 (4)	0.0224 (7)
H19	1.4247	0.5944	-0.5207	0.027*
C20	1.2973 (5)	0.52985 (17)	-0.4347 (4)	0.0265 (8)
H20	1.2704	0.5115	-0.5171	0.032*
C21	1.2453 (5)	0.50774 (16)	-0.3114 (4)	0.0280 (8)
H21	1.1833	0.4741	-0.3098	0.034*
C22	1.2830 (5)	0.53436 (15)	-0.1908 (4)	0.0235 (8)
H22	1.2468	0.5191	-0.1068	0.028*
C23	0.5896 (4)	0.33461 (14)	-0.0012 (3)	0.0152 (6)
C24	0.5720 (5)	0.28392 (14)	-0.0668 (4)	0.0208 (7)
H24	0.6619	0.2574	-0.0607	0.025*
C25	0.4246 (5)	0.27233 (15)	-0.1403 (4)	0.0237 (8)
H25	0.4145	0.2383	-0.1869	0.028*
C26	0.2900 (5)	0.31064 (17)	-0.1465 (4)	0.0279 (9)
H26	0.1865	0.3019	-0.1938	0.033*
C27	0.3071 (5)	0.36096 (15)	-0.0840 (4)	0.0240 (8)
H27	0.2174	0.3875	-0.0911	0.029*
C28	0.4566 (4)	0.37286 (14)	-0.0104 (4)	0.0196 (7)
H28	0.4677	0.4074	0.0337	0.024*

C29	0.7538 (4)	0.34428 (14)	0.0742 (3)	0.0158 (6)
C30	0.9368 (4)	0.41485 (13)	0.1778 (3)	0.0153 (6)
C31	0.8927 (5)	0.35268 (14)	0.3720 (4)	0.0208 (7)
H31A	0.9526	0.3173	0.3878	0.025*
H31B	0.7812	0.3449	0.3265	0.025*
C32	0.8599 (5)	0.38108 (14)	0.5071 (3)	0.0182 (7)
C33	0.9196 (5)	0.35777 (15)	0.6274 (4)	0.0199 (7)
H33	0.9831	0.3244	0.6251	0.024*
C34	0.8866 (5)	0.38311 (16)	0.7499 (4)	0.0221 (7)
H34	0.9260	0.3668	0.8319	0.026*
C35	0.7964 (5)	0.43216 (16)	0.7538 (4)	0.0246 (8)
H35	0.7749	0.4497	0.8382	0.030*
C36	0.7367 (5)	0.45597 (15)	0.6329 (4)	0.0263 (8)
H36	0.6738	0.4895	0.6351	0.032*
C37	0.7702 (5)	0.43033 (16)	0.5101 (4)	0.0271 (8)
H37	0.7315	0.4466	0.4278	0.033*
C38	1.1725 (5)	0.40062 (15)	0.3372 (4)	0.0230 (8)
H38A	1.1607	0.4216	0.4227	0.028*
H38B	1.2339	0.4242	0.2712	0.028*
C39	1.2794 (5)	0.35003 (15)	0.3639 (4)	0.0212 (7)
C40	1.3815 (5)	0.34745 (16)	0.4800 (4)	0.0244 (8)
H40	1.3763	0.3760	0.5457	0.029*
C41	1.4926 (5)	0.30219 (17)	0.4997 (5)	0.0304 (9)
H41	1.5618	0.3001	0.5793	0.036*
C42	1.5013 (5)	0.26096 (16)	0.4041 (4)	0.0288 (9)
H42	1.5789	0.2311	0.4167	0.035*
C43	1.3972 (6)	0.26289 (17)	0.2894 (5)	0.0308 (9)
H43	1.4020	0.2341	0.2246	0.037*
C44	1.2852 (5)	0.30727 (16)	0.2692 (4)	0.0255 (8)
H44	1.2130	0.3084	0.1912	0.031*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0164 (4)	0.0154 (4)	0.0173 (4)	0.0015 (3)	-0.0008 (3)	0.0026 (3)
S2	0.0153 (4)	0.0147 (4)	0.0199 (4)	-0.0025 (3)	0.0031 (3)	0.0014 (3)
O1	0.0184 (12)	0.0191 (12)	0.0310 (15)	-0.0043 (10)	0.0077 (11)	-0.0040 (11)
O2	0.0204 (13)	0.0205 (12)	0.0309 (15)	0.0037 (10)	-0.0070 (11)	-0.0070 (11)
N1	0.0158 (14)	0.0158 (13)	0.0154 (14)	0.0053 (11)	0.0039 (11)	0.0037 (10)
N2	0.0145 (13)	0.0118 (12)	0.0157 (14)	-0.0011 (10)	0.0029 (11)	-0.0022 (10)
N3	0.0171 (14)	0.0161 (13)	0.0206 (15)	-0.0071 (11)	-0.0011 (12)	0.0006 (11)
N4	0.0141 (13)	0.0154 (13)	0.0191 (14)	-0.0001 (10)	0.0016 (11)	-0.0022 (11)
C1	0.0141 (15)	0.0159 (15)	0.0162 (16)	0.0013 (12)	-0.0003 (12)	-0.0014 (12)
C2	0.0191 (17)	0.0161 (16)	0.0259 (19)	-0.0012 (13)	0.0042 (14)	-0.0053 (13)
C3	0.0165 (17)	0.0258 (18)	0.030 (2)	-0.0058 (14)	0.0078 (15)	-0.0075 (15)
C4	0.0172 (17)	0.0254 (18)	0.029 (2)	0.0016 (14)	0.0050 (15)	-0.0027 (15)
C5	0.027 (2)	0.0170 (17)	0.027 (2)	0.0042 (14)	0.0069 (16)	-0.0024 (14)
C6	0.0264 (19)	0.0152 (16)	0.0209 (18)	-0.0008 (13)	0.0076 (14)	-0.0031 (13)



## supplementary materials

---

C7	0.0164 (16)	0.0163 (15)	0.0125 (15)	0.0004 (12)	-0.0008 (12)	0.0005 (12)
C8	0.0109 (14)	0.0167 (15)	0.0141 (15)	-0.0023 (12)	-0.0017 (12)	-0.0025 (12)
C9	0.0160 (16)	0.0184 (16)	0.0242 (18)	0.0031 (13)	0.0087 (14)	0.0039 (14)
C10	0.0139 (15)	0.0160 (15)	0.0228 (18)	-0.0004 (12)	0.0015 (13)	0.0021 (13)
C11	0.0162 (16)	0.0210 (17)	0.0271 (19)	-0.0009 (13)	0.0072 (14)	0.0023 (14)
C12	0.0160 (16)	0.0312 (19)	0.0231 (19)	0.0009 (14)	0.0057 (14)	0.0039 (15)
C13	0.0193 (18)	0.0255 (18)	0.031 (2)	-0.0004 (14)	0.0003 (15)	0.0101 (16)
C14	0.031 (2)	0.0213 (18)	0.029 (2)	-0.0020 (15)	-0.0030 (16)	0.0002 (15)
C15	0.030 (2)	0.0186 (17)	0.0242 (19)	-0.0007 (14)	0.0005 (15)	0.0011 (14)
C16	0.0190 (16)	0.0179 (15)	0.0163 (16)	0.0058 (12)	0.0048 (13)	0.0052 (13)
C17	0.0130 (15)	0.0192 (15)	0.0166 (16)	0.0035 (12)	0.0004 (12)	0.0034 (13)
C18	0.0195 (17)	0.0227 (17)	0.0180 (17)	-0.0022 (13)	-0.0012 (13)	0.0027 (14)
C19	0.0211 (18)	0.032 (2)	0.0141 (17)	-0.0012 (15)	0.0003 (14)	0.0044 (14)
C20	0.0238 (19)	0.031 (2)	0.0248 (19)	-0.0036 (15)	-0.0021 (15)	-0.0018 (16)
C21	0.031 (2)	0.0244 (19)	0.029 (2)	-0.0068 (15)	-0.0015 (16)	0.0006 (15)
C22	0.0289 (19)	0.0224 (17)	0.0192 (18)	-0.0051 (15)	0.0052 (15)	0.0039 (14)
C23	0.0145 (15)	0.0179 (15)	0.0132 (15)	-0.0013 (12)	0.0003 (12)	-0.0014 (12)
C24	0.0203 (18)	0.0170 (16)	0.0252 (19)	-0.0012 (13)	-0.0008 (15)	-0.0043 (14)
C25	0.0228 (18)	0.0198 (17)	0.028 (2)	-0.0004 (14)	-0.0053 (15)	-0.0087 (15)
C26	0.0183 (18)	0.0298 (19)	0.036 (2)	-0.0001 (15)	-0.0116 (16)	-0.0074 (17)
C27	0.0180 (18)	0.0263 (19)	0.028 (2)	0.0045 (14)	-0.0039 (15)	-0.0032 (15)
C28	0.0181 (17)	0.0207 (16)	0.0199 (17)	0.0000 (13)	-0.0019 (14)	-0.0019 (13)
C29	0.0175 (16)	0.0178 (15)	0.0121 (15)	0.0004 (12)	0.0002 (12)	0.0000 (12)
C30	0.0155 (15)	0.0130 (14)	0.0172 (16)	-0.0015 (12)	0.0033 (12)	-0.0041 (12)
C31	0.0266 (18)	0.0189 (16)	0.0170 (17)	-0.0093 (14)	0.0004 (14)	0.0010 (13)
C32	0.0219 (17)	0.0185 (16)	0.0142 (16)	-0.0085 (13)	0.0002 (13)	-0.0008 (13)
C33	0.0189 (17)	0.0204 (17)	0.0204 (18)	-0.0023 (13)	-0.0012 (13)	0.0030 (13)
C34	0.0172 (17)	0.0334 (19)	0.0156 (17)	0.0018 (14)	-0.0013 (13)	0.0018 (15)
C35	0.0258 (19)	0.029 (2)	0.0187 (18)	-0.0012 (15)	-0.0002 (15)	-0.0018 (15)
C36	0.027 (2)	0.0238 (19)	0.028 (2)	0.0093 (15)	-0.0025 (16)	-0.0043 (15)
C37	0.027 (2)	0.0275 (19)	0.027 (2)	-0.0015 (15)	-0.0071 (16)	0.0053 (16)
C38	0.0217 (17)	0.0205 (17)	0.027 (2)	-0.0067 (14)	-0.0072 (15)	-0.0003 (14)
C39	0.0210 (18)	0.0235 (17)	0.0190 (17)	-0.0059 (14)	-0.0022 (14)	0.0027 (14)
C40	0.0188 (18)	0.0309 (19)	0.0234 (19)	-0.0093 (15)	-0.0040 (14)	0.0052 (15)
C41	0.0180 (18)	0.036 (2)	0.037 (2)	-0.0102 (16)	-0.0069 (16)	0.0093 (17)
C42	0.0201 (18)	0.0238 (18)	0.043 (2)	-0.0037 (15)	0.0075 (17)	0.0078 (16)
C43	0.034 (2)	0.0236 (19)	0.035 (2)	0.0015 (17)	0.0039 (18)	-0.0001 (16)
C44	0.031 (2)	0.0240 (18)	0.0214 (19)	-0.0031 (15)	-0.0013 (15)	-0.0002 (15)

### *Geometric parameters (Å, °)*

S1—C8	1.676 (3)	C18—H18	0.9500
S2—C30	1.678 (3)	C19—C20	1.393 (5)
O1—C7	1.214 (4)	C19—H19	0.9500
O2—C29	1.221 (4)	C20—C21	1.390 (6)
N1—C8	1.337 (4)	C20—H20	0.9500
N1—C16	1.480 (4)	C21—C22	1.385 (5)
N1—C9	1.484 (4)	C21—H21	0.9500
N2—C7	1.382 (4)	C22—H22	0.9500

N2—C8	1.396 (4)	C23—C28	1.390 (5)
N2—H2	0.8600	C23—C24	1.400 (5)
N3—C30	1.329 (4)	C23—C29	1.489 (5)
N3—C38	1.462 (4)	C24—C25	1.380 (5)
N3—C31	1.477 (4)	C24—H24	0.9500
N4—C29	1.386 (4)	C25—C26	1.399 (5)
N4—C30	1.410 (4)	C25—H25	0.9500
N4—H4	0.8600	C26—C27	1.378 (5)
C1—C2	1.393 (5)	C26—H26	0.9500
C1—C6	1.404 (5)	C27—C28	1.395 (5)
C1—C7	1.495 (5)	C27—H27	0.9500
C2—C3	1.386 (5)	C28—H28	0.9500
C2—H2A	0.9500	C31—C32	1.522 (5)
C3—C4	1.387 (5)	C31—H31A	0.9900
C3—H3	0.9500	C31—H31B	0.9900
C4—C5	1.384 (5)	C32—C37	1.386 (5)
C4—H4A	0.9500	C32—C33	1.393 (5)
C5—C6	1.384 (5)	C33—C34	1.381 (5)
C5—H5	0.9500	C33—H33	0.9500
C6—H6	0.9500	C34—C35	1.384 (5)
C9—C10	1.504 (5)	C34—H34	0.9500
C9—H9A	0.9900	C35—C36	1.403 (5)
C9—H9B	0.9900	C35—H35	0.9500
C10—C15	1.401 (5)	C36—C37	1.387 (6)
C10—C11	1.405 (5)	C36—H36	0.9500
C11—C12	1.394 (5)	C37—H37	0.9500
C11—H11	0.9500	C38—C39	1.507 (5)
C12—C13	1.379 (6)	C38—H38A	0.9900
C12—H12	0.9500	C38—H38B	0.9900
C13—C14	1.384 (6)	C39—C40	1.391 (5)
C13—H13	0.9500	C39—C44	1.399 (5)
C14—C15	1.389 (5)	C40—C41	1.411 (6)
C14—H14	0.9500	C40—H40	0.9500
C15—H15	0.9500	C41—C42	1.379 (6)
C16—C17	1.522 (5)	C41—H41	0.9500
C16—H16A	0.9900	C42—C43	1.388 (6)
C16—H16B	0.9900	C42—H42	0.9500
C17—C18	1.376 (5)	C43—C44	1.399 (6)
C17—C22	1.406 (5)	C43—H43	0.9500
C18—C19	1.398 (5)	C44—H44	0.9500
C8—N1—C16	123.3 (3)	C22—C21—C20	120.6 (3)
C8—N1—C9	119.3 (3)	C22—C21—H21	119.7
C16—N1—C9	115.5 (3)	C20—C21—H21	119.7
C7—N2—C8	124.4 (3)	C21—C22—C17	119.6 (3)
C7—N2—H2	117.8	C21—C22—H22	120.2
C8—N2—H2	117.8	C17—C22—H22	120.2
C30—N3—C38	121.0 (3)	C28—C23—C24	119.3 (3)
C30—N3—C31	122.8 (3)	C28—C23—C29	123.9 (3)
C38—N3—C31	115.3 (3)	C24—C23—C29	116.9 (3)

## supplementary materials

---

C29—N4—C30	122.3 (3)	C25—C24—C23	120.2 (3)
C29—N4—H4	118.8	C25—C24—H24	119.9
C30—N4—H4	118.8	C23—C24—H24	119.9
C2—C1—C6	118.9 (3)	C24—C25—C26	120.1 (3)
C2—C1—C7	124.1 (3)	C24—C25—H25	120.0
C6—C1—C7	117.0 (3)	C26—C25—H25	120.0
C3—C2—C1	120.5 (3)	C27—C26—C25	120.2 (3)
C3—C2—H2A	119.8	C27—C26—H26	119.9
C1—C2—H2A	119.8	C25—C26—H26	119.9
C2—C3—C4	119.9 (3)	C26—C27—C28	119.8 (3)
C2—C3—H3	120.0	C26—C27—H27	120.1
C4—C3—H3	120.0	C28—C27—H27	120.1
C5—C4—C3	120.3 (3)	C23—C28—C27	120.5 (3)
C5—C4—H4A	119.9	C23—C28—H28	119.7
C3—C4—H4A	119.9	C27—C28—H28	119.7
C6—C5—C4	120.0 (3)	O2—C29—N4	121.4 (3)
C6—C5—H5	120.0	O2—C29—C23	122.5 (3)
C4—C5—H5	120.0	N4—C29—C23	116.1 (3)
C5—C6—C1	120.3 (3)	N3—C30—N4	116.9 (3)
C5—C6—H6	119.8	N3—C30—S2	126.2 (2)
C1—C6—H6	119.8	N4—C30—S2	116.9 (3)
O1—C7—N2	121.7 (3)	N3—C31—C32	109.9 (3)
O1—C7—C1	122.9 (3)	N3—C31—H31A	109.7
N2—C7—C1	115.4 (3)	C32—C31—H31A	109.7
N1—C8—N2	117.1 (3)	N3—C31—H31B	109.7
N1—C8—S1	124.9 (3)	C32—C31—H31B	109.7
N2—C8—S1	117.9 (2)	H31A—C31—H31B	108.2
N1—C9—C10	113.8 (3)	C37—C32—C33	120.0 (3)
N1—C9—H9A	108.8	C37—C32—C31	119.7 (3)
C10—C9—H9A	108.8	C33—C32—C31	120.3 (3)
N1—C9—H9B	108.8	C34—C33—C32	120.1 (3)
C10—C9—H9B	108.8	C34—C33—H33	119.9
H9A—C9—H9B	107.7	C32—C33—H33	119.9
C15—C10—C11	119.3 (3)	C33—C34—C35	120.2 (3)
C15—C10—C9	120.9 (3)	C33—C34—H34	119.9
C11—C10—C9	119.7 (3)	C35—C34—H34	119.9
C12—C11—C10	120.1 (3)	C34—C35—C36	119.9 (3)
C12—C11—H11	120.0	C34—C35—H35	120.1
C10—C11—H11	120.0	C36—C35—H35	120.1
C13—C12—C11	120.0 (4)	C37—C36—C35	119.6 (3)
C13—C12—H12	120.0	C37—C36—H36	120.2
C11—C12—H12	120.0	C35—C36—H36	120.2
C12—C13—C14	120.4 (4)	C32—C37—C36	120.1 (4)
C12—C13—H13	119.8	C32—C37—H37	119.9
C14—C13—H13	119.8	C36—C37—H37	119.9
C13—C14—C15	120.5 (4)	N3—C38—C39	113.1 (3)
C13—C14—H14	119.8	N3—C38—H38A	109.0
C15—C14—H14	119.8	C39—C38—H38A	109.0
C14—C15—C10	119.8 (4)	N3—C38—H38B	109.0

C14—C15—H15	120.1	C39—C38—H38B	109.0
C10—C15—H15	120.1	H38A—C38—H38B	107.8
N1—C16—C17	110.5 (3)	C40—C39—C44	119.8 (4)
N1—C16—H16A	109.6	C40—C39—C38	119.4 (3)
C17—C16—H16A	109.6	C44—C39—C38	120.7 (3)
N1—C16—H16B	109.6	C39—C40—C41	119.6 (4)
C17—C16—H16B	109.6	C39—C40—H40	120.2
H16A—C16—H16B	108.1	C41—C40—H40	120.2
C18—C17—C22	119.9 (3)	C42—C41—C40	120.3 (4)
C18—C17—C16	120.3 (3)	C42—C41—H41	119.8
C22—C17—C16	119.7 (3)	C40—C41—H41	119.8
C17—C18—C19	120.3 (3)	C41—C42—C43	120.2 (4)
C17—C18—H18	119.8	C41—C42—H42	119.9
C19—C18—H18	119.8	C43—C42—H42	119.9
C20—C19—C18	119.9 (3)	C42—C43—C44	120.0 (4)
C20—C19—H19	120.0	C42—C43—H43	120.0
C18—C19—H19	120.0	C44—C43—H43	120.0
C21—C20—C19	119.6 (4)	C43—C44—C39	120.0 (4)
C21—C20—H20	120.2	C43—C44—H44	120.0
C19—C20—H20	120.2	C39—C44—H44	120.0
C6—C1—C2—C3	-0.2 (5)	C28—C23—C24—C25	0.3 (5)
C7—C1—C2—C3	-177.6 (3)	C29—C23—C24—C25	-179.0 (3)
C1—C2—C3—C4	-0.2 (6)	C23—C24—C25—C26	-1.8 (6)
C2—C3—C4—C5	0.5 (6)	C24—C25—C26—C27	2.8 (7)
C3—C4—C5—C6	-0.3 (6)	C25—C26—C27—C28	-2.3 (6)
C4—C5—C6—C1	-0.1 (6)	C24—C23—C28—C27	0.1 (5)
C2—C1—C6—C5	0.4 (5)	C29—C23—C28—C27	179.4 (3)
C7—C1—C6—C5	177.9 (3)	C26—C27—C28—C23	0.9 (6)
C8—N2—C7—O1	12.0 (5)	C30—N4—C29—O2	11.8 (5)
C8—N2—C7—C1	-168.0 (3)	C30—N4—C29—C23	-168.1 (3)
C2—C1—C7—O1	-178.0 (3)	C28—C23—C29—O2	174.3 (3)
C6—C1—C7—O1	4.6 (5)	C24—C23—C29—O2	-6.4 (5)
C2—C1—C7—N2	2.1 (5)	C28—C23—C29—N4	-5.8 (5)
C6—C1—C7—N2	-175.3 (3)	C24—C23—C29—N4	173.5 (3)
C16—N1—C8—N2	-19.0 (5)	C38—N3—C30—N4	174.2 (3)
C9—N1—C8—N2	177.2 (3)	C31—N3—C30—N4	-17.7 (5)
C16—N1—C8—S1	158.1 (3)	C38—N3—C30—S2	-7.3 (5)
C9—N1—C8—S1	-5.7 (5)	C31—N3—C30—S2	160.7 (3)
C7—N2—C8—N1	-61.4 (4)	C29—N4—C30—N3	-62.3 (4)
C7—N2—C8—S1	121.3 (3)	C29—N4—C30—S2	119.1 (3)
C8—N1—C9—C10	-146.6 (3)	C30—N3—C31—C32	-106.5 (4)
C16—N1—C9—C10	48.4 (4)	C38—N3—C31—C32	62.2 (4)
N1—C9—C10—C15	55.3 (5)	N3—C31—C32—C37	62.2 (4)
N1—C9—C10—C11	-128.9 (3)	N3—C31—C32—C33	-117.7 (3)
C15—C10—C11—C12	0.4 (5)	C37—C32—C33—C34	1.3 (5)
C9—C10—C11—C12	-175.5 (3)	C31—C32—C33—C34	-178.8 (3)
C10—C11—C12—C13	-0.2 (5)	C32—C33—C34—C35	-1.0 (5)
C11—C12—C13—C14	0.1 (6)	C33—C34—C35—C36	0.6 (6)
C12—C13—C14—C15	-0.2 (6)	C34—C35—C36—C37	-0.6 (6)

## supplementary materials

---

C13—C14—C15—C10	0.4 (6)	C33—C32—C37—C36	-1.3 (6)
C11—C10—C15—C14	-0.5 (5)	C31—C32—C37—C36	178.8 (4)
C9—C10—C15—C14	175.3 (3)	C35—C36—C37—C32	0.9 (6)
C8—N1—C16—C17	-103.7 (4)	C30—N3—C38—C39	-132.9 (3)
C9—N1—C16—C17	60.7 (4)	C31—N3—C38—C39	58.2 (4)
N1—C16—C17—C18	-123.1 (3)	N3—C38—C39—C40	-139.2 (3)
N1—C16—C17—C22	59.2 (4)	N3—C38—C39—C44	44.8 (5)
C22—C17—C18—C19	1.2 (5)	C44—C39—C40—C41	1.3 (5)
C16—C17—C18—C19	-176.5 (3)	C38—C39—C40—C41	-174.8 (3)
C17—C18—C19—C20	-1.0 (5)	C39—C40—C41—C42	0.5 (6)
C18—C19—C20—C21	0.2 (6)	C40—C41—C42—C43	-1.8 (6)
C19—C20—C21—C22	0.4 (6)	C41—C42—C43—C44	1.2 (6)
C20—C21—C22—C17	-0.2 (6)	C42—C43—C44—C39	0.7 (6)
C18—C17—C22—C21	-0.7 (5)	C40—C39—C44—C43	-1.9 (6)
C16—C17—C22—C21	177.1 (3)	C38—C39—C44—C43	174.1 (4)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N2—H2 $\cdots$ S2	0.86	2.54	3.334 (3)	154
N4—H4 $\cdots$ S1 <sup>i</sup>	0.86	2.54	3.334 (3)	154
C13—H13 $\cdots$ O2 <sup>ii</sup>	0.95	2.57	3.193 (5)	124
C14—H14 $\cdots$ O2 <sup>ii</sup>	0.95	2.60	3.207 (5)	122
C25—H25 $\cdots$ O1 <sup>iii</sup>	0.95	2.55	3.228 (4)	129

Symmetry codes: (i)  $x-1, y, z$ ; (ii)  $-x+3, y+1/2, -z$ ; (iii)  $-x+2, y-1/2, -z$ .

Fig. 1




Fig. 2




Fig. 3






Fig. 4




Fig. 5

