

## 2,2'-(Disulfanediy)dibenzoic acid–*N,N'*-bis(4-pyridylmethyl)ethanedithioamide (1/1)

Hadi D. Arman,<sup>a</sup> Tyler Miller,<sup>a</sup> Pavel Poplaukhin<sup>b</sup> and Edward R. T. Tiekink<sup>c\*</sup>

<sup>a</sup>Department of Chemistry, The University of Texas at San Antonio, One UTSA Circle, San Antonio, Texas 78249-0698, USA, <sup>b</sup>Chemical Abstracts Service, 2540 Olentangy River Rd, Columbus, Ohio 43202, USA, and <sup>c</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: edward.tiekink@gmail.com


Received 14 September 2010; accepted 14 September 2010

Key indicators: single-crystal X-ray study;  $T = 98$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.063;  $wR$  factor = 0.156; data-to-parameter ratio = 17.0.

The asymmetric unit of the title co-crystal,  $\text{C}_{14}\text{H}_{14}\text{N}_4\text{S}_2 \cdot \text{C}_{14}\text{H}_{10}\text{O}_4\text{S}_2$ , comprises a twisted 2,2'-(disulfanediy)dibenzoic acid molecule [dihedral angle between the benzene rings =  $83.53(14)^\circ$ ] and a U-shaped molecule of *N,N'*-bis(4-pyridylmethyl)ethanedithioamide in which intramolecular  $\text{N}-\text{H} \cdots \text{S}$  hydrogen bonds are observed. Two molecules of each form a centrosymmetric ring, with an extended chair conformation, mediated by carboxyl–pyridine  $\text{O}-\text{H} \cdots \text{N}$  hydrogen bonds between the carboxylic acid groups of two 2,2'-(disulfanediy)dibenzoic acid molecules and pyridine-N atoms of two *N,N'*-bis(4-pyridylmethyl)ethanedithioamide molecules. The tetrameric aggregates are linked into a supramolecular chain along the  $b$  axis via amide–carbonyl  $\text{N}-\text{H} \cdots \text{O}$  hydrogen bonds.

### Related literature

For related studies on co-crystal formation involving 2-[(2-carboxyphenyl)disulfanyl]benzoic acid, see: Broker & Tiekink (2007, 2010); Broker *et al.* (2008); Arman *et al.* (2010).



### Experimental

#### Crystal data

$\text{C}_{14}\text{H}_{14}\text{N}_4\text{S}_2 \cdot \text{C}_{14}\text{H}_{10}\text{O}_4\text{S}_2$   
 $M_r = 608.75$   
 Monoclinic,  $P2_1/c$   
 $a = 18.502(5)$  Å  
 $b = 10.624(3)$  Å  
 $c = 15.026(4)$  Å  
 $\beta = 110.235(5)^\circ$

$V = 2771.3(13)$  Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.39$  mm<sup>-1</sup>  
 $T = 98$  K  
 $0.38 \times 0.26 \times 0.10$  mm

#### Data collection

Rigaku AFC12/SATURN724 diffractometer  
 16758 measured reflections

6348 independent reflections  
 5349 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.050$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.063$   
 $wR(F^2) = 0.156$   
 $S = 1.13$   
 6348 reflections  
 373 parameters  
 4 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.40$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.30$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O}2-\text{H}1\text{o} \cdots \text{N}4^{\text{i}}$	0.84 (3)	1.73 (3)	2.565 (4)	168 (4)
$\text{O}4-\text{H}2\text{o} \cdots \text{N}1^{\text{ii}}$	0.85 (3)	1.76 (3)	2.529 (3)	151 (4)
$\text{N}2-\text{H}1\text{n} \cdots \text{S}2$	0.88 (2)	2.37 (2)	2.930 (3)	121 (2)
$\text{N}3-\text{H}2\text{n} \cdots \text{O}3^{\text{iii}}$	0.88 (3)	2.58 (3)	3.312 (4)	142 (2)
$\text{N}3-\text{H}2\text{n} \cdots \text{S}1$	0.88 (3)	2.45 (3)	2.959 (3)	117 (2)

Symmetry codes: (i)  $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $x, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (iii)  $x, -y + \frac{3}{2}, z - \frac{1}{2}$ .

Data collection: *CrystalClear* (Molecular Structure Corporation & Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPII* (Johnson, 1976) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2714).

### References

- Arman, H. D., Kaulgud, T. & Tiekink, E. R. T. (2010). *Acta Cryst.* **E66**, o2117.  
 Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.  
 Broker, G. A., Bettens, R. P. A. & Tiekink, E. R. T. (2008). *CrystEngComm*, **10**, 879–887.  
 Broker, G. A. & Tiekink, E. R. T. (2007). *CrystEngComm*, **9**, 1096–1109.  
 Broker, G. A. & Tiekink, E. R. T. (2010). *Acta Cryst.* **E66**, o705.  
 Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.  
 Molecular Structure Corporation & Rigaku (2005). *CrystalClear*. MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

**supplementary materials**

*Acta Cryst.* (2010). E66, o2592 [ doi:10.1107/S1600536810036755 ]

## 2,2'-(Disulfanediy)dibenzoic acid-*N,N'*-bis(4-pyridylmethyl)ethanedithioamide (1/1)

H. D. Arman, T. Miller, P. Poplaukhin and E. R. T. Tiekink

### Comment

Co-crystallization of 2-[(2-carboxyphenyl)disulfanyl]benzoic acid with various pyridine donors has led to the isolation of a variety of supramolecular motifs (Broker & Tiekink, 2007; Broker *et al.*, 2008; Broker & Tiekink, 2010; Arman *et al.*, 2010). Herein, the structure determination of the 1:1 co-crystal obtained from the co-crystallization of 2,2'-(disulfanediy)dibenzoic acid with *N,N'*-bis(4-pyridylmethyl)ethanedithioamide is described. The asymmetric unit of the resulting 1:1 co-crystal contains one molecule of 2,2'-(disulfanediy)dibenzoic acid, Fig. 1, and *N,N'*-bis(4-pyridylmethyl)ethanedithioamide, Fig. 2.

In the acid, the expected conformation is observed (Broker & Tiekink, 2007), stabilized in part by two close *S*⋯*O*(carbonyl) interactions, *i.e.* S3⋯O1 = 2.759 (2) Å and S4⋯O3 = 2.687 (3) Å; the dihedral angle formed between the benzene rings = 83.53 (14)°. The molecule of *N,N'*-bis(4-pyridylmethyl)ethanedithioamide adopts a U-shaped conformation as both pyridyl groups lie to the same side of the molecule with the C2—C1—C6—N2 and N3—C9—C10—C11 torsion angles being 40.8 (4) and 146.4 (3)°, respectively. Intramolecular N—H⋯S hydrogen bonds are noted, Table 1.

The components of the co-crystal are linked into a supramolecular ring whereby each carboxylic acid group of two 2,2'-(disulfanediy)dibenzoic acid molecules form an O—H⋯N hydrogen bond with a pyridine-N of two *N,N'*-bis(4-pyridylmethyl)ethanedithioamide molecules, Fig. 3 and Table 1. The ring has an extended chair conformation as seen from the view in Fig. 4. Chairs stack to form a supramolecular chain with the main connections between the tetrameric aggregates being amide-N3—H⋯O3—carbonyl hydrogen bonds, Figs 5 & 6 and Table 1.

### Experimental

Equimolar amounts of 2-[(2-carboxyphenyl)disulfanyl]benzoic acid (Fluka) and *N,N'*-bis(4-pyridylmethyl)ethanedithioamide were dissolved in a 1:1 ethanol/chloroform mixture. Crystals were harvested after a few days of slow evaporation.

### Refinement

C-bound H-atoms were placed in calculated positions (C—H 0.95–0.99 Å) and were included in the refinement in the riding model approximation with  $U_{\text{iso}}(\text{H})$  set to  $1.2U_{\text{eq}}(\text{C})$ . The O- and N-bound H-atoms were located in a difference Fourier map and were refined with distance restraints of O—H 0.84±0.01 Å and N—H = 0.88±0.01 Å, and with  $U_{\text{iso}}(\text{H}) = \gamma U_{\text{eq}}(\text{carrier atom})$ ;  $\gamma = 1.5$  for O and  $\gamma = 1.2$  for N. In the final refinement a low angle reflection evidently effected by the beam stop was omitted, *i.e.* (1 1 0).

Figures




Fig. 1. Molecular structure of 2-[(2-carboxyphenyl)disulfanyl]benzoic acid found in the co-crystal of (I) showing atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 2. Molecular structure of *N,N'*-bis(4-pyridylmethyl)ethanedithioamide found in the co-crystal of (I) showing atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 3. Supramolecular ring in (I). The O—H···N hydrogen bonds are shown as orange dashed lines.




Fig. 4. Supramolecular ring in (I) viewed side-on to emphasize the chair-like conformation. The O—H···N hydrogen bonds are shown as orange dashed lines.




Fig. 5. View of the supramolecular chain along the *b* axis in (I). The O—H···N hydrogen bonds are shown as orange dashed lines. Every second tetrameric aggregate is shown in orange. The N—H···O hydrogen bonds are obscured in this view.



Fig. 6. Side-on view of the supramolecular chain along the *b* axis in (I). The O—H···N and N—H···O hydrogen bonds are shown as orange and green dashed lines, respectively. Every second tetrameric aggregate is shown in orange.

**2,2'-[(Disulfanediyloxy)bis(4-pyridylmethyl)]ethanedithioamide (1/1)**

*Crystal data*

$C_{14}H_{14}N_4S_2 \cdot C_{14}H_{10}O_4S_2$

$M_r = 608.75$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 18.502$  (5) Å

$b = 10.624$  (3) Å

$c = 15.026$  (4) Å

$\beta = 110.235$  (5)°

$V = 2771.3$  (13) Å<sup>3</sup>

$Z = 4$

$F(000) = 1264$

$D_x = 1.459$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71069$  Å

Cell parameters from 11634 reflections

$\theta = 2.2$ – $40.8$ °

$\mu = 0.39$  mm<sup>-1</sup>

$T = 98$  K

Block, red

$0.38 \times 0.26 \times 0.10$  mm

*Data collection*

Rigaku AFC12K/SATURN724 diffractometer	5349 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.050$
graphite	$\theta_{\text{max}} = 27.5^\circ$ , $\theta_{\text{min}} = 2.3^\circ$
$\omega$ scans	$h = -22 \rightarrow 24$
16758 measured reflections	$k = -13 \rightarrow 12$
6348 independent reflections	$l = -19 \rightarrow 12$

*Refinement*

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.063$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.156$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.13$	$w = 1/[\sigma^2(F_o^2) + (0.0595P)^2 + 1.5417P]$
6348 reflections	where $P = (F_o^2 + 2F_c^2)/3$
373 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
4 restraints	$\Delta\rho_{\text{max}} = 0.40 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.30 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.14789 (4)	1.09050 (7)	0.26162 (6)	0.0419 (2)
S2	0.38384 (5)	1.05034 (10)	0.43964 (7)	0.0577 (3)
S3	0.28281 (4)	0.41538 (6)	0.47637 (5)	0.03595 (18)
S4	0.23818 (4)	0.28425 (6)	0.54351 (5)	0.03659 (18)
O1	0.36687 (12)	0.57687 (19)	0.40613 (16)	0.0439 (5)
O2	0.45371 (13)	0.7063 (3)	0.50168 (18)	0.0542 (6)
H1o	0.469 (2)	0.707 (4)	0.455 (2)	0.081*
O3	0.18572 (13)	0.1116 (2)	0.63693 (18)	0.0487 (5)

## supplementary materials

---

O4	0.06256 (14)	0.06603 (19)	0.61426 (18)	0.0482 (6)
H2o	0.0915 (19)	0.016 (3)	0.655 (2)	0.072*
N1	0.10469 (15)	0.6043 (2)	0.23918 (19)	0.0389 (6)
N2	0.22644 (15)	0.9846 (3)	0.42551 (19)	0.0445 (6)
H1n	0.2727 (10)	0.978 (3)	0.4689 (19)	0.053*
N3	0.30049 (14)	1.2123 (2)	0.3103 (2)	0.0401 (6)
H2n	0.2547 (10)	1.233 (3)	0.271 (2)	0.048*
N4	0.48896 (14)	1.2348 (3)	0.12994 (19)	0.0439 (6)
C1	0.13990 (17)	0.8058 (3)	0.3642 (2)	0.0363 (6)
C2	0.19777 (17)	0.7309 (3)	0.3524 (2)	0.0389 (6)
H2	0.2504	0.7483	0.3872	0.047*
C3	0.17793 (17)	0.6311 (3)	0.2896 (2)	0.0376 (6)
H3	0.2176	0.5801	0.2820	0.045*
C4	0.04915 (17)	0.6752 (3)	0.2516 (2)	0.0389 (6)
H4	-0.0030	0.6554	0.2163	0.047*
C5	0.06427 (16)	0.7753 (3)	0.3131 (2)	0.0367 (6)
H5	0.0233	0.8230	0.3205	0.044*
C6	0.16015 (19)	0.9164 (3)	0.4317 (2)	0.0465 (7)
H6A	0.1155	0.9740	0.4164	0.056*
H6B	0.1717	0.8858	0.4974	0.056*
C7	0.22516 (16)	1.0588 (3)	0.3541 (2)	0.0365 (6)
C8	0.30383 (17)	1.1141 (3)	0.3643 (2)	0.0395 (7)
C9	0.36528 (19)	1.2963 (3)	0.3191 (3)	0.0514 (9)
H9A	0.4025	1.2903	0.3847	0.062*
H9B	0.3459	1.3839	0.3093	0.062*
C10	0.40742 (17)	1.2706 (3)	0.2513 (2)	0.0414 (7)
C11	0.43798 (18)	1.3709 (3)	0.2185 (3)	0.0476 (7)
H11	0.4306	1.4543	0.2365	0.057*
C12	0.47942 (19)	1.3496 (3)	0.1591 (3)	0.0509 (8)
H12	0.5018	1.4192	0.1385	0.061*
C13	0.45899 (17)	1.1392 (3)	0.1596 (2)	0.0447 (7)
H13	0.4657	1.0572	0.1384	0.054*
C14	0.41768 (18)	1.1519 (3)	0.2208 (3)	0.0470 (7)
H14	0.3969	1.0802	0.2411	0.056*
C15	0.35166 (15)	0.6427 (2)	0.5491 (2)	0.0336 (6)
C16	0.29850 (16)	0.5498 (2)	0.5523 (2)	0.0337 (6)
C17	0.25988 (17)	0.5629 (3)	0.6169 (2)	0.0384 (6)
H17	0.2235	0.5011	0.6193	0.046*
C18	0.27440 (19)	0.6650 (3)	0.6771 (2)	0.0460 (7)
H18	0.2473	0.6728	0.7203	0.055*
C19	0.32744 (19)	0.7565 (3)	0.6763 (3)	0.0482 (8)
H19	0.3373	0.8259	0.7187	0.058*
C20	0.36576 (17)	0.7446 (3)	0.6123 (2)	0.0407 (7)
H20	0.4024	0.8067	0.6112	0.049*
C21	0.39214 (16)	0.6382 (3)	0.4790 (2)	0.0352 (6)
C22	0.13608 (16)	0.3115 (3)	0.4988 (2)	0.0346 (6)
C23	0.08651 (17)	0.2328 (2)	0.5271 (2)	0.0351 (6)
C24	0.00720 (18)	0.2556 (3)	0.4900 (2)	0.0418 (7)
H24	-0.0266	0.2019	0.5078	0.050*

C25	-0.02326 (18)	0.3530 (3)	0.4287 (2)	0.0450 (7)
H25	-0.0773	0.3669	0.4047	0.054*
C26	0.02518 (19)	0.4301 (3)	0.4026 (2)	0.0466 (7)
H26	0.0046	0.4987	0.3610	0.056*
C27	0.10406 (17)	0.4090 (3)	0.4363 (2)	0.0415 (7)
H27	0.1367	0.4622	0.4161	0.050*
C28	0.11699 (17)	0.1310 (3)	0.5979 (2)	0.0369 (6)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0356 (4)	0.0497 (4)	0.0434 (5)	-0.0029 (3)	0.0175 (3)	0.0003 (3)
S2	0.0405 (4)	0.0839 (6)	0.0467 (5)	-0.0025 (4)	0.0127 (4)	-0.0042 (5)
S3	0.0454 (4)	0.0380 (4)	0.0333 (4)	-0.0046 (3)	0.0250 (3)	-0.0045 (3)
S4	0.0435 (4)	0.0368 (3)	0.0357 (4)	-0.0001 (3)	0.0215 (3)	0.0020 (3)
O1	0.0499 (12)	0.0498 (12)	0.0446 (13)	-0.0110 (9)	0.0321 (11)	-0.0096 (10)
O2	0.0440 (12)	0.0843 (16)	0.0417 (14)	-0.0211 (11)	0.0244 (11)	-0.0103 (13)
O3	0.0585 (14)	0.0433 (11)	0.0520 (14)	0.0064 (10)	0.0292 (12)	0.0153 (10)
O4	0.0566 (14)	0.0402 (11)	0.0561 (15)	-0.0004 (10)	0.0300 (12)	0.0117 (10)
N1	0.0505 (14)	0.0356 (12)	0.0407 (14)	-0.0035 (10)	0.0285 (12)	-0.0009 (10)
N2	0.0473 (14)	0.0528 (15)	0.0324 (14)	-0.0135 (12)	0.0125 (12)	-0.0055 (12)
N3	0.0408 (13)	0.0452 (13)	0.0427 (15)	-0.0110 (11)	0.0253 (12)	-0.0135 (11)
N4	0.0355 (12)	0.0631 (16)	0.0361 (15)	0.0073 (11)	0.0160 (11)	0.0018 (12)
C1	0.0486 (16)	0.0363 (13)	0.0296 (15)	-0.0040 (12)	0.0205 (13)	0.0023 (11)
C2	0.0376 (14)	0.0440 (15)	0.0368 (17)	0.0005 (12)	0.0152 (13)	0.0100 (12)
C3	0.0442 (15)	0.0341 (13)	0.0431 (17)	0.0045 (11)	0.0262 (14)	0.0100 (12)
C4	0.0417 (15)	0.0393 (14)	0.0399 (17)	-0.0047 (12)	0.0197 (13)	-0.0018 (12)
C5	0.0422 (15)	0.0372 (13)	0.0379 (16)	-0.0005 (11)	0.0231 (13)	-0.0007 (12)
C6	0.0573 (19)	0.0523 (17)	0.0356 (17)	-0.0154 (15)	0.0231 (15)	-0.0088 (14)
C7	0.0423 (15)	0.0377 (14)	0.0346 (16)	-0.0074 (11)	0.0200 (13)	-0.0092 (12)
C8	0.0415 (15)	0.0451 (15)	0.0371 (16)	-0.0088 (12)	0.0202 (13)	-0.0160 (13)
C9	0.0545 (19)	0.0558 (18)	0.058 (2)	-0.0189 (15)	0.0367 (18)	-0.0220 (16)
C10	0.0378 (15)	0.0480 (16)	0.0438 (18)	-0.0066 (12)	0.0210 (14)	-0.0065 (13)
C11	0.0480 (17)	0.0491 (17)	0.051 (2)	0.0005 (14)	0.0241 (16)	-0.0003 (15)
C12	0.0492 (18)	0.0589 (19)	0.052 (2)	0.0012 (15)	0.0273 (17)	0.0020 (16)
C13	0.0417 (16)	0.0509 (17)	0.0427 (18)	0.0026 (13)	0.0160 (14)	-0.0075 (14)
C14	0.0492 (17)	0.0469 (16)	0.051 (2)	-0.0080 (13)	0.0248 (16)	-0.0093 (14)
C15	0.0333 (13)	0.0394 (14)	0.0313 (15)	-0.0002 (11)	0.0154 (12)	-0.0004 (11)
C16	0.0378 (14)	0.0370 (13)	0.0300 (15)	0.0011 (11)	0.0165 (12)	-0.0008 (11)
C17	0.0444 (15)	0.0408 (14)	0.0382 (17)	-0.0022 (12)	0.0246 (14)	-0.0037 (12)
C18	0.0568 (18)	0.0523 (17)	0.0413 (18)	-0.0017 (14)	0.0326 (16)	-0.0079 (14)
C19	0.0574 (19)	0.0482 (17)	0.0459 (19)	-0.0053 (14)	0.0267 (16)	-0.0164 (14)
C20	0.0425 (15)	0.0432 (15)	0.0402 (17)	-0.0062 (12)	0.0190 (14)	-0.0060 (13)
C21	0.0363 (14)	0.0389 (14)	0.0347 (16)	0.0012 (11)	0.0178 (12)	0.0008 (12)
C22	0.0418 (14)	0.0375 (13)	0.0291 (15)	-0.0049 (11)	0.0182 (12)	-0.0040 (11)
C23	0.0470 (15)	0.0350 (13)	0.0279 (14)	-0.0040 (11)	0.0187 (12)	-0.0023 (11)
C24	0.0481 (16)	0.0431 (15)	0.0391 (17)	-0.0109 (13)	0.0213 (14)	-0.0024 (13)
C25	0.0416 (16)	0.0552 (18)	0.0354 (17)	-0.0042 (13)	0.0099 (14)	0.0021 (14)

## supplementary materials

---

C26	0.0497 (17)	0.0499 (17)	0.0370 (18)	-0.0036 (14)	0.0108 (14)	0.0088 (13)
C27	0.0449 (16)	0.0456 (16)	0.0359 (17)	-0.0068 (13)	0.0166 (14)	0.0065 (13)
C28	0.0494 (17)	0.0323 (13)	0.0373 (16)	-0.0010 (12)	0.0255 (14)	-0.0020 (11)

### *Geometric parameters (Å, °)*

S1—C7	1.648 (3)	C9—C10	1.506 (4)
S2—C8	1.665 (3)	C9—H9A	0.9900
S3—C16	1.788 (3)	C9—H9B	0.9900
S3—S4	2.0535 (10)	C10—C11	1.375 (4)
S4—C22	1.796 (3)	C10—C14	1.378 (4)
O1—C21	1.219 (4)	C11—C12	1.381 (4)
O2—C21	1.292 (3)	C11—H11	0.9500
O2—H1o	0.84 (3)	C12—H12	0.9500
O3—C28	1.220 (4)	C13—C14	1.390 (4)
O4—C28	1.312 (3)	C13—H13	0.9500
O4—H2o	0.85 (3)	C14—H14	0.9500
N1—C3	1.334 (4)	C15—C20	1.404 (4)
N1—C4	1.338 (4)	C15—C16	1.406 (4)
N2—C7	1.325 (4)	C15—C21	1.490 (4)
N2—C6	1.455 (4)	C16—C17	1.397 (4)
N2—H1n	0.88 (3)	C17—C18	1.379 (4)
N3—C8	1.310 (4)	C17—H17	0.9500
N3—C9	1.464 (4)	C18—C19	1.384 (4)
N3—H2n	0.88 (3)	C18—H18	0.9500
N4—C13	1.308 (4)	C19—C20	1.384 (4)
N4—C12	1.329 (4)	C19—H19	0.9500
C1—C5	1.382 (4)	C20—H20	0.9500
C1—C2	1.394 (4)	C22—C27	1.386 (4)
C1—C6	1.512 (4)	C22—C23	1.411 (4)
C2—C3	1.382 (4)	C23—C24	1.399 (4)
C2—H2	0.9500	C23—C28	1.485 (4)
C3—H3	0.9500	C24—C25	1.371 (4)
C4—C5	1.373 (4)	C24—H24	0.9500
C4—H4	0.9500	C25—C26	1.367 (4)
C5—H5	0.9500	C25—H25	0.9500
C6—H6A	0.9900	C26—C27	1.388 (4)
C6—H6B	0.9900	C26—H26	0.9500
C7—C8	1.527 (4)	C27—H27	0.9500
C16—S3—S4	103.44 (10)	N4—C12—C11	122.0 (3)
C22—S4—S3	104.86 (10)	N4—C12—H12	119.0
C21—O2—H1O	108 (3)	C11—C12—H12	119.0
C28—O4—H2O	98 (3)	N4—C13—C14	123.0 (3)
C3—N1—C4	118.6 (3)	N4—C13—H13	118.5
C7—N2—C6	124.7 (3)	C14—C13—H13	118.5
C7—N2—H1N	113 (2)	C10—C14—C13	118.7 (3)
C6—N2—H1N	122 (2)	C10—C14—H14	120.6
C8—N3—C9	124.6 (3)	C13—C14—H14	120.6
C8—N3—H2N	117 (2)	C20—C15—C16	119.1 (3)



C9—N3—H2N	118 (2)	C20—C15—C21	118.8 (2)
C13—N4—C12	118.7 (3)	C16—C15—C21	122.0 (2)
C5—C1—C2	118.1 (3)	C17—C16—C15	119.1 (3)
C5—C1—C6	121.5 (3)	C17—C16—S3	120.9 (2)
C2—C1—C6	120.4 (3)	C15—C16—S3	120.0 (2)
C3—C2—C1	119.4 (3)	C18—C17—C16	120.3 (3)
C3—C2—H2	120.3	C18—C17—H17	119.9
C1—C2—H2	120.3	C16—C17—H17	119.9
N1—C3—C2	121.9 (3)	C17—C18—C19	121.6 (3)
N1—C3—H3	119.0	C17—C18—H18	119.2
C2—C3—H3	119.0	C19—C18—H18	119.2
N1—C4—C5	122.9 (3)	C20—C19—C18	118.5 (3)
N1—C4—H4	118.6	C20—C19—H19	120.7
C5—C4—H4	118.6	C18—C19—H19	120.7
C4—C5—C1	119.1 (3)	C19—C20—C15	121.4 (3)
C4—C5—H5	120.4	C19—C20—H20	119.3
C1—C5—H5	120.4	C15—C20—H20	119.3
N2—C6—C1	111.3 (2)	O1—C21—O2	124.4 (3)
N2—C6—H6A	109.4	O1—C21—C15	121.5 (3)
C1—C6—H6A	109.4	O2—C21—C15	114.1 (3)
N2—C6—H6B	109.4	C27—C22—C23	118.5 (3)
C1—C6—H6B	109.4	C27—C22—S4	121.5 (2)
H6A—C6—H6B	108.0	C23—C22—S4	120.1 (2)
N2—C7—C8	113.5 (3)	C24—C23—C22	118.6 (3)
N2—C7—S1	124.9 (2)	C24—C23—C28	119.9 (2)
C8—C7—S1	121.6 (2)	C22—C23—C28	121.5 (3)
N3—C8—C7	113.9 (3)	C25—C24—C23	122.0 (3)
N3—C8—S2	125.8 (2)	C25—C24—H24	119.0
C7—C8—S2	120.3 (2)	C23—C24—H24	119.0
N3—C9—C10	115.3 (3)	C26—C25—C24	119.1 (3)
N3—C9—H9A	108.4	C26—C25—H25	120.4
C10—C9—H9A	108.4	C24—C25—H25	120.4
N3—C9—H9B	108.4	C25—C26—C27	120.7 (3)
C10—C9—H9B	108.4	C25—C26—H26	119.7
H9A—C9—H9B	107.5	C27—C26—H26	119.7
C11—C10—C14	117.9 (3)	C22—C27—C26	121.1 (3)
C11—C10—C9	118.4 (3)	C22—C27—H27	119.4
C14—C10—C9	123.7 (3)	C26—C27—H27	119.4
C10—C11—C12	119.6 (3)	O3—C28—O4	124.2 (3)
C10—C11—H11	120.2	O3—C28—C23	122.8 (3)
C12—C11—H11	120.2	O4—C28—C23	113.0 (3)
C16—S3—S4—C22	88.06 (13)	C20—C15—C16—S3	-177.1 (2)
C5—C1—C2—C3	1.1 (4)	C21—C15—C16—S3	4.5 (4)
C6—C1—C2—C3	-179.4 (3)	S4—S3—C16—C17	-20.3 (3)
C4—N1—C3—C2	-1.3 (4)	S4—S3—C16—C15	158.2 (2)
C1—C2—C3—N1	0.3 (4)	C15—C16—C17—C18	-0.6 (4)
C3—N1—C4—C5	0.8 (4)	S3—C16—C17—C18	177.9 (2)
N1—C4—C5—C1	0.6 (4)	C16—C17—C18—C19	-0.5 (5)
C2—C1—C5—C4	-1.5 (4)	C17—C18—C19—C20	0.7 (5)

## supplementary materials

C6—C1—C5—C4	178.9 (3)	C18—C19—C20—C15	0.1 (5)
C7—N2—C6—C1	73.4 (4)	C16—C15—C20—C19	-1.2 (5)
C5—C1—C6—N2	-139.7 (3)	C21—C15—C20—C19	177.3 (3)
C2—C1—C6—N2	40.8 (4)	C20—C15—C21—O1	-157.7 (3)
C6—N2—C7—C8	-177.5 (3)	C16—C15—C21—O1	20.7 (4)
C6—N2—C7—S1	2.1 (4)	C20—C15—C21—O2	20.7 (4)
C9—N3—C8—C7	168.0 (2)	C16—C15—C21—O2	-160.9 (3)
C9—N3—C8—S2	-11.6 (4)	S3—S4—C22—C27	-3.0 (3)
N2—C7—C8—N3	-162.1 (3)	S3—S4—C22—C23	177.0 (2)
S1—C7—C8—N3	18.3 (3)	C27—C22—C23—C24	0.8 (4)
N2—C7—C8—S2	17.4 (3)	S4—C22—C23—C24	-179.2 (2)
S1—C7—C8—S2	-162.20 (17)	C27—C22—C23—C28	-176.5 (3)
C8—N3—C9—C10	98.3 (4)	S4—C22—C23—C28	3.5 (4)
N3—C9—C10—C11	146.4 (3)	C22—C23—C24—C25	-1.2 (4)
N3—C9—C10—C14	-34.1 (5)	C28—C23—C24—C25	176.1 (3)
C14—C10—C11—C12	-2.0 (5)	C23—C24—C25—C26	0.3 (5)
C9—C10—C11—C12	177.6 (3)	C24—C25—C26—C27	1.1 (5)
C13—N4—C12—C11	-1.0 (5)	C23—C22—C27—C26	0.6 (5)
C10—C11—C12—N4	2.1 (5)	S4—C22—C27—C26	-179.5 (3)
C12—N4—C13—C14	-0.2 (5)	C25—C26—C27—C22	-1.5 (5)
C11—C10—C14—C13	0.8 (5)	C24—C23—C28—O3	-175.5 (3)
C9—C10—C14—C13	-178.7 (3)	C22—C23—C28—O3	1.7 (4)
N4—C13—C14—C10	0.3 (5)	C24—C23—C28—O4	3.9 (4)
C20—C15—C16—C17	1.4 (4)	C22—C23—C28—O4	-178.8 (3)
C21—C15—C16—C17	-177.0 (3)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2—H1o $\cdots$ N4 <sup>i</sup>	0.84 (3)	1.73 (3)	2.565 (4)	168 (4)
O4—H2o $\cdots$ N1 <sup>ii</sup>	0.85 (3)	1.76 (3)	2.529 (3)	151 (4)
N2—H1n $\cdots$ S2	0.88 (2)	2.37 (2)	2.930 (3)	121 (2)
N3—H2n $\cdots$ O3 <sup>iii</sup>	0.88 (3)	2.58 (3)	3.312 (4)	142 (2)
N3—H2n $\cdots$ S1	0.88 (3)	2.45 (3)	2.959 (3)	117 (2)

Symmetry codes: (i)  $-x+1, y-1/2, -z+1/2$ ; (ii)  $x, -y+1/2, z+1/2$ ; (iii)  $x, -y+3/2, z-1/2$ .

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6

