

2,2'-(Disulfanediy)dibenzoic acid–2,9-dimethylphenanthroline–tetrahydrofuran (1/2/1)

Hadi D. Arman,^a Trupta Kaulgud^a and Edward R. T. Tiekink^{b*}

^aDepartment of Chemistry, The University of Texas at San Antonio, One UTSA Circle, San Antonio, Texas 78249-0698, USA, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: edward.tiekink@gmail.com


Received 14 September 2010; accepted 16 September 2010

Key indicators: single-crystal X-ray study; $T = 98$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.068; wR factor = 0.169; data-to-parameter ratio = 17.3.

The asymmetric unit of the title co-crystal solvate, $\text{C}_{14}\text{H}_{10}\text{O}_4\text{S}_2 \cdot 2\text{C}_{14}\text{H}_{12}\text{N}_2 \cdot \text{C}_4\text{H}_8\text{O}$, comprises a 2,2'-(disulfanediy)dibenzoic acid molecule, two molecules of 2,9-dimethylphenanthroline and a tetrahydrofuran (THF) solvent molecule. Each end of the twisted diacid [dihedral angle between the benzene rings = 74.33 (17°)] forms a strong $\text{O}-\text{H} \cdots \text{N}$ hydrogen bond with a 2,9-dimethylphenanthroline molecule, forming a trimeric aggregate. The crystal structure comprises layers of acid and THF molecules, and layers of 2,9-dimethylphenanthroline molecules that alternate along the a axis, the main connections between them being of the type $\text{C}-\text{H} \cdots \text{O}$.

Related literature

For related studies on co-crystal formation involving 2-[(2-carboxyphenyl)disulfanyl]benzoic acid, see: Broker & Tiekink (2007, 2010); Broker *et al.* (2008). For a co-crystal involving 2,9-dimethylphenanthroline, see: Arman *et al.* (2010).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{10}\text{O}_4\text{S}_2 \cdot 2\text{C}_{14}\text{H}_{12}\text{N}_2 \cdot \text{C}_4\text{H}_8\text{O}$

$M_r = 794.96$

Monoclinic, $P2_1$
 $a = 14.011$ (4) Å
 $b = 8.516$ (3) Å
 $c = 17.403$ (5) Å
 $\beta = 109.637$ (6) $^\circ$
 $V = 1955.7$ (10) Å³

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.19$ mm⁻¹
 $T = 98$ K
 $0.26 \times 0.21 \times 0.10$ mm

Data collection

Rigaku AFC12/SATURN724
diffractometer
13023 measured reflections

8637 independent reflections
7988 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.068$
 $wR(F^2) = 0.169$
 $S = 1.05$
8637 reflections
499 parameters
8 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.83$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.85$ e Å⁻³
Absolute structure: Flack (1983),
3550 Friedel pairs
Flack parameter: 0.01 (9)

Table 1

Hydrogen-bond geometry (Å, $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O}2-\text{H}2\text{o} \cdots \text{N}1^{\text{i}}$	0.85 (4)	1.91 (4)	2.734 (4)	163 (4)
$\text{O}2-\text{H}2\text{o} \cdots \text{N}2^{\text{i}}$	0.85 (4)	2.46 (4)	2.982 (5)	121 (4)
$\text{O}4-\text{H}4\text{o} \cdots \text{N}4^{\text{ii}}$	0.84 (4)	1.86 (3)	2.691 (4)	170 (4)
$\text{C}19-\text{H}19 \cdots \text{O}1^{\text{iii}}$	0.95	2.59	3.448 (5)	150
$\text{C}22-\text{H}22 \cdots \text{O}5$	0.95	2.52	3.383 (7)	150
$\text{C}23-\text{H}23 \cdots \text{O}3$	0.95	2.56	3.345 (5)	140

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 2, y - \frac{1}{2}, -z + 1$; (iii) $-x + 1, y + \frac{1}{2}, -z$.

Data collection: *CrystalClear* (Molecular Structure Corporation & Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2213).

References

- Arman, H. D., Kaulgud, T. & Tiekink, E. R. T. (2010). *Acta Cryst.* **E66**, o2117.
Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Broker, G. A., Bettens, R. P. A. & Tiekink, E. R. T. (2008). *CrystEngComm*, **10**, 879–887.
Broker, G. A. & Tiekink, E. R. T. (2007). *CrystEngComm*, **9**, 1096–1109.
Broker, G. A. & Tiekink, E. R. T. (2010). *Acta Cryst.* **E66**, o705.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
Molecular Structure Corporation & Rigaku (2005). *CrystalClear*. MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2010). E66, o2602 [doi:10.1107/S1600536810037165]

2,2'-(Disulfanediy)dibenzoic acid-2,9-dimethylphenanthroline-tetrahydrofuran (1/2/1)

H. D. Arman, T. Kaulgud and E. R. T. Tiekink

Comment

As a continuation of studies into the phenomenon of co-crystallization of 2-[(2-carboxyphenyl)disulfanyl]benzoic acid (Broker & Tiekink, 2007; Broker *et al.*, 2008; Broker & Tiekink, 2010; Arman *et al.*, 2010), the co-crystallization of this dithiodibenzoic acid and 2,9-dimethylphenanthroline was investigated. This led to the isolation of the title co-crystal Tetrahydrofuran (thf) solvate.

The crystallographic asymmetric unit of the title compound comprises one molecule of dithiodibenzoic acid (Fig. 1), two molecules of 2,9-dimethylphenanthroline (Figs. 2 and 3), and a solvent thf molecule. The acid adopts the expected conformation (Broker & Tiekink, 2007), stabilized in part by two close *S*⋯*O*(carbonyl) interactions, *i.e.* S1⋯O1 = 2.713 (3) Å and S2⋯O3 = 2.711 (3) Å; the dihedral angle formed between the benzene rings = 74.33 (17)°. Each carboxylic acid-H forms a close hydrogen bond to a phenanthroline-N (Table 1), and in the case of the N1-phenanthroline molecule, a weaker O2—H⋯N2 interaction is noted; the equivalent O4—H⋯N4 contact is longer than 2.66 Å. These interactions result in the formation of a trimeric aggregate, Fig. 4.

In the crystal packing the dithiodibenzoic acid and tetrahydrofuran molecules assemble into layers in the *bc* plane interspersed by layers of 2,9-dimethylphenanthroline molecules, with the most prominent interactions between them being of the type C—H⋯O (see Fig. 5 and Table 1).

Experimental

Gold coloured crystals of the title compound were obtained by the co-crystallization of 2-[(2-carboxyphenyl)disulfanyl]benzoic acid (Fluka, 0.02 mmol) and 2,9-dimethylphenanthroline (ACROS, 0.02 mmol) in tetrahydrofuran. Crystals were obtained by slow evaporation.

Refinement

The O-bound H-atoms were located in a difference Fourier map and were refined with a distance restraint of O—H 0.84 (1) Å, and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. C-bound H-atoms were placed in calculated positions and were included in the refinement in the riding model approximation: C—H 0.95, 0.99 and 0.98 Å for CH, CH₂ and CH₃ H-atoms, respectively, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$, where $k = 1.5$ for CH₃ H-atoms and 1.2 for all other H-atoms. High thermal motion was associated with the tetrahydrofuran molecule but an alternate conformation could not be resolved. The constituent atoms were refined isotropically with O—C and C—C distance restraints of 1.43 (1) and 1.50 (1) Å, respectively. In the final refinement a low angle reflection evidently effected by the beam stop was omitted, *i.e.* ($\bar{1}01$).

Figures




Fig. 1. Molecular structure of 2-[(2-carboxyphenyl)disulfanyl]benzoic acid found in the title co-crystal solvate, showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 2. Molecular structure of first independent molecule of 2,9-dimethylphenanthroline found in the title co-crystal solvate, showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 3. Molecular structure of second independent molecule of 2,9-dimethylphenanthroline found in the title co-crystal solvate, showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 4. The trimeric aggregate, comprising a molecule of 2-[(2-carboxyphenyl)disulfanyl]benzoic acid and the two independent molecules of 2,9-dimethylphenanthroline, sustained by O—H \cdots N hydrogen bonds (dashed lines) in the structure of the title co-crystal solvate [see Table 1 for details].




Fig. 5. Stacking of alternating layers along the *a* axis in the title co-crystal solvate. The O—H \cdots N hydrogen bonding and C—H \cdots O interactions are shown as orange and blue dashed lines, respectively [see Table 1 for details. Hydrogen atoms not involved in intermolecular interactions have been removed for reasons of clarity].

2,2'-(Disulfanediy) dibenzoic acid–2,9-dimethylphenanthroline–tetrahydrofuran (1/2/1)

Crystal data

$C_{14}H_{10}O_4S_2 \cdot 2C_{14}H_{12}N_2 \cdot C_4H_8O$

$M_r = 794.96$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 14.011 (4) \text{ \AA}$

$F(000) = 836$

$D_x = 1.350 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 8113 reflections

$\theta = 2.3\text{--}40.2^\circ$

$b = 8.516 (3) \text{ \AA}$
 $c = 17.403 (5) \text{ \AA}$
 $\beta = 109.637 (6)^\circ$
 $V = 1955.7 (10) \text{ \AA}^3$
 $Z = 2$

$\mu = 0.19 \text{ mm}^{-1}$
 $T = 98 \text{ K}$
 Block, gold
 $0.26 \times 0.21 \times 0.10 \text{ mm}$

Data collection

Rigaku AFC12K/SATURN724 diffractometer
 Radiation source: fine-focus sealed tube
 graphite
 ω scans
 13023 measured reflections
 8637 independent reflections

7988 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$
 $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 2.3^\circ$
 $h = -15 \rightarrow 18$
 $k = -10 \rightarrow 11$
 $l = -22 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.068$
 $wR(F^2) = 0.169$
 $S = 1.05$
 8637 reflections
 499 parameters
 8 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0751P)^2 + 1.2802P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.83 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.85 \text{ e \AA}^{-3}$
 Absolute structure: Flack (1983), 3550 Friedel pairs
 Flack parameter: 0.01 (9)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.42365 (6)	0.42502 (11)	0.23785 (5)	0.02513 (19)

supplementary materials

S2	0.57129 (6)	0.34896 (10)	0.26918 (6)	0.02557 (19)
O1	0.24189 (19)	0.5734 (3)	0.19397 (16)	0.0292 (6)
O2	0.2027 (2)	0.7481 (3)	0.27555 (17)	0.0333 (6)
H2o	0.154 (3)	0.776 (6)	0.234 (2)	0.050*
O3	0.76279 (19)	0.2507 (3)	0.29142 (16)	0.0289 (6)
O4	0.80075 (19)	0.0087 (3)	0.25967 (17)	0.0283 (6)
H4o	0.855 (2)	0.053 (5)	0.261 (3)	0.042*
C1	0.4064 (2)	0.4616 (4)	0.3339 (2)	0.0239 (7)
C2	0.3269 (2)	0.5589 (4)	0.3380 (2)	0.0230 (7)
C3	0.3127 (3)	0.5853 (5)	0.4125 (2)	0.0279 (8)
H3	0.2585	0.6502	0.4146	0.034*
C4	0.3771 (3)	0.5176 (5)	0.4835 (2)	0.0314 (8)
H4	0.3673	0.5359	0.5342	0.038*
C5	0.4560 (3)	0.4227 (5)	0.4797 (2)	0.0294 (8)
H5	0.4998	0.3750	0.5280	0.035*
C6	0.4715 (2)	0.3967 (4)	0.4064 (2)	0.0242 (7)
H6	0.5271	0.3340	0.4052	0.029*
C7	0.2529 (2)	0.6278 (4)	0.2614 (2)	0.0233 (7)
C8	0.5633 (3)	0.1399 (4)	0.2802 (2)	0.0243 (7)
C9	0.6461 (3)	0.0442 (4)	0.2825 (2)	0.0228 (7)
C10	0.6392 (3)	-0.1175 (4)	0.2914 (2)	0.0273 (8)
H10	0.6940	-0.1827	0.2909	0.033*
C11	0.5540 (3)	-0.1847 (5)	0.3008 (2)	0.0323 (8)
H11	0.5516	-0.2946	0.3094	0.039*
C12	0.4721 (3)	-0.0905 (5)	0.2977 (2)	0.0308 (8)
H12	0.4127	-0.1363	0.3028	0.037*
C13	0.4766 (3)	0.0702 (4)	0.2872 (2)	0.0272 (8)
H13	0.4199	0.1337	0.2847	0.033*
C14	0.7423 (3)	0.1115 (4)	0.2779 (2)	0.0236 (7)
N1	1.0745 (2)	0.8741 (3)	0.13399 (18)	0.0244 (6)
N2	0.9787 (3)	0.7029 (4)	0.21862 (19)	0.0282 (7)
C15	1.1222 (3)	0.9556 (4)	0.0924 (2)	0.0276 (8)
C16	1.0706 (3)	1.0298 (4)	0.0182 (2)	0.0280 (7)
H16	1.1070	1.0878	-0.0095	0.034*
C17	0.9668 (3)	1.0182 (4)	-0.0142 (2)	0.0289 (8)
H17	0.9311	1.0682	-0.0644	0.035*
C18	0.9138 (3)	0.9315 (4)	0.0276 (2)	0.0260 (7)
C19	0.8062 (3)	0.9126 (5)	-0.0028 (2)	0.0313 (8)
H19	0.7678	0.9592	-0.0533	0.038*
C20	0.7588 (3)	0.8293 (5)	0.0396 (2)	0.0323 (8)
H20	0.6873	0.8180	0.0182	0.039*
C21	0.8140 (3)	0.7573 (5)	0.1164 (2)	0.0311 (8)
C22	0.7672 (3)	0.6662 (5)	0.1615 (3)	0.0338 (9)
H22	0.6958	0.6518	0.1424	0.041*
C23	0.8265 (3)	0.5986 (5)	0.2336 (3)	0.0360 (9)
H23	0.7962	0.5384	0.2653	0.043*
C24	0.9327 (3)	0.6194 (4)	0.2602 (2)	0.0309 (8)
C25	0.9211 (3)	0.7715 (4)	0.1483 (2)	0.0239 (7)
C26	0.9722 (3)	0.8614 (4)	0.1026 (2)	0.0242 (7)

C27	1.2360 (3)	0.9665 (5)	0.1295 (3)	0.0362 (9)
H27A	1.2548	0.9921	0.1876	0.054*
H27B	1.2610	1.0488	0.1017	0.054*
H27C	1.2663	0.8656	0.1233	0.054*
C28	0.9995 (4)	0.5427 (5)	0.3367 (3)	0.0404 (10)
H28A	1.0702	0.5712	0.3459	0.061*
H28B	0.9919	0.4284	0.3313	0.061*
H28C	0.9801	0.5781	0.3829	0.061*
N3	1.0152 (2)	0.4436 (4)	0.59852 (18)	0.0247 (6)
N4	1.0154 (2)	0.6163 (4)	0.73176 (19)	0.0246 (6)
C29	1.0161 (3)	0.3580 (5)	0.5358 (2)	0.0269 (7)
C30	0.9254 (3)	0.3064 (4)	0.4751 (2)	0.0274 (7)
H30	0.9281	0.2410	0.4316	0.033*
C31	0.8345 (3)	0.3520 (5)	0.4802 (2)	0.0284 (7)
H31	0.7734	0.3204	0.4393	0.034*
C32	0.8309 (3)	0.4456 (4)	0.5455 (2)	0.0266 (7)
C33	0.7390 (3)	0.5060 (5)	0.5519 (3)	0.0345 (9)
H33	0.6766	0.4803	0.5109	0.041*
C34	0.7386 (3)	0.5989 (5)	0.6146 (2)	0.0306 (8)
H34	0.6761	0.6389	0.6166	0.037*
C35	0.8316 (3)	0.6379 (4)	0.6785 (2)	0.0256 (7)
C36	0.8348 (3)	0.7342 (4)	0.7439 (2)	0.0304 (8)
H36	0.7739	0.7763	0.7481	0.036*
C37	0.9266 (3)	0.7683 (5)	0.8025 (2)	0.0309 (8)
H37	0.9294	0.8323	0.8479	0.037*
C38	1.0165 (3)	0.7075 (4)	0.7948 (2)	0.0263 (7)
C39	0.9249 (3)	0.5806 (4)	0.6737 (2)	0.0234 (7)
C40	0.9247 (3)	0.4860 (4)	0.6051 (2)	0.0229 (7)
C41	1.1178 (3)	0.3183 (5)	0.5301 (3)	0.0333 (9)
H41A	1.1660	0.4023	0.5552	0.050*
H41B	1.1119	0.3074	0.4726	0.050*
H41C	1.1420	0.2192	0.5587	0.050*
C42	1.1181 (3)	0.7423 (5)	0.8571 (2)	0.0313 (8)
H42A	1.1562	0.6443	0.8734	0.047*
H42B	1.1088	0.7915	0.9050	0.047*
H42C	1.1557	0.8138	0.8336	0.047*
O5	0.5266 (4)	0.6646 (6)	0.0293 (3)	0.0881 (15)*
C43	0.4873 (6)	0.6959 (12)	-0.0568 (5)	0.106 (3)*
H43A	0.5205	0.7896	-0.0701	0.127*
H43B	0.4994	0.6053	-0.0879	0.127*
C44	0.3809 (5)	0.7227 (10)	-0.0764 (4)	0.089 (2)*
H44A	0.3429	0.6232	-0.0927	0.106*
H44B	0.3563	0.7982	-0.1221	0.106*
C45	0.3658 (5)	0.7886 (8)	-0.0006 (4)	0.0757 (18)*
H45A	0.3626	0.9047	-0.0023	0.091*
H45B	0.3034	0.7468	0.0064	0.091*
C46	0.4594 (5)	0.7317 (10)	0.0663 (4)	0.086 (2)*
H46A	0.4928	0.8205	0.1018	0.103*
H46B	0.4406	0.6521	0.1001	0.103*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0238 (4)	0.0282 (4)	0.0225 (4)	0.0022 (3)	0.0066 (3)	0.0010 (3)
S2	0.0232 (4)	0.0245 (4)	0.0300 (5)	0.0000 (3)	0.0103 (3)	0.0003 (3)
O1	0.0279 (13)	0.0325 (14)	0.0231 (13)	0.0051 (11)	0.0031 (10)	0.0004 (11)
O2	0.0331 (14)	0.0318 (14)	0.0282 (14)	0.0131 (12)	0.0013 (10)	-0.0029 (12)
O3	0.0247 (13)	0.0295 (13)	0.0332 (14)	-0.0019 (11)	0.0105 (10)	-0.0017 (11)
O4	0.0208 (12)	0.0308 (14)	0.0338 (15)	0.0012 (10)	0.0100 (11)	-0.0037 (11)
C1	0.0205 (15)	0.0267 (17)	0.0247 (18)	0.0008 (13)	0.0078 (13)	-0.0005 (13)
C2	0.0181 (15)	0.0241 (17)	0.0249 (17)	0.0001 (12)	0.0049 (12)	-0.0004 (13)
C3	0.0241 (17)	0.0323 (19)	0.0251 (18)	0.0027 (15)	0.0053 (13)	-0.0032 (15)
C4	0.0263 (18)	0.044 (2)	0.0239 (19)	0.0037 (16)	0.0084 (14)	-0.0019 (16)
C5	0.0250 (17)	0.0366 (19)	0.0228 (17)	-0.0004 (15)	0.0029 (13)	0.0014 (16)
C6	0.0162 (14)	0.0308 (19)	0.0239 (17)	0.0011 (12)	0.0043 (12)	0.0010 (13)
C7	0.0193 (15)	0.0226 (16)	0.0272 (18)	-0.0007 (13)	0.0068 (13)	-0.0005 (14)
C8	0.0256 (17)	0.0248 (17)	0.0214 (17)	-0.0037 (13)	0.0063 (13)	-0.0030 (13)
C9	0.0228 (16)	0.0272 (17)	0.0168 (16)	-0.0022 (13)	0.0045 (12)	-0.0016 (13)
C10	0.0261 (17)	0.0273 (19)	0.0257 (18)	0.0025 (13)	0.0048 (13)	0.0003 (13)
C11	0.0299 (19)	0.0268 (19)	0.035 (2)	-0.0048 (14)	0.0034 (15)	-0.0009 (15)
C12	0.0277 (17)	0.0354 (19)	0.0283 (18)	-0.0086 (15)	0.0083 (14)	-0.0029 (16)
C13	0.0217 (16)	0.0313 (19)	0.0252 (18)	-0.0025 (14)	0.0035 (13)	0.0005 (14)
C14	0.0224 (16)	0.0287 (18)	0.0191 (16)	0.0014 (13)	0.0062 (12)	0.0014 (13)
N1	0.0277 (14)	0.0212 (14)	0.0230 (15)	0.0033 (11)	0.0067 (11)	-0.0009 (11)
N2	0.0391 (18)	0.0235 (15)	0.0240 (16)	0.0011 (13)	0.0134 (13)	0.0011 (12)
C15	0.0317 (19)	0.0226 (17)	0.0290 (19)	0.0001 (14)	0.0109 (14)	-0.0020 (14)
C16	0.0353 (19)	0.0247 (17)	0.0277 (18)	0.0009 (14)	0.0154 (15)	0.0027 (14)
C17	0.038 (2)	0.0249 (18)	0.0220 (18)	0.0016 (15)	0.0077 (15)	0.0010 (14)
C18	0.0264 (17)	0.0234 (17)	0.0243 (17)	0.0017 (14)	0.0035 (13)	-0.0030 (14)
C19	0.0314 (19)	0.0303 (18)	0.0266 (18)	0.0046 (16)	0.0026 (14)	-0.0062 (16)
C20	0.0268 (18)	0.033 (2)	0.033 (2)	0.0011 (15)	0.0049 (14)	-0.0031 (16)
C21	0.035 (2)	0.0244 (17)	0.036 (2)	0.0039 (15)	0.0147 (15)	-0.0015 (15)
C22	0.030 (2)	0.0266 (19)	0.048 (2)	-0.0044 (15)	0.0171 (17)	-0.0074 (17)
C23	0.050 (2)	0.029 (2)	0.039 (2)	-0.0038 (18)	0.0269 (19)	-0.0044 (17)
C24	0.044 (2)	0.0213 (17)	0.031 (2)	0.0006 (15)	0.0175 (16)	-0.0005 (15)
C25	0.0318 (18)	0.0192 (16)	0.0221 (17)	0.0008 (13)	0.0108 (13)	-0.0007 (13)
C26	0.0314 (17)	0.0207 (15)	0.0204 (16)	0.0041 (14)	0.0086 (13)	-0.0019 (13)
C27	0.0285 (19)	0.039 (2)	0.040 (2)	0.0000 (16)	0.0105 (16)	0.0030 (17)
C28	0.062 (3)	0.035 (2)	0.026 (2)	0.001 (2)	0.0163 (19)	0.0050 (17)
N3	0.0235 (14)	0.0261 (15)	0.0256 (15)	0.0019 (12)	0.0099 (11)	-0.0006 (12)
N4	0.0258 (14)	0.0224 (14)	0.0271 (15)	0.0008 (11)	0.0111 (11)	0.0022 (12)
C29	0.0271 (17)	0.0280 (17)	0.0263 (18)	0.0003 (15)	0.0097 (13)	0.0024 (15)
C30	0.0296 (18)	0.0293 (18)	0.0236 (18)	-0.0045 (14)	0.0092 (14)	-0.0013 (14)
C31	0.0284 (17)	0.0299 (17)	0.0253 (18)	-0.0022 (15)	0.0068 (13)	-0.0008 (15)
C32	0.0244 (16)	0.0262 (18)	0.0286 (18)	-0.0032 (14)	0.0081 (13)	0.0023 (14)
C33	0.0212 (18)	0.044 (2)	0.036 (2)	-0.0018 (16)	0.0069 (15)	0.0039 (18)
C34	0.0222 (17)	0.0341 (19)	0.035 (2)	-0.0004 (15)	0.0086 (14)	0.0037 (16)

C35	0.0261 (17)	0.0228 (16)	0.0296 (19)	0.0042 (13)	0.0118 (14)	0.0078 (14)
C36	0.0304 (19)	0.0273 (18)	0.036 (2)	0.0088 (15)	0.0144 (15)	0.0049 (16)
C37	0.037 (2)	0.0285 (19)	0.031 (2)	0.0044 (16)	0.0167 (15)	0.0021 (15)
C38	0.0301 (18)	0.0248 (17)	0.0248 (18)	-0.0017 (14)	0.0103 (14)	0.0029 (14)
C39	0.0242 (16)	0.0200 (16)	0.0285 (18)	0.0017 (13)	0.0120 (13)	0.0038 (13)
C40	0.0234 (16)	0.0211 (16)	0.0257 (18)	-0.0009 (13)	0.0101 (13)	0.0020 (13)
C41	0.0319 (19)	0.037 (2)	0.032 (2)	0.0035 (16)	0.0118 (15)	-0.0070 (16)
C42	0.034 (2)	0.0302 (18)	0.029 (2)	-0.0022 (16)	0.0102 (15)	-0.0015 (16)

Geometric parameters (Å, °)

S1—C1	1.795 (4)	C23—H23	0.9500
S1—S2	2.0586 (13)	C24—C28	1.496 (6)
S2—C8	1.798 (4)	C25—C26	1.455 (5)
O1—C7	1.222 (4)	C27—H27A	0.9800
O2—C7	1.312 (4)	C27—H27B	0.9800
O2—H2o	0.85 (4)	C27—H27C	0.9800
O3—C14	1.224 (5)	C28—H28A	0.9800
O4—C14	1.309 (4)	C28—H28B	0.9800
O4—H4o	0.84 (4)	C28—H28C	0.9800
C1—C6	1.398 (5)	N3—C29	1.316 (5)
C1—C2	1.409 (5)	N3—C40	1.360 (4)
C2—C3	1.395 (5)	N4—C38	1.340 (5)
C2—C7	1.506 (5)	N4—C39	1.365 (4)
C3—C4	1.387 (5)	C29—C30	1.423 (5)
C3—H3	0.9500	C29—C41	1.500 (5)
C4—C5	1.389 (5)	C30—C31	1.362 (5)
C4—H4	0.9500	C30—H30	0.9500
C5—C6	1.382 (5)	C31—C32	1.404 (5)
C5—H5	0.9500	C31—H31	0.9500
C6—H6	0.9500	C32—C40	1.417 (5)
C8—C13	1.393 (5)	C32—C33	1.424 (5)
C8—C9	1.408 (5)	C33—C34	1.351 (6)
C9—C10	1.392 (5)	C33—H33	0.9500
C9—C14	1.490 (5)	C34—C35	1.439 (5)
C10—C11	1.383 (5)	C34—H34	0.9500
C10—H10	0.9500	C35—C36	1.391 (5)
C11—C12	1.386 (6)	C35—C39	1.425 (5)
C11—H11	0.9500	C36—C37	1.378 (6)
C12—C13	1.385 (6)	C36—H36	0.9500
C12—H12	0.9500	C37—C38	1.409 (5)
C13—H13	0.9500	C37—H37	0.9500
N1—C15	1.333 (5)	C38—C42	1.502 (5)
N1—C26	1.357 (4)	C39—C40	1.439 (5)
N2—C24	1.326 (5)	C41—H41A	0.9800
N2—C25	1.351 (5)	C41—H41B	0.9800
C15—C16	1.401 (5)	C41—H41C	0.9800
C15—C27	1.509 (5)	C42—H42A	0.9800
C16—C17	1.375 (5)	C42—H42B	0.9800

supplementary materials

C16—H16	0.9500	C42—H42C	0.9800
C17—C18	1.411 (5)	O5—C46	1.426 (7)
C17—H17	0.9500	O5—C43	1.436 (7)
C18—C26	1.417 (5)	C43—C44	1.432 (7)
C18—C19	1.429 (5)	C43—H43A	0.9900
C19—C20	1.348 (6)	C43—H43B	0.9900
C19—H19	0.9500	C44—C45	1.512 (7)
C20—C21	1.437 (5)	C44—H44A	0.9900
C20—H20	0.9500	C44—H44B	0.9900
C21—C22	1.413 (6)	C45—C46	1.511 (7)
C21—C25	1.419 (5)	C45—H45A	0.9900
C22—C23	1.377 (6)	C45—H45B	0.9900
C22—H22	0.9500	C46—H46A	0.9900
C23—C24	1.413 (6)	C46—H46B	0.9900
C1—S1—S2	104.22 (12)	H27A—C27—H27B	109.5
C8—S2—S1	104.28 (13)	C15—C27—H27C	109.5
C7—O2—H2O	113 (4)	H27A—C27—H27C	109.5
C14—O4—H4O	109 (3)	H27B—C27—H27C	109.5
C6—C1—C2	118.3 (3)	C24—C28—H28A	109.5
C6—C1—S1	121.3 (3)	C24—C28—H28B	109.5
C2—C1—S1	120.3 (3)	H28A—C28—H28B	109.5
C3—C2—C1	120.3 (3)	C24—C28—H28C	109.5
C3—C2—C7	119.3 (3)	H28A—C28—H28C	109.5
C1—C2—C7	120.4 (3)	H28B—C28—H28C	109.5
C4—C3—C2	120.5 (3)	C29—N3—C40	119.1 (3)
C4—C3—H3	119.7	C38—N4—C39	119.2 (3)
C2—C3—H3	119.7	N3—C29—C30	122.1 (3)
C3—C4—C5	119.3 (4)	N3—C29—C41	117.0 (3)
C3—C4—H4	120.4	C30—C29—C41	120.8 (3)
C5—C4—H4	120.4	C31—C30—C29	119.0 (4)
C6—C5—C4	120.8 (3)	C31—C30—H30	120.5
C6—C5—H5	119.6	C29—C30—H30	120.5
C4—C5—H5	119.6	C30—C31—C32	120.3 (3)
C5—C6—C1	120.8 (3)	C30—C31—H31	119.9
C5—C6—H6	119.6	C32—C31—H31	119.9
C1—C6—H6	119.6	C31—C32—C40	116.9 (3)
O1—C7—O2	125.0 (3)	C31—C32—C33	123.2 (3)
O1—C7—C2	122.0 (3)	C40—C32—C33	119.8 (4)
O2—C7—C2	113.0 (3)	C34—C33—C32	121.6 (4)
C13—C8—C9	119.1 (3)	C34—C33—H33	119.2
C13—C8—S2	121.2 (3)	C32—C33—H33	119.2
C9—C8—S2	119.7 (3)	C33—C34—C35	120.7 (4)
C10—C9—C8	119.2 (3)	C33—C34—H34	119.6
C10—C9—C14	118.9 (3)	C35—C34—H34	119.6
C8—C9—C14	121.9 (3)	C36—C35—C39	118.2 (3)
C11—C10—C9	121.2 (3)	C36—C35—C34	122.7 (3)
C11—C10—H10	119.4	C39—C35—C34	119.0 (4)
C9—C10—H10	119.4	C37—C36—C35	119.7 (4)
C10—C11—C12	119.5 (4)	C37—C36—H36	120.2

C10—C11—H11	120.3	C35—C36—H36	120.2
C12—C11—H11	120.3	C36—C37—C38	119.6 (4)
C13—C12—C11	120.2 (4)	C36—C37—H37	120.2
C13—C12—H12	119.9	C38—C37—H37	120.2
C11—C12—H12	119.9	N4—C38—C37	121.8 (3)
C12—C13—C8	120.8 (3)	N4—C38—C42	116.9 (3)
C12—C13—H13	119.6	C37—C38—C42	121.3 (4)
C8—C13—H13	119.6	N4—C39—C35	121.5 (3)
O3—C14—O4	124.6 (3)	N4—C39—C40	118.8 (3)
O3—C14—C9	121.3 (3)	C35—C39—C40	119.7 (3)
O4—C14—C9	114.0 (3)	N3—C40—C32	122.4 (3)
C15—N1—C26	118.5 (3)	N3—C40—C39	118.5 (3)
C24—N2—C25	118.3 (3)	C32—C40—C39	119.0 (3)
N1—C15—C16	122.5 (3)	C29—C41—H41A	109.5
N1—C15—C27	116.8 (3)	C29—C41—H41B	109.5
C16—C15—C27	120.8 (4)	H41A—C41—H41B	109.5
C17—C16—C15	119.6 (4)	C29—C41—H41C	109.5
C17—C16—H16	120.2	H41A—C41—H41C	109.5
C15—C16—H16	120.2	H41B—C41—H41C	109.5
C16—C17—C18	119.6 (3)	C38—C42—H42A	109.5
C16—C17—H17	120.2	C38—C42—H42B	109.5
C18—C17—H17	120.2	H42A—C42—H42B	109.5
C17—C18—C26	117.0 (3)	C38—C42—H42C	109.5
C17—C18—C19	122.7 (3)	H42A—C42—H42C	109.5
C26—C18—C19	120.3 (3)	H42B—C42—H42C	109.5
C20—C19—C18	120.6 (4)	C46—O5—C43	108.4 (6)
C20—C19—H19	119.7	C44—C43—O5	106.4 (6)
C18—C19—H19	119.7	C44—C43—H43A	110.5
C19—C20—C21	121.5 (4)	O5—C43—H43A	110.5
C19—C20—H20	119.2	C44—C43—H43B	110.5
C21—C20—H20	119.2	O5—C43—H43B	110.5
C22—C21—C25	117.1 (4)	H43A—C43—H43B	108.6
C22—C21—C20	123.1 (4)	C43—C44—C45	106.8 (6)
C25—C21—C20	119.7 (4)	C43—C44—H44A	110.4
C23—C22—C21	119.1 (4)	C45—C44—H44A	110.4
C23—C22—H22	120.5	C43—C44—H44B	110.4
C21—C22—H22	120.5	C45—C44—H44B	110.4
C22—C23—C24	119.4 (4)	H44A—C44—H44B	108.6
C22—C23—H23	120.3	C46—C45—C44	102.4 (6)
C24—C23—H23	120.3	C46—C45—H45A	111.3
N2—C24—C23	122.7 (4)	C44—C45—H45A	111.3
N2—C24—C28	116.5 (4)	C46—C45—H45B	111.3
C23—C24—C28	120.7 (4)	C44—C45—H45B	111.3
N2—C25—C21	123.4 (3)	H45A—C45—H45B	109.2
N2—C25—C26	118.0 (3)	O5—C46—C45	108.3 (6)
C21—C25—C26	118.7 (3)	O5—C46—H46A	110.0
N1—C26—C18	122.8 (3)	C45—C46—H46A	110.0
N1—C26—C25	118.0 (3)	O5—C46—H46B	110.0
C18—C26—C25	119.2 (3)	C45—C46—H46B	110.0

supplementary materials

C15—C27—H27A	109.5	H46A—C46—H46B	108.4
C15—C27—H27B	109.5		
C1—S1—S2—C8	88.74 (17)	C24—N2—C25—C26	-179.3 (3)
S2—S1—C1—C6	-18.7 (3)	C22—C21—C25—N2	-0.1 (5)
S2—S1—C1—C2	160.5 (2)	C20—C21—C25—N2	-177.7 (3)
C6—C1—C2—C3	-1.8 (5)	C22—C21—C25—C26	178.6 (3)
S1—C1—C2—C3	179.0 (3)	C20—C21—C25—C26	1.0 (5)
C6—C1—C2—C7	-178.2 (3)	C15—N1—C26—C18	-0.2 (5)
S1—C1—C2—C7	2.6 (4)	C15—N1—C26—C25	179.7 (3)
C1—C2—C3—C4	0.7 (6)	C17—C18—C26—N1	-0.5 (5)
C7—C2—C3—C4	177.1 (3)	C19—C18—C26—N1	179.3 (3)
C2—C3—C4—C5	-0.1 (6)	C17—C18—C26—C25	179.7 (3)
C3—C4—C5—C6	0.7 (6)	C19—C18—C26—C25	-0.5 (5)
C4—C5—C6—C1	-1.9 (6)	N2—C25—C26—N1	-1.4 (5)
C2—C1—C6—C5	2.4 (5)	C21—C25—C26—N1	179.9 (3)
S1—C1—C6—C5	-178.4 (3)	N2—C25—C26—C18	178.5 (3)
C3—C2—C7—O1	-157.5 (4)	C21—C25—C26—C18	-0.2 (5)
C1—C2—C7—O1	19.0 (5)	C40—N3—C29—C30	-0.9 (5)
C3—C2—C7—O2	21.9 (5)	C40—N3—C29—C41	178.2 (3)
C1—C2—C7—O2	-161.7 (3)	N3—C29—C30—C31	2.7 (6)
S1—S2—C8—C13	-15.5 (3)	C41—C29—C30—C31	-176.5 (4)
S1—S2—C8—C9	165.0 (2)	C29—C30—C31—C32	-1.5 (6)
C13—C8—C9—C10	0.2 (5)	C30—C31—C32—C40	-1.1 (5)
S2—C8—C9—C10	179.7 (3)	C30—C31—C32—C33	175.7 (4)
C13—C8—C9—C14	-178.0 (3)	C31—C32—C33—C34	-178.3 (4)
S2—C8—C9—C14	1.4 (4)	C40—C32—C33—C34	-1.5 (6)
C8—C9—C10—C11	-2.3 (5)	C32—C33—C34—C35	-1.1 (6)
C14—C9—C10—C11	176.0 (3)	C33—C34—C35—C36	179.4 (4)
C9—C10—C11—C12	2.9 (6)	C33—C34—C35—C39	1.6 (6)
C10—C11—C12—C13	-1.6 (6)	C39—C35—C36—C37	-1.5 (5)
C11—C12—C13—C8	-0.5 (6)	C34—C35—C36—C37	-179.3 (4)
C9—C8—C13—C12	1.1 (5)	C35—C36—C37—C38	1.2 (6)
S2—C8—C13—C12	-178.4 (3)	C39—N4—C38—C37	-0.4 (5)
C10—C9—C14—O3	-160.7 (3)	C39—N4—C38—C42	179.7 (3)
C8—C9—C14—O3	17.6 (5)	C36—C37—C38—N4	-0.2 (6)
C10—C9—C14—O4	18.5 (5)	C36—C37—C38—C42	179.7 (4)
C8—C9—C14—O4	-163.2 (3)	C38—N4—C39—C35	0.0 (5)
C26—N1—C15—C16	0.7 (5)	C38—N4—C39—C40	178.4 (3)
C26—N1—C15—C27	-179.9 (3)	C36—C35—C39—N4	0.9 (5)
N1—C15—C16—C17	-0.6 (6)	C34—C35—C39—N4	178.8 (3)
C27—C15—C16—C17	-179.9 (4)	C36—C35—C39—C40	-177.4 (3)
C15—C16—C17—C18	-0.1 (5)	C34—C35—C39—C40	0.4 (5)
C16—C17—C18—C26	0.6 (5)	C29—N3—C40—C32	-1.9 (5)
C16—C17—C18—C19	-179.2 (4)	C29—N3—C40—C39	-179.5 (3)
C17—C18—C19—C20	-179.6 (4)	C31—C32—C40—N3	3.0 (5)
C26—C18—C19—C20	0.6 (6)	C33—C32—C40—N3	-174.0 (3)
C18—C19—C20—C21	0.2 (6)	C31—C32—C40—C39	-179.4 (3)
C19—C20—C21—C22	-178.5 (4)	C33—C32—C40—C39	3.6 (5)
C19—C20—C21—C25	-1.0 (6)	N4—C39—C40—N3	-3.7 (5)

C25—C21—C22—C23	0.9 (6)	C35—C39—C40—N3	174.7 (3)
C20—C21—C22—C23	178.5 (4)	N4—C39—C40—C32	178.6 (3)
C21—C22—C23—C24	-1.1 (6)	C35—C39—C40—C32	-3.0 (5)
C25—N2—C24—C23	0.4 (5)	C46—O5—C43—C44	-23.0 (9)
C25—N2—C24—C28	178.7 (3)	O5—C43—C44—C45	28.7 (9)
C22—C23—C24—N2	0.4 (6)	C43—C44—C45—C46	-22.9 (9)
C22—C23—C24—C28	-177.8 (4)	C43—O5—C46—C45	8.1 (9)
C24—N2—C25—C21	-0.6 (5)	C44—C45—C46—O5	8.9 (8)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O2—H2o...N1 ⁱ	0.85 (4)	1.91 (4)	2.734 (4)	163 (4)
O2—H2o...N2 ⁱ	0.85 (4)	2.46 (4)	2.982 (5)	121 (4)
O4—H4o...N4 ⁱⁱ	0.84 (4)	1.86 (3)	2.691 (4)	170 (4)
C19—H19...O1 ⁱⁱⁱ	0.95	2.59	3.448 (5)	150
C22—H22...O5	0.95	2.52	3.383 (7)	150
C23—H23...O3	0.95	2.56	3.345 (5)	140

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+2, y-1/2, -z+1$; (iii) $-x+1, y+1/2, -z$.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

