

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

tert-Butyl 4-cyanophenyl carbonate

Malcolm J. Applewhite

Department of Chemistry and Polymer Science, Stellenbosch University, Private Bag X1, Matieland 7602, South Africa

Correspondence e-mail: malcolma@sun.ac.za


Received 15 September 2010; accepted 27 September 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.034; wR factor = 0.091; data-to-parameter ratio = 16.4.

The title compound, $\text{C}_{12}\text{H}_{13}\text{NO}_3$, was prepared by reacting one equivalent of di-*tert*-butyl dicarbonate with 4-cyanophenol. Herringbone crystal packing is observed and there are no significant intermolecular interactions.

Related literature

For a similar packing arrangement in related structures, see: Girard *et al.* (2005); Nagata *et al.* (2008). For reference structural data, see: Allen *et al.* (1987).



Experimental

Crystal data

 $\text{C}_{12}\text{H}_{13}\text{NO}_3$
 $M_r = 219.23$

 Monoclinic, $P2_1/c$
 $a = 5.7347$ (7) Å

 $b = 14.3237$ (16) Å
 $c = 13.7727$ (16) Å
 $\beta = 101.110$ (1)°
 $V = 1110.1$ (2) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 296$ K
 $0.45 \times 0.25 \times 0.12$ mm

Data collection

 Bruker APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.959$, $T_{\max} = 0.989$

 6536 measured reflections
 2427 independent reflections
 2100 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.017$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.091$
 $S = 1.05$
 2427 reflections

 148 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.31$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.23$ e Å⁻³

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001; Atwood & Barbour, 2003); software used to prepare material for publication: X-SEED (Barbour, 2001; Atwood & Barbour, 2003).

The author would like to thank the Central Analytical Facility (CAF) at the University of Stellenbosch (US) for the use of their diffractometer as well as the US and the NRF for funding.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2315).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Atwood, J. L. & Barbour, L. J. (2003). *Cryst. Growth. Des.* **3**, 3–8.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Girard, M., Murphy, P. & Tsou, N. N. (2005). *Tetrahedron Lett.* **46**, 2449–2452.
- Nagata, Y., Otaka, H. & Chujo, Y. (2008). *Macromolecules*, **41**, 737–740.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, o2681 [doi:10.1107/S1600536810038602]

tert-Butyl 4-cyanophenyl carbonate

M. J. Applewhite

Comment

In the title molecule, all bond lengths after refinement are within the normal values as previously collected by Allen *et al.* (1987).

The two-dimensional herring-bone type packing along the *b* and *c* axis observed has two molecules oriented in opposite directions in each turn. Such an arrangement is similar to that described by Girard *et al.* (2005) and Nagata *et al.* (2008) for related organics.

Experimental

Di-*tert*-butyl dicarbonate (279 mg, 1.28 mmol) was added to 4-cyano phenol in 5 ml of dichloromethane. 4-Dimethylamino pyridine (5.20 mg, 0.043 mmol) as a catalyst and triethylamine (0.263 ml, 1.87 mmol) as a base were also added. The solution was stirred at room temperature for six hours.

The reaction was then worked-up using 20 ml of H₂O and 10 ml of 2M HCl which was repeated. The solution was then dried over MgSO₄ after which the solvent was reduced. The resultant white solid was then dried thoroughly under vacuum.

Refinement

Structure solution and refinement were performed using the *SHELX97* suite of programs (Sheldrick, 2008). The H atoms were placed in calculated positions, using and refined using a riding model, with C—H bond lengths fixed to 0.93 (aromatic CH) or 0.96 Å (methyl CH₃).

Figures




Fig. 1. The molecular structure of the title compound showing the 50% probability ellipsoids for non-H atoms.




Fig. 2. The crystal packing viewed down the *a* axis which shows the herring-bone type packing and the two oppositely oriented molecules in each turn.

tert-Butyl 4-cyanophenyl carbonate

Crystal data

$C_{12}H_{13}NO_3$	$F(000) = 464$
$M_r = 219.23$	$D_x = 1.312 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 3143 reflections
$a = 5.7347 (7) \text{ \AA}$	$\theta = 2.8\text{--}27.5^\circ$
$b = 14.3237 (16) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$c = 13.7727 (16) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 101.110 (1)^\circ$	Plate, colourless
$V = 1110.1 (2) \text{ \AA}^3$	$0.45 \times 0.25 \times 0.12 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEX CCD area-detector diffractometer	2427 independent reflections
Radiation source: fine-focus sealed tube graphite	2100 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.017$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 27.9^\circ$, $\theta_{\text{min}} = 2.1^\circ$
$T_{\text{min}} = 0.959$, $T_{\text{max}} = 0.989$	$h = -7 \rightarrow 7$
6536 measured reflections	$k = -18 \rightarrow 14$
	$l = -17 \rightarrow 15$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.034$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.091$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0448P)^2 + 0.3462P]$
2427 reflections	where $P = (F_o^2 + 2F_c^2)/3$
148 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.31 \text{ e \AA}^{-3}$
0 constraints	$\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.33241 (18)	0.70351 (7)	0.16270 (7)	0.0241 (2)

C2	0.4174 (2)	0.69212 (8)	0.24446 (8)	0.0182 (2)
C3	0.52163 (19)	0.67936 (8)	0.34783 (8)	0.0163 (2)
C4	0.74719 (19)	0.71664 (8)	0.38553 (8)	0.0180 (2)
H4	0.8322	0.7469	0.3438	0.022*
C5	0.84213 (19)	0.70792 (8)	0.48552 (8)	0.0187 (2)
H5	0.9911	0.7325	0.5117	0.022*
C6	0.7118 (2)	0.66212 (8)	0.54586 (8)	0.0170 (2)
C7	0.4902 (2)	0.62391 (8)	0.50998 (8)	0.0187 (2)
H7	0.4072	0.5931	0.5521	0.022*
C8	0.3944 (2)	0.63258 (8)	0.40994 (8)	0.0182 (2)
H8	0.2459	0.6073	0.3843	0.022*
O9	0.79952 (14)	0.65945 (6)	0.64836 (6)	0.0199 (2)
C10	0.97513 (19)	0.59643 (8)	0.68019 (8)	0.0159 (2)
O11	1.04663 (14)	0.53999 (6)	0.62887 (6)	0.0211 (2)
O12	1.04148 (14)	0.61088 (6)	0.77662 (5)	0.01731 (19)
C13	1.22277 (19)	0.54939 (8)	0.83729 (8)	0.0161 (2)
C14	1.2463 (2)	0.59433 (9)	0.93865 (8)	0.0210 (3)
H14A	1.3021	0.6573	0.9359	0.032*
H14B	1.3575	0.5594	0.9860	0.032*
H14C	1.0943	0.5947	0.9581	0.032*
C15	1.1255 (2)	0.45053 (8)	0.83660 (9)	0.0203 (2)
H15A	0.9751	0.4516	0.8574	0.030*
H15B	1.2350	0.4123	0.8811	0.030*
H15C	1.1050	0.4254	0.7709	0.030*
C16	1.45548 (19)	0.55369 (8)	0.79966 (8)	0.0189 (2)
H16A	1.4385	0.5204	0.7382	0.028*
H16B	1.5798	0.5257	0.8475	0.028*
H16C	1.4948	0.6177	0.7896	0.028*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0254 (5)	0.0257 (6)	0.0193 (5)	-0.0037 (4)	-0.0003 (4)	0.0010 (4)
C2	0.0180 (5)	0.0158 (5)	0.0206 (6)	-0.0013 (4)	0.0029 (4)	-0.0008 (4)
C3	0.0179 (5)	0.0151 (5)	0.0150 (5)	0.0023 (4)	0.0012 (4)	-0.0004 (4)
C4	0.0181 (5)	0.0186 (5)	0.0176 (5)	-0.0002 (4)	0.0039 (4)	0.0022 (4)
C5	0.0150 (5)	0.0210 (6)	0.0189 (6)	-0.0004 (4)	0.0007 (4)	0.0001 (4)
C6	0.0187 (5)	0.0184 (5)	0.0135 (5)	0.0058 (4)	0.0019 (4)	0.0013 (4)
C7	0.0201 (5)	0.0181 (6)	0.0191 (6)	0.0015 (4)	0.0070 (4)	0.0025 (4)
C8	0.0158 (5)	0.0175 (5)	0.0207 (6)	-0.0009 (4)	0.0022 (4)	-0.0012 (4)
O9	0.0201 (4)	0.0259 (4)	0.0133 (4)	0.0071 (3)	0.0020 (3)	0.0025 (3)
C10	0.0144 (5)	0.0171 (5)	0.0160 (5)	-0.0008 (4)	0.0029 (4)	0.0025 (4)
O11	0.0240 (4)	0.0221 (4)	0.0167 (4)	0.0043 (3)	0.0026 (3)	-0.0016 (3)
O12	0.0181 (4)	0.0204 (4)	0.0130 (4)	0.0056 (3)	0.0018 (3)	0.0013 (3)
C13	0.0153 (5)	0.0179 (5)	0.0144 (5)	0.0036 (4)	0.0008 (4)	0.0032 (4)
C14	0.0238 (6)	0.0241 (6)	0.0144 (5)	0.0055 (5)	0.0016 (4)	0.0012 (4)
C15	0.0192 (5)	0.0197 (6)	0.0221 (6)	0.0001 (4)	0.0040 (4)	0.0035 (4)
C16	0.0168 (5)	0.0191 (6)	0.0208 (6)	0.0010 (4)	0.0037 (4)	0.0014 (4)

supplementary materials

Geometric parameters (Å, °)

N1—C2	1.1493 (15)	C10—O12	1.3254 (13)
C2—C3	1.4455 (15)	O12—C13	1.4899 (13)
C3—C8	1.3979 (16)	C13—C14	1.5194 (15)
C3—C4	1.4032 (15)	C13—C15	1.5214 (16)
C4—C5	1.3855 (15)	C13—C16	1.5231 (15)
C4—H4	0.9300	C14—H14A	0.9600
C5—C6	1.3850 (16)	C14—H14B	0.9600
C5—H5	0.9300	C14—H14C	0.9600
C6—C7	1.3846 (16)	C15—H15A	0.9600
C6—O9	1.4049 (13)	C15—H15B	0.9600
C7—C8	1.3871 (16)	C15—H15C	0.9600
C7—H7	0.9300	C16—H16A	0.9600
C8—H8	0.9300	C16—H16B	0.9600
O9—C10	1.3605 (13)	C16—H16C	0.9600
C10—O11	1.1971 (14)		
N1—C2—C3	178.86 (13)	O12—C13—C14	101.18 (8)
C8—C3—C4	120.57 (10)	O12—C13—C15	109.34 (9)
C8—C3—C2	119.85 (10)	C14—C13—C15	111.57 (9)
C4—C3—C2	119.55 (10)	O12—C13—C16	110.35 (9)
C5—C4—C3	119.42 (10)	C14—C13—C16	111.52 (9)
C5—C4—H4	120.3	C15—C13—C16	112.32 (9)
C3—C4—H4	120.3	C13—C14—H14A	109.5
C6—C5—C4	119.01 (10)	C13—C14—H14B	109.5
C6—C5—H5	120.5	H14A—C14—H14B	109.5
C4—C5—H5	120.5	C13—C14—H14C	109.5
C7—C6—C5	122.50 (10)	H14A—C14—H14C	109.5
C7—C6—O9	118.29 (10)	H14B—C14—H14C	109.5
C5—C6—O9	119.07 (10)	C13—C15—H15A	109.5
C6—C7—C8	118.67 (10)	C13—C15—H15B	109.5
C6—C7—H7	120.7	H15A—C15—H15B	109.5
C8—C7—H7	120.7	C13—C15—H15C	109.5
C7—C8—C3	119.82 (10)	H15A—C15—H15C	109.5
C7—C8—H8	120.1	H15B—C15—H15C	109.5
C3—C8—H8	120.1	C13—C16—H16A	109.5
C10—O9—C6	116.18 (9)	C13—C16—H16B	109.5
O11—C10—O12	129.21 (10)	H16A—C16—H16B	109.5
O11—C10—O9	125.05 (10)	C13—C16—H16C	109.5
O12—C10—O9	105.73 (9)	H16A—C16—H16C	109.5
C10—O12—C13	120.28 (8)	H16B—C16—H16C	109.5
C8—C3—C4—C5	−0.93 (17)	C7—C6—O9—C10	105.80 (12)
C2—C3—C4—C5	177.14 (10)	C5—C6—O9—C10	−78.37 (13)
C3—C4—C5—C6	0.27 (17)	C6—O9—C10—O11	−5.43 (16)
C4—C5—C6—C7	0.48 (17)	C6—O9—C10—O12	175.49 (9)
C4—C5—C6—O9	−175.16 (10)	O11—C10—O12—C13	−2.85 (17)
C5—C6—C7—C8	−0.55 (17)	O9—C10—O12—C13	176.18 (8)
O9—C6—C7—C8	175.12 (10)	C10—O12—C13—C14	177.84 (9)

supplementary materials

C6—C7—C8—C3	-0.12 (17)	C10—O12—C13—C15	-64.35 (12)
C4—C3—C8—C7	0.86 (17)	C10—O12—C13—C16	59.67 (12)
C2—C3—C8—C7	-177.20 (10)		

Fig. 1




Fig. 2

