

## Tris[2-ethoxy-6-(methyliminomethyl)-phenolato- $\kappa^2 N,O^1$ ]cobalt(III) monohydrate

Yin Dan Huang,<sup>a</sup> Shu-Hua Zhang,<sup>a\*</sup> Jiang Ke Qin<sup>b\*</sup> and Fu Li Chen<sup>a</sup>

<sup>a</sup>College of Chemistry and Bioengineering, Guilin University of Technology, Key Laboratory of Non-ferrous Metal Materials and Processing Technology, Ministry of Education, Guilin 541004, People's Republic of China, and <sup>b</sup>School of Chemistry and Chemical Engineering, Guangxi Normal University, Guilin 541004, People's Republic of China

Correspondence e-mail: zsh720108@163.com


Received 29 September 2010; accepted 7 October 2010

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(C-C) = 0.004$  Å; disorder in main residue;  $R$  factor = 0.037;  $wR$  factor = 0.100; data-to-parameter ratio = 14.0.

In the title compound,  $[Co(C_{10}H_{12}NO_2)_3] \cdot H_2O$ , the Co<sup>III</sup> ion is coordinated by three O atoms and three N atoms from three bidentate 2-ethoxy-6-(methyliminomethyl)phenolate ligands in a slightly distorted octahedral environment. The water molecule connects two ligands by O–H···O hydrogen bonds. One terminal methyl group is disordered over two positions, with site-occupancy factors of 0.412 (15) and 0.588 (15).

### Related literature

For Co<sup>III</sup> complexes, see: Park *et al.* (2008); Galezowski *et al.* (2008); Gupta *et al.* (2007). For Schiff-base compounds, see: Gupta & Sutar (2008); Sreenivasulu *et al.* (2005); Zhang & Feng (2010); Zhang *et al.* (2010).



### Experimental

#### Crystal data

$[Co(C_{10}H_{12}NO_2)_3] \cdot H_2O$	$V = 2994.1$ (17) Å <sup>3</sup>
$M_r = 611.56$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 14.022$ (5) Å	$\mu = 0.62$ mm <sup>-1</sup>
$b = 16.969$ (6) Å	$T = 296$ K
$c = 14.233$ (5) Å	$0.42 \times 0.20 \times 0.09$ mm
$\beta = 117.857$ (4)°	

#### Data collection

Bruker SMART CCD area-detector diffractometer	21931 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	5317 independent reflections
$T_{min} = 0.860$ , $T_{max} = 0.944$	3929 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.035$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$	12 restraints
$wR(F^2) = 0.100$	H-atom parameters constrained
$S = 1.01$	$\Delta\rho_{\text{max}} = 0.31$ e Å <sup>-3</sup>
5317 reflections	$\Delta\rho_{\text{min}} = -0.21$ e Å <sup>-3</sup>
379 parameters	

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1W–H1WA···O3	0.85	2.01	2.821 (3)	159
O1W–H1WB···O6	0.85	2.24	3.038 (3)	155

Data collection: *SMART* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The work was supported financially by the Guangxi Key Laboratory for Advanced Materials and New Preparation Technology (No. 0842003–25), the Young Science Foundation of Guangxi Province of China (No. 0832085) and the Doctoral start-up research fund of Guilin University of Technology.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5367).

### References

- Bruker (2004). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Galezowski, W., Kuta, J. & Kozlowski, P. M. (2008). *J. Phys. Chem. B*, **112**, 3177–3183.
- Gupta, S. K., Hitchcock, P. B., Kushwah, Y. S. & Argal, G. S. (2007). *Inorg. Chim. Acta*, **360**, 2145–2152.
- Gupta, K. C. & Sutar, A. K. (2008). *Coord. Chem. Rev.* **252**, 1420–1450.
- Park, J., Lang, K., Abboud, K. A. & Hong, S. (2008). *J. Am. Chem. Soc.* **130**, 16484–16485.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sreenivasulu, B., Vetrichelvan, M., Zhao, F., Gao, S. & Vittal, J. J. (2005). *Eur. J. Inorg. Chem.* pp. 4635–4645.
- Zhang, S. H. & Feng, C. (2010). *J. Mol. Struct.* **977**, 62–66.
- Zhang, S. H., Song, Y., Liang, H. & Zeng, M. H. (2009). *CrystEngComm*, **11**, 865–872.

## **supplementary materials**

*Acta Cryst.* (2010). E66, m1399 [doi:10.1107/S1600536810040043]

## Tris[2-ethoxy-6-(methyliminomethyl)phenolato- $\kappa^2N,O^1$ ]cobalt(III) monohydrate

**Y. D. Huang, S.-H. Zhang, J. K. Qin and F. L. Chen**

### Comment

Schiff base complexes have been studied for many years (Gupta & Sutar, 2008; Sreenivasulu *et al.*, 2005; Zhang *et al.*, 2009; Zhang *et al.*, 2010) and have attracted interest because of their anticancer, catalytic and fluorescent properties. Using 2-hydrogen-3-ethoxy-benzaldehyde, methylamine and  $\text{Co}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ , we have hydrothermally prepared the title compound, (I). As an example of  $\text{Co}^{\text{III}}$  compounds there is a series of  $\text{Co}^{\text{III}}$  complexes that were synthesized and characterized (Park *et al.* 2008; Galezowski *et al.* 2008; Gupta *et al.*, 2007).

In the molecular structure of (I), the  $\text{Co}^{\text{III}}$  ion is coordinated by three O atoms and three N atoms from three bidentate *L* ligand forming a slightly distorted octahedral geometry (Fig. 1). The  $\text{Co}-X$  ( $X = \text{O}, \text{N}$ ) bond lengths lie in the range 1.878 (2)–1.954 (2) Å, and the angles subtended at the  $\text{Co}^{\text{III}}$  atom range from 85.61 (9) to 94.25 (10) °. The Co ion is in the 3+ oxidation state, as evidenced by bond valence summation calculations, charge balance considerations, and the presence of typical bond lengths for  $\text{Co}^{\text{III}}$  (Park, *et al.* 2008; Galezowski, *et al.* 2008; Gupta *et al.*, 2007). The water molecule connects two ligands by O–H···O hydrogen bonds (Table 1).

### Experimental

Complex I was prepared from a mixture of 2-hydrogen-3-ethoxy-benzaldehyde (0.166 g, 1 mmol), methylamine solution (0.5 ml),  $\text{Co}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$  (0.360 g, 1 mmol), and methanol (8 ml) sealed in a 15 ml Teflon-lined stainless steel bomb, and kept at 120 °C for 72 h under autogenous pressure. After the reaction was slowly cooled to room temperature, red block crystals were produced (yield: 52%, based on 2-hydrogen-3-ethoxy-benzaldehyde). Anal. Calcd for  $\text{C}_{30}\text{H}_{38}\text{CoN}_3\text{O}_7$  (%): C, 58.91; H, 6.26; N, 6.88. Found (%): C, 58.88; H, 6.31; N, 6.92.

### Refinement

H atoms were positioned geometrically and were treated as riding atoms, with C–H distances of 0.93–0.97 Å, O–H 0.85 Å with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}_{\text{sp}2})$  and  $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C}_{\text{sp}3} \text{ or } \text{O})$ . One terminal methyl group is disordered over two positions with site occupation factors of 0.412 (15) and 0.588 (15), respectively. The displacement parameters of the disordered atoms were restrained to an approximately isotropic behaviour.

# supplementary materials

---

## Figures




Fig. 1. Molecular structure of the title compound with displacement ellipsoids drawn at the 50% probability level.

## Tris[2-ethoxy-6-(methyliminomethyl)phenolato- $\kappa^2N,O^1$ ]cobalt(III) monohydrate

### Crystal data

[Co(C <sub>10</sub> H <sub>12</sub> NO <sub>2</sub> ) <sub>3</sub> ] <sub>·</sub> H <sub>2</sub> O	$F(000) = 1288$
$M_r = 611.56$	$D_x = 1.357 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 7434 reflections
$a = 14.022 (5) \text{ \AA}$	$\theta = 2.8\text{--}28.4^\circ$
$b = 16.969 (6) \text{ \AA}$	$\mu = 0.62 \text{ mm}^{-1}$
$c = 14.233 (5) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 117.857 (4)^\circ$	Block, red
$V = 2994.1 (17) \text{ \AA}^3$	$0.42 \times 0.20 \times 0.09 \text{ mm}$
$Z = 4$	

### Data collection

Bruker SMART CCD area-detector diffractometer	5317 independent reflections
Radiation source: fine-focus sealed tube graphite	3929 reflections with $I > 2\sigma(I)$
phi and $\omega$ scans	$R_{\text{int}} = 0.035$
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 25.1^\circ, \theta_{\text{min}} = 2.8^\circ$
$T_{\text{min}} = 0.860, T_{\text{max}} = 0.944$	$h = -16 \rightarrow 16$
21931 measured reflections	$k = -20 \rightarrow 19$
	$l = -16 \rightarrow 16$

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.037$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.100$	H-atom parameters constrained
$S = 1.01$	$w = 1/[\sigma^2(F_o^2) + (0.0481P)^2 + 0.9938P]$
5317 reflections	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} = 0.001$

379 parameters  $\Delta\rho_{\max} = 0.31 \text{ e } \text{\AA}^{-3}$   
 12 restraints  $\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Co1	0.25696 (2)	0.943185 (18)	0.22915 (2)	0.04238 (12)	
C1	0.45768 (19)	0.96749 (15)	0.22521 (19)	0.0484 (6)	
C2	0.5048 (2)	0.89830 (16)	0.2845 (2)	0.0536 (7)	
C3	0.6109 (2)	0.87721 (19)	0.3075 (2)	0.0698 (9)	
H3	0.6412	0.8315	0.3461	0.084*	
C4	0.6695 (2)	0.9220 (2)	0.2748 (3)	0.0802 (10)	
H4	0.7388	0.9068	0.2896	0.096*	
C5	0.6255 (2)	0.9904 (2)	0.2193 (3)	0.0735 (9)	
H5	0.6666	1.0217	0.1983	0.088*	
C6	0.5225 (2)	1.01357 (16)	0.1942 (2)	0.0560 (7)	
C7	0.4056 (3)	1.0805 (2)	0.0383 (3)	0.0950 (12)	
H7A	0.3539	1.0397	0.0305	0.142*	
H7B	0.4399	1.0657	-0.0046	0.142*	
C9	0.4483 (2)	0.84859 (16)	0.3222 (2)	0.0583 (8)	
H9	0.4858	0.8048	0.3611	0.070*	
C10	0.3095 (3)	0.79479 (17)	0.3507 (3)	0.0754 (9)	
H10A	0.3647	0.7562	0.3871	0.113*	
H10B	0.2486	0.7701	0.2931	0.113*	
H10C	0.2874	0.8174	0.3994	0.113*	
C11	0.19289 (18)	0.90745 (15)	0.01213 (19)	0.0451 (6)	
C12	0.14933 (18)	0.98317 (15)	-0.02230 (19)	0.0469 (6)	
C13	0.1192 (2)	1.00754 (18)	-0.1260 (2)	0.0583 (7)	
H13	0.0903	1.0576	-0.1475	0.070*	
C14	0.1313 (2)	0.9596 (2)	-0.1968 (2)	0.0673 (8)	
H14	0.1114	0.9768	-0.2655	0.081*	
C15	0.1737 (2)	0.88476 (19)	-0.1645 (2)	0.0624 (8)	
H15	0.1816	0.8515	-0.2124	0.075*	
C16	0.2044 (2)	0.85881 (16)	-0.0627 (2)	0.0541 (7)	
C17	0.3562 (3)	0.7792 (2)	0.0374 (3)	0.0882 (11)	
H17A	0.3757	0.8204	0.0905	0.132*	

## supplementary materials

---

H17B	0.3972	0.7870	-0.0010	0.132*	
C18	0.3805 (4)	0.7025 (3)	0.0884 (4)	0.1320 (17)	
H18A	0.3397	0.6952	0.1264	0.198*	
H18B	0.4561	0.6993	0.1372	0.198*	
H18C	0.3617	0.6621	0.0355	0.198*	
C19	0.12739 (19)	1.03513 (14)	0.0451 (2)	0.0484 (6)	
H19	0.0843	1.0787	0.0125	0.058*	
C20	0.1228 (2)	1.08673 (15)	0.1956 (2)	0.0552 (7)	
H20A	0.1819	1.1201	0.2407	0.083*	
H20B	0.0950	1.0604	0.2374	0.083*	
H20C	0.0669	1.1181	0.1420	0.083*	
C21	0.1177 (2)	0.90195 (14)	0.32031 (19)	0.0456 (6)	
C22	0.1746 (2)	0.95016 (15)	0.4104 (2)	0.0514 (7)	
C23	0.1439 (3)	0.95312 (17)	0.4916 (2)	0.0644 (8)	
H23	0.1814	0.9858	0.5500	0.077*	
C24	0.0613 (3)	0.90943 (19)	0.4861 (2)	0.0690 (8)	
H24	0.0411	0.9128	0.5396	0.083*	
C25	0.0061 (2)	0.85931 (16)	0.4001 (2)	0.0616 (7)	
H25	-0.0501	0.8286	0.3971	0.074*	
C26	0.0336 (2)	0.85452 (15)	0.3196 (2)	0.0504 (6)	
C27	-0.0963 (3)	0.75265 (17)	0.2315 (2)	0.0699 (8)	
H27A	-0.1573	0.7824	0.2271	0.105*	
H27B	-0.0671	0.7213	0.2960	0.105*	
C28	-0.1311 (3)	0.7007 (2)	0.1376 (3)	0.0891 (11)	
H28A	-0.1621	0.7320	0.0741	0.134*	
H28B	-0.1838	0.6639	0.1360	0.134*	
H28C	-0.0699	0.6724	0.1418	0.134*	
C29	0.2648 (2)	0.99735 (16)	0.4233 (2)	0.0556 (7)	
H29	0.2972	1.0275	0.4849	0.067*	
C30	0.3961 (2)	1.05901 (18)	0.3891 (3)	0.0714 (9)	
H30A	0.4084	1.0867	0.4526	0.107*	
H30B	0.3782	1.0961	0.3323	0.107*	
H30C	0.4602	1.0307	0.4012	0.107*	
N1	0.35163 (18)	0.85684 (12)	0.30894 (16)	0.0514 (5)	
N2	0.16059 (15)	1.02795 (11)	0.14483 (16)	0.0435 (5)	
N3	0.30595 (16)	1.00281 (12)	0.35992 (17)	0.0505 (5)	
O1	0.36036 (12)	0.99193 (9)	0.19967 (13)	0.0493 (4)	
O2	0.48376 (15)	1.08425 (12)	0.14429 (17)	0.0684 (6)	
O3	0.21754 (13)	0.87875 (9)	0.10683 (12)	0.0486 (4)	
O4	0.24247 (16)	0.78279 (11)	-0.03486 (15)	0.0663 (5)	
O5	0.13858 (13)	0.89719 (9)	0.23996 (13)	0.0475 (4)	
O6	-0.01521 (15)	0.80540 (10)	0.23346 (14)	0.0584 (5)	
O1W	0.0519 (2)	0.76823 (16)	0.0638 (2)	0.1173 (10)	
H1WA	0.1112	0.7907	0.0777	0.176*	
H1WB	0.0468	0.7664	0.1214	0.176*	
C8	0.3302 (13)	1.1506 (9)	0.0252 (13)	0.110 (3)	0.412 (15)
H8A	0.2703	1.1327	0.0347	0.165*	0.412 (15)
H8B	0.3041	1.1727	-0.0446	0.165*	0.412 (15)
H8C	0.3693	1.1901	0.0775	0.165*	0.412 (15)

C8'	0.3642 (9)	1.1540 (6)	-0.0134 (9)	0.110 (3)	0.588 (15)
H8'A	0.4168	1.1790	-0.0287	0.165*	0.588 (15)
H8'B	0.3487	1.1874	0.0323	0.165*	0.588 (15)
H8'C	0.2993	1.1451	-0.0784	0.165*	0.588 (15)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Co1	0.04589 (19)	0.03229 (19)	0.03721 (19)	0.00247 (14)	0.00959 (14)	-0.00092 (14)
C1	0.0427 (14)	0.0450 (15)	0.0415 (14)	0.0012 (11)	0.0063 (11)	-0.0116 (11)
C2	0.0462 (14)	0.0468 (16)	0.0471 (15)	0.0083 (12)	0.0045 (12)	-0.0074 (13)
C3	0.0526 (17)	0.0628 (19)	0.065 (2)	0.0147 (15)	0.0036 (15)	-0.0126 (16)
C4	0.0423 (16)	0.089 (3)	0.088 (2)	0.0107 (17)	0.0131 (17)	-0.022 (2)
C5	0.0472 (16)	0.082 (2)	0.082 (2)	-0.0067 (16)	0.0221 (15)	-0.0183 (19)
C6	0.0468 (15)	0.0523 (17)	0.0553 (17)	-0.0053 (12)	0.0123 (13)	-0.0132 (13)
C7	0.091 (3)	0.080 (2)	0.079 (3)	-0.006 (2)	0.010 (2)	0.005 (2)
C9	0.0626 (18)	0.0465 (16)	0.0420 (15)	0.0183 (14)	0.0046 (13)	-0.0014 (12)
C10	0.101 (2)	0.0520 (18)	0.077 (2)	0.0271 (17)	0.0448 (19)	0.0278 (16)
C11	0.0425 (13)	0.0451 (14)	0.0374 (14)	-0.0065 (11)	0.0101 (11)	-0.0028 (12)
C12	0.0384 (12)	0.0487 (15)	0.0412 (14)	-0.0022 (11)	0.0082 (11)	0.0031 (12)
C13	0.0508 (15)	0.0639 (18)	0.0447 (16)	-0.0012 (13)	0.0095 (13)	0.0111 (14)
C14	0.0594 (17)	0.089 (2)	0.0431 (16)	-0.0056 (16)	0.0151 (14)	0.0104 (16)
C15	0.0616 (17)	0.078 (2)	0.0435 (16)	-0.0102 (15)	0.0213 (14)	-0.0110 (15)
C16	0.0540 (15)	0.0520 (17)	0.0472 (16)	-0.0055 (12)	0.0161 (13)	-0.0054 (13)
C17	0.097 (3)	0.070 (2)	0.082 (2)	0.0242 (19)	0.029 (2)	-0.0117 (19)
C18	0.146 (4)	0.102 (3)	0.136 (4)	0.039 (3)	0.055 (3)	0.041 (3)
C19	0.0393 (13)	0.0409 (14)	0.0512 (17)	0.0029 (10)	0.0097 (12)	0.0071 (12)
C20	0.0560 (15)	0.0399 (14)	0.0639 (18)	0.0072 (12)	0.0232 (14)	-0.0006 (13)
C21	0.0536 (14)	0.0365 (14)	0.0390 (14)	0.0087 (11)	0.0151 (12)	0.0041 (11)
C22	0.0596 (16)	0.0447 (15)	0.0424 (15)	0.0068 (12)	0.0177 (13)	-0.0027 (12)
C23	0.080 (2)	0.0571 (18)	0.0532 (18)	0.0083 (15)	0.0281 (16)	-0.0088 (14)
C24	0.090 (2)	0.067 (2)	0.0576 (19)	0.0080 (18)	0.0416 (17)	-0.0022 (16)
C25	0.0729 (19)	0.0503 (17)	0.0662 (19)	0.0057 (14)	0.0364 (16)	0.0074 (15)
C26	0.0601 (16)	0.0382 (14)	0.0457 (15)	0.0055 (12)	0.0186 (13)	0.0061 (12)
C27	0.0747 (19)	0.0583 (19)	0.070 (2)	-0.0154 (15)	0.0278 (16)	0.0028 (16)
C28	0.095 (3)	0.076 (2)	0.083 (2)	-0.0341 (19)	0.031 (2)	-0.0142 (19)
C29	0.0616 (17)	0.0471 (16)	0.0433 (16)	0.0008 (13)	0.0122 (14)	-0.0111 (12)
C30	0.0659 (18)	0.069 (2)	0.067 (2)	-0.0190 (15)	0.0213 (15)	-0.0273 (16)
N1	0.0646 (14)	0.0390 (12)	0.0384 (12)	0.0099 (10)	0.0137 (10)	0.0026 (9)
N2	0.0417 (11)	0.0325 (11)	0.0474 (13)	-0.0011 (8)	0.0132 (10)	0.0001 (9)
N3	0.0500 (12)	0.0426 (12)	0.0448 (12)	-0.0006 (10)	0.0102 (10)	-0.0085 (10)
O1	0.0423 (9)	0.0417 (10)	0.0525 (10)	0.0045 (7)	0.0126 (8)	0.0039 (8)
O2	0.0624 (12)	0.0585 (12)	0.0731 (14)	-0.0073 (10)	0.0224 (11)	-0.0016 (11)
O3	0.0635 (11)	0.0344 (9)	0.0394 (10)	0.0011 (8)	0.0170 (8)	-0.0007 (7)
O4	0.0811 (14)	0.0505 (12)	0.0580 (12)	-0.0003 (10)	0.0247 (11)	-0.0132 (9)
O5	0.0579 (10)	0.0390 (10)	0.0387 (9)	-0.0051 (8)	0.0167 (8)	-0.0030 (8)
O6	0.0678 (12)	0.0476 (11)	0.0572 (11)	-0.0114 (9)	0.0270 (10)	-0.0023 (9)
O1W	0.137 (2)	0.126 (2)	0.1031 (19)	-0.0782 (19)	0.0675 (18)	-0.0525 (17)

## supplementary materials

---

C8	0.108 (5)	0.083 (3)	0.097 (5)	0.009 (4)	0.013 (3)	0.016 (4)
C8'	0.108 (5)	0.083 (3)	0.097 (5)	0.009 (4)	0.013 (3)	0.016 (4)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

Co1—O1	1.8792 (18)	C17—H17B	0.9700
Co1—O5	1.9044 (18)	C18—H18A	0.9600
Co1—O3	1.9061 (17)	C18—H18B	0.9600
Co1—N3	1.940 (2)	C18—H18C	0.9600
Co1—N1	1.947 (2)	C19—N2	1.276 (3)
Co1—N2	1.954 (2)	C19—H19	0.9300
C1—O1	1.305 (3)	C20—N2	1.468 (3)
C1—C2	1.416 (4)	C20—H20A	0.9600
C1—C6	1.416 (4)	C20—H20B	0.9600
C2—C3	1.411 (4)	C20—H20C	0.9600
C2—C9	1.423 (4)	C21—O5	1.309 (3)
C3—C4	1.352 (5)	C21—C22	1.412 (3)
C3—H3	0.9300	C21—C26	1.424 (4)
C4—C5	1.376 (5)	C22—C23	1.411 (4)
C4—H4	0.9300	C22—C29	1.434 (4)
C5—C6	1.374 (4)	C23—C24	1.347 (4)
C5—H5	0.9300	C23—H23	0.9300
C6—O2	1.370 (3)	C24—C25	1.391 (4)
C7—O2	1.391 (4)	C24—H24	0.9300
C7—C8	1.546 (16)	C25—C26	1.371 (4)
C7—C8'	1.427 (10)	C25—H25	0.9300
C7—H7A	0.9700	C26—O6	1.372 (3)
C7—H7B	0.9700	C27—O6	1.437 (3)
C9—N1	1.286 (3)	C27—C28	1.481 (4)
C9—H9	0.9300	C27—H27A	0.9700
C10—N1	1.463 (4)	C27—H27B	0.9700
C10—H10A	0.9600	C28—H28A	0.9600
C10—H10B	0.9600	C28—H28B	0.9600
C10—H10C	0.9600	C28—H28C	0.9600
C11—O3	1.318 (3)	C29—N3	1.281 (3)
C11—C12	1.409 (3)	C29—H29	0.9300
C11—C16	1.415 (4)	C30—N3	1.480 (3)
C12—C13	1.395 (4)	C30—H30A	0.9600
C12—C19	1.438 (4)	C30—H30B	0.9600
C13—C14	1.365 (4)	C30—H30C	0.9600
C13—H13	0.9300	O1W—H1WA	0.8507
C14—C15	1.386 (4)	O1W—H1WB	0.8550
C14—H14	0.9300	C8—H8A	0.9600
C15—C16	1.377 (4)	C8—H8B	0.9600
C15—H15	0.9300	C8—H8C	0.9600
C16—O4	1.381 (3)	C8'—H8'A	0.9600
C17—O4	1.439 (4)	C8'—H8'B	0.9600
C17—C18	1.452 (5)	C8'—H8'C	0.9600
C17—H17A	0.9700		

O1—Co1—O5	172.32 (7)	H18A—C18—H18C	109.5
O1—Co1—O3	88.46 (8)	H18B—C18—H18C	109.5
O5—Co1—O3	86.96 (7)	N2—C19—C12	127.1 (2)
O1—Co1—N3	91.17 (9)	N2—C19—H19	116.5
O5—Co1—N3	93.87 (9)	C12—C19—H19	116.5
O3—Co1—N3	175.49 (8)	N2—C20—H20A	109.5
O1—Co1—N1	94.28 (9)	N2—C20—H20B	109.5
O5—Co1—N1	91.54 (9)	H20A—C20—H20B	109.5
O3—Co1—N1	85.92 (8)	N2—C20—H20C	109.5
N3—Co1—N1	89.63 (9)	H20A—C20—H20C	109.5
O1—Co1—N2	85.59 (8)	H20B—C20—H20C	109.5
O5—Co1—N2	88.43 (8)	O5—C21—C22	124.5 (2)
O3—Co1—N2	92.24 (8)	O5—C21—C26	118.8 (2)
N3—Co1—N2	92.21 (9)	C22—C21—C26	116.7 (3)
N1—Co1—N2	178.15 (9)	C23—C22—C21	120.3 (3)
O1—C1—C2	124.5 (3)	C23—C22—C29	117.6 (3)
O1—C1—C6	118.5 (2)	C21—C22—C29	122.1 (3)
C2—C1—C6	117.0 (2)	C24—C23—C22	121.1 (3)
C3—C2—C1	119.8 (3)	C24—C23—H23	119.5
C3—C2—C9	118.3 (3)	C22—C23—H23	119.5
C1—C2—C9	121.9 (2)	C23—C24—C25	119.8 (3)
C4—C3—C2	121.4 (3)	C23—C24—H24	120.1
C4—C3—H3	119.3	C25—C24—H24	120.1
C2—C3—H3	119.3	C26—C25—C24	121.0 (3)
C3—C4—C5	119.4 (3)	C26—C25—H25	119.5
C3—C4—H4	120.3	C24—C25—H25	119.5
C5—C4—H4	120.3	C25—C26—O6	124.4 (3)
C6—C5—C4	121.6 (3)	C25—C26—C21	121.0 (3)
C6—C5—H5	119.2	O6—C26—C21	114.6 (2)
C4—C5—H5	119.2	O6—C27—C28	108.2 (3)
O2—C6—C5	119.9 (3)	O6—C27—H27A	110.1
O2—C6—C1	119.2 (2)	C28—C27—H27A	110.1
C5—C6—C1	120.7 (3)	O6—C27—H27B	110.1
O2—C7—C8	104.0 (6)	C28—C27—H27B	110.1
O2—C7—C8'	116.3 (5)	H27A—C27—H27B	108.4
O2—C7—H7A	108.9	C27—C28—H28A	109.5
C8—C7—H7A	95.8	C27—C28—H28B	109.5
C8'—C7—H7A	117.5	H28A—C28—H28B	109.5
O2—C7—H7B	108.9	C27—C28—H28C	109.5
C8—C7—H7B	129.5	H28A—C28—H28C	109.5
C8'—C7—H7B	96.0	H28B—C28—H28C	109.5
H7A—C7—H7B	107.9	N3—C29—C22	127.6 (2)
N1—C9—C2	128.0 (2)	N3—C29—H29	116.2
N1—C9—H9	116.0	C22—C29—H29	116.2
C2—C9—H9	116.0	N3—C30—H30A	109.5
N1—C10—H10A	109.5	N3—C30—H30B	109.5
N1—C10—H10B	109.5	H30A—C30—H30B	109.5
H10A—C10—H10B	109.5	N3—C30—H30C	109.5
N1—C10—H10C	109.5	H30A—C30—H30C	109.5

## supplementary materials

---

H10A—C10—H10C	109.5	H30B—C30—H30C	109.5
H10B—C10—H10C	109.5	C9—N1—C10	117.4 (2)
O3—C11—C12	123.8 (2)	C9—N1—C <sub>0</sub> 1	123.5 (2)
O3—C11—C16	119.1 (2)	C10—N1—C <sub>0</sub> 1	118.94 (19)
C12—C11—C16	117.0 (2)	C19—N2—C20	117.0 (2)
C13—C12—C11	120.4 (3)	C19—N2—C <sub>0</sub> 1	122.78 (18)
C13—C12—C19	118.1 (2)	C20—N2—C <sub>0</sub> 1	120.17 (17)
C11—C12—C19	121.4 (2)	C29—N3—C30	117.1 (2)
C14—C13—C12	121.6 (3)	C29—N3—C <sub>0</sub> 1	124.30 (18)
C14—C13—H13	119.2	C30—N3—C <sub>0</sub> 1	118.6 (2)
C12—C13—H13	119.2	C1—O1—C <sub>0</sub> 1	127.84 (17)
C13—C14—C15	118.9 (3)	C6—O2—C7	116.3 (2)
C13—C14—H14	120.6	C11—O3—C <sub>0</sub> 1	123.20 (15)
C15—C14—H14	120.6	C16—O4—C17	113.3 (2)
C16—C15—C14	121.1 (3)	C21—O5—C <sub>0</sub> 1	127.20 (15)
C16—C15—H15	119.4	C26—O6—C27	117.1 (2)
C14—C15—H15	119.4	H1WA—O1W—H1WB	107.7
C15—C16—O4	119.3 (3)	C7—C8—H8A	109.5
C15—C16—C11	121.0 (3)	C7—C8—H8B	109.5
O4—C16—C11	119.6 (2)	H8A—C8—H8B	109.5
O4—C17—C18	108.6 (3)	C7—C8—H8C	109.5
O4—C17—H17A	110.0	H8A—C8—H8C	109.5
C18—C17—H17A	110.0	H8B—C8—H8C	109.5
O4—C17—H17B	110.0	C7—C8'—H8'A	109.5
C18—C17—H17B	110.0	C7—C8'—H8'B	109.5
H17A—C17—H17B	108.4	H8'A—C8'—H8'B	109.5
C17—C18—H18A	109.5	C7—C8'—H8'C	109.5
C17—C18—H18B	109.5	H8'A—C8'—H8'C	109.5
H18A—C18—H18B	109.5	H8'B—C8'—H8'C	109.5
C17—C18—H18C	109.5		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1W—H1WA…O3	0.85	2.01	2.821 (3)	159
O1W—H1WB…O6	0.85	2.24	3.038 (3)	155

Fig. 1

