

Bis{2-[bis(2-hydroxyethyl)amino]acetato- $\kappa^3 O,N,O'$ }zinc(II)

Kong Mun Lo and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 22 October 2010; accepted 23 October 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$;
 R factor = 0.021; wR factor = 0.062; data-to-parameter ratio = 15.6.

In the crystal structure of the zinc(II) complex of bicine, $[\text{Zn}(\text{C}_6\text{H}_{12}\text{NO}_4)_2]$, the deprotonated amino acid O,N,O' -chelates to the metal atom through a carboxylate O atom, a hydroxy O atom and the N atom, the three atoms occupying *fac* positions of the distorted octahedron surrounding the metal atom. The metal atom lies on a center of inversion. The uncoordinated carboxylate O atom is hydrogen bonded to the hydroxy groups of adjacent molecules, these two hydrogen bonds leading to the formation of a three-dimensional network.

Related literature

For the isostructural cobalt(II) analog, see: Zhao & Liu (2010).



Experimental

Crystal data

$[\text{Zn}(\text{C}_6\text{H}_{12}\text{NO}_4)_2]$	$V = 772.28 (9)\text{ \AA}^3$
$M_r = 389.70$	$Z = 2$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 9.7863 (7)\text{ \AA}$	$\mu = 1.64\text{ mm}^{-1}$
$b = 11.3715 (8)\text{ \AA}$	$T = 100\text{ K}$
$c = 7.3462 (5)\text{ \AA}$	$0.25 \times 0.20 \times 0.15\text{ mm}$
$\beta = 109.1495 (8)^\circ$	

Data collection

Bruker SMART APEX	7174 measured reflections
diffractometer	1773 independent reflections
Absorption correction: multi-scan	1634 reflections with $I > 2\sigma(I)$
(SADABS; Sheldrick, 1996)	$R_{\text{int}} = 0.026$
	$T_{\text{min}} = 0.685, T_{\text{max}} = 0.792$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$	H atoms treated by a mixture of
$wR(F^2) = 0.062$	independent and constrained
$S = 1.12$	refinement
1773 reflections	$\Delta\rho_{\text{max}} = 0.43\text{ e \AA}^{-3}$
114 parameters	$\Delta\rho_{\text{min}} = -0.35\text{ e \AA}^{-3}$
2 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O}3-\text{H}3 \cdots \text{O}2^{\text{i}}$	0.84 (1)	1.85 (1)	2.651 (2)	161 (2)
$\text{O}4-\text{H}4 \cdots \text{O}2^{\text{ii}}$	0.83 (1)	1.89 (1)	2.715 (2)	178 (3)

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank the University of Malaya (grant No. RG020/09AFR) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5062).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zhao, J.-P. & Liu, F.-C. (2010). *Acta Cryst. E* **66**, m848.

supplementary materials

Acta Cryst. (2010). E66, m1485 [doi:10.1107/S1600536810043217]

Bis{2-[bis(2-hydroxyethyl)amino]acetato- κ^3O,N,O' }zinc(II)

K. M. Lo and S. W. Ng

Comment

In the crystal structure of the cobalt(II) derivative of bicine, the deprotonated aminoacid N,O,O' -chelates to the metal atom through the nitrogen, carboxyl oxygen and hydroxyl oxygen atoms, the three atoms occupying *fac* positions of the octahedron (Zhao & Liu, 2010). The present zinc analog (Scheme I, Fig. 1) is isostructural. The double-bond carboxyl oxygen atom is hydrogen-bond acceptor to the coordinated as well as the free hydroxyl unit of adjacent molecules, these two hydrogen bonds leading to the formation of a three-dimensional network (Table 1).

Experimental

N,N-Bis(2-hydroxyethyl)glycine (0.94 g, 5.7 mmol) and zinc carbonate (0.36 g, 2.8 mmol) were heated in a 1:1 water:DMSO mixture (100 ml) for 1 h. The solution was filtered; colorless crystals were obtained upon slow evaporation of the filtrate.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.99 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to $1.2U(C)$.

The hydroxy H-atoms were located in a difference Fourier map, and were refined with the O—H distance restrained to 0.84 ± 0.01 Å; its temperature factor was refined.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $Zn(C_6H_{12}NO_4)_2$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

Bis{2-[bis(2-hydroxyethyl)amino]acetato- κ^3O,N,O' }zinc(II)

Crystal data

$[Zn(C_6H_{12}NO_4)_2]$	$F(000) = 408$
$M_r = 389.70$	$D_x = 1.676 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Hall symbol: -P 2ybc	Cell parameters from 4608 reflections
$a = 9.7863 (7)$ Å	$\theta = 2.2\text{--}28.3^\circ$

supplementary materials

$b = 11.3715 (8) \text{ \AA}$	$\mu = 1.64 \text{ mm}^{-1}$
$c = 7.3462 (5) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 109.1495 (8)^\circ$	Block, colorless
$V = 772.28 (9) \text{ \AA}^3$	$0.25 \times 0.20 \times 0.15 \text{ mm}$
$Z = 2$	

Data collection

Bruker SMART APEX diffractometer	1773 independent reflections
Radiation source: fine-focus sealed tube graphite	1634 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.026$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.2^\circ$
$T_{\text{min}} = 0.685, T_{\text{max}} = 0.792$	$h = -12 \rightarrow 12$
7174 measured reflections	$k = -14 \rightarrow 14$
	$l = -9 \rightarrow 9$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.021$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.062$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.12$	$w = 1/[\sigma^2(F_o^2) + (0.0283P)^2 + 0.4612P]$
1773 reflections	where $P = (F_o^2 + 2F_c^2)/3$
114 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
2 restraints	$\Delta\rho_{\text{max}} = 0.43 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.35 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	0.5000	0.5000	0.00889 (9)
O1	0.45689 (11)	0.33271 (9)	0.56937 (15)	0.0132 (2)
O2	0.28013 (11)	0.21247 (9)	0.57458 (17)	0.0167 (2)
O3	0.51490 (11)	0.55292 (9)	0.78390 (15)	0.0129 (2)
H3	0.564 (2)	0.6126 (14)	0.831 (3)	0.030 (6)*
O4	0.00548 (11)	0.66106 (11)	0.01696 (15)	0.0160 (2)
H4	-0.0808 (12)	0.678 (2)	-0.009 (3)	0.040 (7)*
N1	0.27253 (13)	0.52389 (11)	0.44766 (18)	0.0095 (2)
C1	0.32624 (16)	0.31061 (13)	0.5430 (2)	0.0116 (3)
C2	0.21180 (15)	0.40649 (12)	0.4649 (2)	0.0114 (3)
H2A	0.1466	0.3821	0.3363	0.014*
H2B	0.1528	0.4130	0.5511	0.014*

C3	0.26823 (16)	0.60594 (13)	0.6035 (2)	0.0128 (3)
H3A	0.1690	0.6082	0.6106	0.015*
H3B	0.2935	0.6862	0.5732	0.015*
C4	0.37309 (16)	0.56760 (14)	0.7966 (2)	0.0141 (3)
H4A	0.3757	0.6275	0.8955	0.017*
H4B	0.3400	0.4925	0.8360	0.017*
C5	0.20539 (15)	0.57679 (13)	0.2541 (2)	0.0114 (3)
H5A	0.2129	0.5197	0.1561	0.014*
H5B	0.2620	0.6472	0.2442	0.014*
C6	0.04786 (16)	0.61242 (13)	0.2055 (2)	0.0131 (3)
H6A	-0.0125	0.5431	0.2091	0.016*
H6B	0.0370	0.6713	0.2990	0.016*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.00708 (13)	0.00885 (13)	0.01018 (13)	-0.00067 (8)	0.00204 (9)	0.00051 (8)
O1	0.0092 (5)	0.0119 (5)	0.0173 (5)	-0.0003 (4)	0.0026 (4)	0.0023 (4)
O2	0.0106 (5)	0.0123 (5)	0.0236 (6)	-0.0021 (4)	0.0008 (4)	0.0057 (4)
O3	0.0102 (5)	0.0136 (5)	0.0133 (5)	-0.0013 (4)	0.0019 (4)	-0.0020 (4)
O4	0.0111 (5)	0.0223 (6)	0.0127 (5)	0.0045 (4)	0.0014 (4)	0.0056 (4)
N1	0.0094 (6)	0.0089 (5)	0.0098 (6)	-0.0002 (4)	0.0026 (5)	0.0004 (4)
C1	0.0120 (7)	0.0118 (6)	0.0100 (6)	0.0003 (5)	0.0021 (5)	0.0005 (5)
C2	0.0091 (7)	0.0114 (6)	0.0131 (7)	-0.0011 (5)	0.0029 (5)	0.0013 (5)
C3	0.0125 (7)	0.0127 (6)	0.0129 (7)	0.0015 (5)	0.0037 (6)	-0.0013 (5)
C4	0.0124 (7)	0.0184 (7)	0.0117 (7)	0.0001 (5)	0.0042 (6)	-0.0013 (6)
C5	0.0092 (7)	0.0133 (7)	0.0109 (7)	0.0007 (5)	0.0021 (5)	0.0016 (5)
C6	0.0106 (7)	0.0151 (7)	0.0118 (7)	0.0022 (5)	0.0012 (5)	0.0016 (5)

Geometric parameters (\AA , $^\circ$)

Zn1—O1	2.0488 (10)	N1—C3	1.4882 (18)
Zn1—O1 ⁱ	2.0488 (10)	C1—C2	1.533 (2)
Zn1—O3 ⁱ	2.1292 (11)	C2—H2A	0.9900
Zn1—O3	2.1292 (11)	C2—H2B	0.9900
Zn1—N1 ⁱ	2.1484 (13)	C3—C4	1.516 (2)
Zn1—N1	2.1484 (13)	C3—H3A	0.9900
O1—C1	1.2542 (18)	C3—H3B	0.9900
O2—C1	1.2537 (18)	C4—H4A	0.9900
O3—C4	1.4308 (18)	C4—H4B	0.9900
O3—H3	0.84 (1)	C5—C6	1.519 (2)
O4—C6	1.4212 (17)	C5—H5A	0.9900
O4—H4	0.83 (1)	C5—H5B	0.9900
N1—C2	1.4836 (18)	C6—H6A	0.9900
N1—C5	1.4839 (18)	C6—H6B	0.9900
O1—Zn1—O1 ⁱ	180.0	N1—C2—H2A	108.7
O1—Zn1—O3 ⁱ	91.59 (4)	C1—C2—H2A	108.7

supplementary materials

O1 ⁱ —Zn1—O3 ⁱ	88.41 (4)	N1—C2—H2B	108.7
O1—Zn1—O3	88.41 (4)	C1—C2—H2B	108.7
O1 ⁱ —Zn1—O3	91.59 (4)	H2A—C2—H2B	107.6
O3 ⁱ —Zn1—O3	180.0	N1—C3—C4	110.98 (12)
O1—Zn1—N1 ⁱ	97.12 (4)	N1—C3—H3A	109.4
O1 ⁱ —Zn1—N1 ⁱ	82.88 (4)	C4—C3—H3A	109.4
O3 ⁱ —Zn1—N1 ⁱ	82.74 (4)	N1—C3—H3B	109.4
O3—Zn1—N1 ⁱ	97.26 (4)	C4—C3—H3B	109.4
O1—Zn1—N1	82.88 (4)	H3A—C3—H3B	108.0
O1 ⁱ —Zn1—N1	97.12 (4)	O3—C4—C3	110.24 (12)
O3 ⁱ —Zn1—N1	97.26 (4)	O3—C4—H4A	109.6
O3—Zn1—N1	82.74 (4)	C3—C4—H4A	109.6
N1 ⁱ —Zn1—N1	180.0	O3—C4—H4B	109.6
C1—O1—Zn1	115.48 (9)	C3—C4—H4B	109.6
C4—O3—Zn1	109.91 (8)	H4A—C4—H4B	108.1
C4—O3—H3	108.3 (16)	N1—C5—C6	115.44 (12)
Zn1—O3—H3	118.6 (16)	N1—C5—H5A	108.4
C6—O4—H4	105.4 (17)	C6—C5—H5A	108.4
C2—N1—C5	112.49 (11)	N1—C5—H5B	108.4
C2—N1—C3	112.67 (11)	C6—C5—H5B	108.4
C5—N1—C3	111.57 (11)	H5A—C5—H5B	107.5
C2—N1—Zn1	106.95 (8)	O4—C6—C5	106.39 (12)
C5—N1—Zn1	109.26 (9)	O4—C6—H6A	110.5
C3—N1—Zn1	103.34 (9)	C5—C6—H6A	110.5
O2—C1—O1	124.08 (13)	O4—C6—H6B	110.5
O2—C1—C2	116.06 (13)	C5—C6—H6B	110.5
O1—C1—C2	119.86 (13)	H6A—C6—H6B	108.6
N1—C2—C1	114.08 (12)		
O3 ⁱ —Zn1—O1—C1	−92.62 (10)	O3 ⁱ —Zn1—N1—C3	−157.45 (8)
O3—Zn1—O1—C1	87.38 (10)	O3—Zn1—N1—C3	22.55 (8)
N1 ⁱ —Zn1—O1—C1	−175.50 (10)	Zn1—O1—C1—O2	178.78 (12)
N1—Zn1—O1—C1	4.50 (10)	Zn1—O1—C1—C2	−0.37 (17)
O1—Zn1—O3—C4	−78.86 (9)	C5—N1—C2—C1	128.95 (13)
O1 ⁱ —Zn1—O3—C4	101.14 (9)	C3—N1—C2—C1	−103.88 (14)
N1 ⁱ —Zn1—O3—C4	−175.83 (9)	Zn1—N1—C2—C1	8.99 (14)
N1—Zn1—O3—C4	4.17 (9)	O2—C1—C2—N1	174.38 (12)
O1—Zn1—N1—C2	−7.26 (9)	O1—C1—C2—N1	−6.4 (2)
O1 ⁱ —Zn1—N1—C2	172.74 (9)	C2—N1—C3—C4	68.96 (15)
O3 ⁱ —Zn1—N1—C2	83.44 (9)	C5—N1—C3—C4	−163.38 (12)
O3—Zn1—N1—C2	−96.56 (9)	Zn1—N1—C3—C4	−46.10 (13)
O1—Zn1—N1—C5	−129.27 (9)	Zn1—O3—C4—C3	−30.45 (14)
O1 ⁱ —Zn1—N1—C5	50.73 (9)	N1—C3—C4—O3	53.60 (16)
O3 ⁱ —Zn1—N1—C5	−38.57 (9)	C2—N1—C5—C6	68.31 (16)
O3—Zn1—N1—C5	141.43 (9)	C3—N1—C5—C6	−59.44 (16)
O1—Zn1—N1—C3	111.84 (9)	Zn1—N1—C5—C6	−173.08 (10)

O1ⁱ—Zn1—N1—C3 —68.16 (9) N1—C5—C6—O4 179.67 (11)
Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O3—H3 \cdots O2 ⁱⁱ	0.84 (1)	1.85 (1)	2.651 (2)	161 (2)
O4—H4 \cdots O2 ⁱⁱⁱ	0.83 (1)	1.89 (1)	2.715 (2)	178 (3)

Symmetry codes: (ii) $-x+1, y+1/2, -z+3/2$; (iii) $-x, y+1/2, -z+1/2$.

supplementary materials

Fig. 1

