

12 β ,14-Dihydroxy-3-oxo-5 β ,20(22)-cardenolide monohydrate

Chao-Jun He, Min Wang,* Hua Sun, Zeng-bing Liu and Yu-wan Fu

College of Biotechnology, Tianjin University of Science and Technology, Tianjin 300457, People's Republic of China

Correspondence e-mail: minw@tust.edu.cn


Received 2 August 2010; accepted 23 September 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.034; wR factor = 0.082; data-to-parameter ratio = 6.6.

The title compound, digoxigenone, $\text{C}_{23}\text{H}_{30}\text{O}_5 \cdot \text{H}_2\text{O}$, was biotransformed from digoxigenin. In the crystal, intermolecular $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds contribute to the formation of a three-dimensional supramolecular structure. The title compound has three fused six-membered rings (A, B, C) and two non-fused five-membered rings (D, E). As in other structures, compound nucleus has a *cis-trans-cis* conformation for the $A-B, B-C, C-D$ ring junctions with rings A, B and C exhibiting chair conformations.

Related literature

Digitoxin and digoxin, the typical clinically used forms (Kreis *et al.*, 1998), are the drugs of choice for the treatment of congestive heart failure, acting as selective inhibitors of the Na^+ , K^+ ATPase enzyme. For the biotransformation of digoxigenin into digoxigenin and digoxigenone by *Fusarium ciliatum* and into 1 β -hydroxydigoxigenin, 7- β -hydroxydigoxigenin, 8- β -hydroxydigoxigenin and digitoxigenone by *Cochliobolus lunatus*, see: Pádua *et al.* (2005, 2007).



Experimental

Crystal data

$\text{C}_{23}\text{H}_{30}\text{O}_5 \cdot \text{H}_2\text{O}$
 $M_r = 404.49$
 Triclinic, $P1$
 $a = 7.4017$ (15) Å
 $b = 7.7450$ (15) Å
 $c = 10.215$ (2) Å
 $\alpha = 99.51$ (3)°
 $\beta = 94.70$ (3)°
 $\gamma = 114.97$ (3)°
 $V = 516.0$ (2) Å³
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 293$ K
 $0.32 \times 0.26 \times 0.20$ mm

Data collection

Rigaku Saturn CCD area-detector diffractometer
 Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.971$, $T_{\max} = 0.982$
 5251 measured reflections
 1809 independent reflections
 1443 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.082$
 $S = 1.00$
 1809 reflections
 275 parameters
 6 restraints
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.15$ e Å⁻³
 $\Delta\rho_{\min} = -0.14$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O}2-\text{H}2 \cdots \text{O}6^i$	0.82	1.99	2.808 (3)	173
$\text{O}3-\text{H}3 \cdots \text{O}2^{ii}$	0.82	2.10	2.900 (3)	164
$\text{O}6-\text{H}1\text{W} \cdots \text{O}1^{iii}$	0.86 (1)	1.94 (1)	2.801 (3)	174 (4)
$\text{O}6-\text{H}2\text{W} \cdots \text{O}5^v$	0.86 (1)	1.91 (2)	2.741 (4)	162 (5)

Symmetry codes: (i) $x-1, y, z-1$; (ii) $x+1, y, z$; (iii) $x+1, y+1, z$; (iv) $x, y-1, z+1$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the Natural Science Foundation of Tianjin (No. 08JCZDJ15200)

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2696).

References

- Kreis, W., Hensel, A. & Stuhlemmer, U. (1998). *Planta Med.* **64**, 491–499.
 Pádua, R. M., Oliveira, A. B., Souza Filho, J. D., Takahashi, J. A., Silva, M. A. & Braga, F. C. (2007). *J. Braz. Chem. Soc.* **18**, 1303–1310.
 Pádua, R. M., Oliveira, A. B., Souza Filho, J. D., Vieira, G. J., Takahashi, J. A. & Braga, F. C. (2005). *J. Braz. Chem. Soc.* **16**, 614–619.
 Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, o2704 [doi:10.1107/S1600536810038031]

12 β ,14-Dihydroxy-3-oxo-5 β ,20(22)-cardenolide monohydrate

C.-J. He, M. Wang, H. Sun, Z. Liu and Y. Fu

Comment

Digitoxin and digoxin, the typical clinically used forms (Kreis *et al.*, 1998), are the drugs of choice for the treatment of congestive heart failure, acting as selective inhibitors of the Na⁺, K⁺ ATPase enzyme. Digitoxigenin has been biotransformed into digoxigenin and digoxigenone by *Fusarium ciliatum* and into 1 beta-hydroxydigitoxigenin, 7 beta-hydroxydigitoxigenin, 8 beta-hydroxydigitoxigenin, and digitoxigenone by *Cochliobolus lunatus* (Pádua *et al.*, 2005; Pádua *et al.*, 2007). In this paper, we studied the biotransformation of digoxigenin by *Arthrobacter simplex* and reported crystal structure of the compound digoxigenone.

The crystal structure is shown in Fig. 1. The crystal structure consists of an digoxigenone molecule and one water molecule. Compound digoxigenone has three fused six-membered rings (A/B/C) and two non-fused five-membered rings (D/E). As in other structures the A, B and C rings have a chair conformation. The 12-hydroxy is beta configuration with the torsion angles C9—C11—C12—O2 = -179.6 (2)°. The 14-hydroxy is beta configuration with the torsion angles C7—C8—C14—O3 = -64.4 (3)°. The orientation of the lactone ring is determined by the torsion angle C(13)—C(17)—C(20)—C(22) = -113.1 (3)°.

In the crystal packing (Fig. 2), X-ray analysis indicates that there were three types of intermolecular hydrogen bonds contributed to the formation of three-dimensional supramolecular structure: between solvent water molecule and carbonyl of adjacent digoxigenone molecule, hydroxyl of digoxigenone molecule and hydroxyl of adjacent digoxigenone molecule, and hydroxyl of digoxigenone molecule and solvent water molecule. The hydrogen bonding parameters are O(6)—H(6 A)···O(1)#3, 2.800 (3), 173.7; O(6)—H(6B)···O(5)#4, 2.742 (3), 160.3; O(2)—H(2)···O(6)#1 2.805 (2), 172.7; and O(3)—H(3)···O(2)#2, 2.901 (2), 163.7. (Symmetry code: #1 $x - 1, y, z - 1$; #2 $x + 1, y, z$; #3 $x + 1, y + 1, z$; #4 $x, y - 1, z + 1$).

Experimental

The *Arthrobacter simplex* TCCC 11037 (stored in our laboratory) was maintained at 277 K on a slant containing glucose 10.0 g/L, yeast extract 10.0 g L⁻¹ and agar 20.0 g L⁻¹ (pH = 7.2). Seed media consists of glucose 10.0 g/L, corn slurry 10.0 g/L, peptone 5.0 g/L and KH₂PO₄ 2.5 g/L (pH = 7.2).

The *Arthrobacter simplex* (ASP) cells were prepared in two consecutive cultivation steps (18 h for seed culture and 24 h for cell incubation, respectively) in shake flasks. The whole ASP culture using 5% (v/v) of seed culture was grown in 250 ml shake flasks containing 30 ml culture media on a rotary shaker (160 rpm) at 305 K. Digoxigenin dissolved in ethanol (50 g L⁻¹) and distributed among the flasks (0.34 g/L), and then the reaction was allowed to proceed for 7 days at 307 K. The biotransformation products were sequentially extracted with ethyl acetate (3 × 100 ml) in a separator funnel and the solvent was vacuum removed at 333 K, until a residue was produced. The crude extracts were purified by Si gel column using chloroform/methanol (25:1, v/v). The white power was diffused with petroleum ether/acetone (5:3, v/v) at room temperature. Colourless prism crystals were obtained.

Refinement

The C-bound H atoms were geometrically placed (C–H = 0.96 Å for methyl, 0.97 Å for methylene and 0.93 Å for C(sp²), respectively) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for others. The H atoms of hydroxyl were geometrically placed O–H = 0.82 Å and refined as riding with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. The water H atoms were located from a difference map and refined freely.

Figures




Fig. 1. Perspective view of the title compound (noncoordinated water molecule was omitted for clarity)




Fig. 2. View of the packing of the title compound

12β,14-Dihydroxy-3-oxo-5β,20 (22)-cardenolide monohydrate

Crystal data

$\text{C}_{23}\text{H}_{30}\text{O}_5 \cdot \text{H}_2\text{O}$

$M_r = 404.49$

Triclinic, *P*1

Hall symbol: P 1

$a = 7.4017 (15) \text{ \AA}$

$b = 7.7450 (15) \text{ \AA}$

$c = 10.215 (2) \text{ \AA}$

$\alpha = 99.51 (3)^\circ$

$\beta = 94.70 (3)^\circ$

$\gamma = 114.97 (3)^\circ$

$V = 516.0 (2) \text{ \AA}^3$

$Z = 1$

$F(000) = 218$

$D_x = 1.302 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1642 reflections

$\theta = 2.1\text{--}27.8^\circ$

$\mu = 0.09 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Prism, colourless

$0.32 \times 0.26 \times 0.20 \text{ mm}$

Data collection

Rigaku Saturn CCD area-detector diffractometer

Radiation source: rotating anode confocal

Detector resolution: $7.31 \text{ pixels mm}^{-1}$

ω and ϕ scans

Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2005)

$T_{\text{min}} = 0.971$, $T_{\text{max}} = 0.982$

1809 independent reflections

1443 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.1^\circ$

$h = -8 \rightarrow 8$

$k = -9 \rightarrow 8$

$l = -12 \rightarrow 12$

5251 measured reflections

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.082$	$w = 1/[\sigma^2(F_o^2) + (0.0504P)^2 + 0.010P]$
$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
1809 reflections	$(\Delta/\sigma)_{\max} < 0.001$
275 parameters	$\Delta\rho_{\max} = 0.14 \text{ e } \text{\AA}^{-3}$
6 restraints	$\Delta\rho_{\min} = -0.14 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
	Extinction coefficient: 0.43 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.0073 (4)	0.3043 (5)	0.5686 (3)	0.0474 (8)
H1A	-0.1000	0.3610	0.5551	0.057*
H1B	-0.0281	0.2103	0.4864	0.057*
C2	-0.0581 (5)	0.1979 (5)	0.6826 (3)	0.0566 (9)
H2A	-0.0574	0.2868	0.7618	0.068*
H2B	-0.1931	0.0907	0.6569	0.068*
C3	0.0884 (5)	0.1203 (5)	0.7161 (3)	0.0562 (9)
C4	0.2931 (5)	0.2275 (5)	0.6930 (3)	0.0548 (8)
H4	0.3868	0.1813	0.7137	0.066*
C5	0.3553 (4)	0.3895 (4)	0.6435 (3)	0.0394 (7)
C6	0.5751 (4)	0.5096 (5)	0.6423 (3)	0.0510 (8)
H6A	0.6496	0.4382	0.6646	0.061*
H6B	0.6252	0.6295	0.7107	0.061*
C7	0.6133 (4)	0.5598 (5)	0.5056 (3)	0.0426 (7)

supplementary materials

H7A	0.5823	0.4415	0.4395	0.051*
H7B	0.7552	0.6476	0.5118	0.051*
C8	0.4846 (4)	0.6561 (4)	0.4589 (3)	0.0302 (6)
H8	0.5236	0.7780	0.5253	0.036*
C9	0.2593 (3)	0.5255 (4)	0.4572 (3)	0.0299 (6)
H9	0.2219	0.4043	0.3906	0.036*
C10	0.2105 (4)	0.4666 (3)	0.5934 (2)	0.0321 (6)
C11	0.1312 (4)	0.6216 (4)	0.4064 (3)	0.0380 (7)
H11A	-0.0108	0.5345	0.4005	0.046*
H11B	0.1622	0.7411	0.4712	0.046*
C12	0.1679 (4)	0.6685 (4)	0.2704 (3)	0.0333 (6)
H12	0.1254	0.5461	0.2038	0.040*
C13	0.3939 (4)	0.8041 (3)	0.2682 (2)	0.0302 (6)
C14	0.5248 (4)	0.7098 (4)	0.3221 (3)	0.0311 (6)
C15	0.4892 (4)	0.5418 (4)	0.2060 (3)	0.0364 (6)
H15A	0.3658	0.4274	0.2068	0.044*
H15B	0.6013	0.5080	0.2116	0.044*
C16	0.4720 (4)	0.6179 (4)	0.0779 (3)	0.0441 (7)
H16A	0.3672	0.5153	0.0083	0.053*
H16B	0.5991	0.6614	0.0442	0.053*
C17	0.4186 (4)	0.7903 (4)	0.1164 (3)	0.0350 (6)
H17	0.2852	0.7522	0.0644	0.042*
C18	0.2350 (5)	0.6415 (5)	0.7033 (3)	0.0507 (8)
H18A	0.2182	0.6027	0.7878	0.076*
H18B	0.1346	0.6836	0.6789	0.076*
H18C	0.3674	0.7469	0.7116	0.076*
C19	0.4521 (4)	1.0101 (4)	0.3480 (3)	0.0418 (7)
H19A	0.5941	1.0896	0.3529	0.063*
H19B	0.4220	1.0049	0.4373	0.063*
H19C	0.3767	1.0654	0.3039	0.063*
C20	0.5633 (4)	0.9736 (4)	0.0803 (3)	0.0380 (6)
C21	0.7889 (4)	1.0689 (4)	0.1180 (4)	0.0567 (9)
H21A	0.8296	1.1136	0.2150	0.068*
H21B	0.8434	0.9787	0.0863	0.068*
C22	0.5139 (5)	1.0757 (4)	0.0035 (3)	0.0488 (8)
H22	0.3828	1.0478	-0.0331	0.059*
C23	0.6953 (5)	1.2364 (4)	-0.0143 (3)	0.0523 (8)
O1	0.0400 (5)	-0.0230 (4)	0.7666 (3)	0.0905 (10)
O2	0.0392 (3)	0.7578 (3)	0.2377 (2)	0.0497 (6)
H2	0.0251	0.7534	0.1566	0.075*
O3	0.7324 (2)	0.8562 (2)	0.33978 (19)	0.0392 (5)
H3	0.8017	0.8051	0.3095	0.059*
O4	0.8586 (3)	1.2327 (3)	0.0526 (2)	0.0610 (6)
O5	0.7166 (4)	1.3617 (4)	-0.0773 (3)	0.0797 (8)
O6	0.9639 (4)	0.7538 (4)	0.9630 (3)	0.0661 (7)
H1W	0.979 (7)	0.816 (5)	0.899 (3)	0.105 (15)*
H2W	0.901 (7)	0.6285 (16)	0.937 (4)	0.13 (2)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0383 (16)	0.0560 (19)	0.0500 (19)	0.0171 (14)	0.0123 (14)	0.0255 (16)
C2	0.0578 (19)	0.053 (2)	0.061 (2)	0.0180 (16)	0.0204 (16)	0.0302 (16)
C3	0.076 (2)	0.0477 (19)	0.049 (2)	0.0266 (17)	0.0187 (17)	0.0210 (16)
C4	0.070 (2)	0.063 (2)	0.056 (2)	0.0440 (18)	0.0204 (16)	0.0306 (17)
C5	0.0454 (16)	0.0485 (17)	0.0328 (16)	0.0267 (14)	0.0062 (12)	0.0149 (13)
C6	0.0421 (17)	0.075 (2)	0.0494 (19)	0.0327 (16)	0.0062 (14)	0.0296 (17)
C7	0.0332 (16)	0.0563 (18)	0.0503 (18)	0.0265 (14)	0.0103 (13)	0.0232 (15)
C8	0.0299 (13)	0.0342 (14)	0.0291 (14)	0.0165 (11)	0.0042 (10)	0.0076 (11)
C9	0.0297 (13)	0.0332 (14)	0.0311 (14)	0.0168 (11)	0.0066 (10)	0.0099 (11)
C10	0.0342 (15)	0.0356 (16)	0.0300 (15)	0.0179 (12)	0.0057 (11)	0.0089 (12)
C11	0.0336 (15)	0.0534 (17)	0.0390 (16)	0.0261 (13)	0.0104 (12)	0.0205 (14)
C12	0.0263 (13)	0.0449 (16)	0.0383 (16)	0.0213 (12)	0.0098 (11)	0.0166 (13)
C13	0.0309 (13)	0.0318 (14)	0.0306 (15)	0.0156 (11)	0.0074 (11)	0.0083 (11)
C14	0.0234 (13)	0.0334 (14)	0.0380 (15)	0.0127 (11)	0.0087 (11)	0.0096 (12)
C15	0.0371 (14)	0.0359 (15)	0.0399 (16)	0.0179 (12)	0.0138 (12)	0.0093 (13)
C16	0.0528 (17)	0.0400 (15)	0.0382 (17)	0.0183 (13)	0.0133 (13)	0.0083 (13)
C17	0.0322 (13)	0.0384 (15)	0.0311 (14)	0.0117 (12)	0.0056 (11)	0.0096 (12)
C18	0.075 (2)	0.0529 (19)	0.0376 (17)	0.0377 (18)	0.0189 (15)	0.0126 (15)
C19	0.0480 (16)	0.0357 (15)	0.0439 (16)	0.0210 (13)	0.0088 (13)	0.0076 (13)
C20	0.0429 (15)	0.0383 (15)	0.0328 (15)	0.0169 (12)	0.0113 (12)	0.0088 (12)
C21	0.0425 (18)	0.049 (2)	0.074 (2)	0.0101 (15)	0.0141 (15)	0.0295 (17)
C22	0.055 (2)	0.0499 (18)	0.0475 (18)	0.0235 (16)	0.0153 (14)	0.0210 (15)
C23	0.071 (2)	0.0427 (18)	0.0470 (19)	0.0238 (16)	0.0232 (17)	0.0179 (15)
O1	0.119 (2)	0.0708 (17)	0.107 (2)	0.0453 (16)	0.0475 (19)	0.0631 (18)
O2	0.0421 (11)	0.0820 (15)	0.0485 (12)	0.0417 (11)	0.0136 (10)	0.0331 (12)
O3	0.0267 (9)	0.0401 (11)	0.0502 (12)	0.0131 (8)	0.0086 (8)	0.0119 (9)
O4	0.0545 (14)	0.0495 (13)	0.0670 (16)	0.0068 (10)	0.0171 (11)	0.0236 (11)
O5	0.108 (2)	0.0596 (15)	0.083 (2)	0.0347 (15)	0.0367 (15)	0.0438 (14)
O6	0.0627 (15)	0.0775 (19)	0.0564 (15)	0.0232 (15)	0.0116 (12)	0.0316 (14)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.518 (4)	C13—C19	1.526 (4)
C1—C10	1.536 (4)	C13—C14	1.555 (3)
C1—H1A	0.9700	C13—C17	1.567 (3)
C1—H1B	0.9700	C14—O3	1.448 (3)
C2—C3	1.487 (4)	C14—C15	1.523 (4)
C2—H2A	0.9700	C15—C16	1.538 (4)
C2—H2B	0.9700	C15—H15A	0.9700
C3—O1	1.229 (4)	C15—H15B	0.9700
C3—C4	1.450 (5)	C16—C17	1.547 (4)
C4—C5	1.342 (4)	C16—H16A	0.9700
C4—H4	0.9300	C16—H16B	0.9700
C5—C6	1.497 (4)	C17—C20	1.501 (4)
C5—C10	1.523 (3)	C17—H17	0.9800

supplementary materials

C6—C7	1.527 (4)	C18—H18A	0.9600
C6—H6A	0.9700	C18—H18B	0.9600
C6—H6B	0.9700	C18—H18C	0.9600
C7—C8	1.528 (4)	C19—H19A	0.9600
C7—H7A	0.9700	C19—H19B	0.9600
C7—H7B	0.9700	C19—H19C	0.9600
C8—C14	1.539 (3)	C20—C22	1.333 (4)
C8—C9	1.540 (3)	C20—C21	1.497 (4)
C8—H8	0.9800	C21—O4	1.451 (3)
C9—C11	1.537 (3)	C21—H21A	0.9700
C9—C10	1.558 (3)	C21—H21B	0.9700
C9—H9	0.9800	C22—C23	1.445 (4)
C10—C18	1.544 (4)	C22—H22	0.9300
C11—C12	1.510 (4)	C23—O5	1.215 (4)
C11—H11A	0.9700	C23—O4	1.352 (4)
C11—H11B	0.9700	O2—H2	0.8200
C12—O2	1.442 (3)	O3—H3	0.8200
C12—C13	1.560 (3)	O6—H1W	0.861 (11)
C12—H12	0.9800	O6—H2W	0.862 (11)
C2—C1—C10	113.8 (2)	C13—C12—H12	108.7
C2—C1—H1A	108.8	C19—C13—C14	113.6 (2)
C10—C1—H1A	108.8	C19—C13—C12	110.0 (2)
C2—C1—H1B	108.8	C14—C13—C12	108.13 (19)
C10—C1—H1B	108.8	C19—C13—C17	114.5 (2)
H1A—C1—H1B	107.7	C14—C13—C17	103.33 (19)
C3—C2—C1	111.8 (3)	C12—C13—C17	106.8 (2)
C3—C2—H2A	109.3	O3—C14—C15	108.79 (19)
C1—C2—H2A	109.3	O3—C14—C8	107.92 (19)
C3—C2—H2B	109.3	C15—C14—C8	116.0 (2)
C1—C2—H2B	109.3	O3—C14—C13	105.59 (19)
H2A—C2—H2B	107.9	C15—C14—C13	104.0 (2)
O1—C3—C4	121.6 (3)	C8—C14—C13	113.95 (19)
O1—C3—C2	121.4 (3)	C14—C15—C16	105.1 (2)
C4—C3—C2	116.9 (3)	C14—C15—H15A	110.7
C5—C4—C3	123.9 (3)	C16—C15—H15A	110.7
C5—C4—H4	118.1	C14—C15—H15B	110.7
C3—C4—H4	118.1	C16—C15—H15B	110.7
C4—C5—C6	120.8 (3)	H15A—C15—H15B	108.8
C4—C5—C10	122.4 (3)	C15—C16—C17	107.2 (2)
C6—C5—C10	116.7 (2)	C15—C16—H16A	110.3
C5—C6—C7	112.2 (2)	C17—C16—H16A	110.3
C5—C6—H6A	109.2	C15—C16—H16B	110.3
C7—C6—H6A	109.2	C17—C16—H16B	110.3
C5—C6—H6B	109.2	H16A—C16—H16B	108.5
C7—C6—H6B	109.2	C20—C17—C16	112.9 (2)
H6A—C6—H6B	107.9	C20—C17—C13	116.6 (2)
C6—C7—C8	111.6 (2)	C16—C17—C13	105.62 (19)
C6—C7—H7A	109.3	C20—C17—H17	107.1
C8—C7—H7A	109.3	C16—C17—H17	107.1

C6—C7—H7B	109.3	C13—C17—H17	107.1
C8—C7—H7B	109.3	C10—C18—H18A	109.5
H7A—C7—H7B	108.0	C10—C18—H18B	109.5
C7—C8—C14	111.8 (2)	H18A—C18—H18B	109.5
C7—C8—C9	110.3 (2)	C10—C18—H18C	109.5
C14—C8—C9	112.24 (18)	H18A—C18—H18C	109.5
C7—C8—H8	107.4	H18B—C18—H18C	109.5
C14—C8—H8	107.4	C13—C19—H19A	109.5
C9—C8—H8	107.4	C13—C19—H19B	109.5
C11—C9—C8	109.72 (19)	H19A—C19—H19B	109.5
C11—C9—C10	112.22 (19)	C13—C19—H19C	109.5
C8—C9—C10	114.26 (19)	H19A—C19—H19C	109.5
C11—C9—H9	106.7	H19B—C19—H19C	109.5
C8—C9—H9	106.7	C22—C20—C21	107.7 (2)
C10—C9—H9	106.7	C22—C20—C17	125.9 (3)
C5—C10—C1	109.3 (2)	C21—C20—C17	126.4 (2)
C5—C10—C18	107.8 (2)	O4—C21—C20	105.0 (2)
C1—C10—C18	110.2 (2)	O4—C21—H21A	110.7
C5—C10—C9	109.2 (2)	C20—C21—H21A	110.7
C1—C10—C9	108.2 (2)	O4—C21—H21B	110.7
C18—C10—C9	112.1 (2)	C20—C21—H21B	110.7
C12—C11—C9	112.8 (2)	H21A—C21—H21B	108.8
C12—C11—H11A	109.0	C20—C22—C23	109.6 (3)
C9—C11—H11A	109.0	C20—C22—H22	125.2
C12—C11—H11B	109.0	C23—C22—H22	125.2
C9—C11—H11B	109.0	O5—C23—O4	120.3 (3)
H11A—C11—H11B	107.8	O5—C23—C22	130.4 (3)
O2—C12—C11	106.0 (2)	O4—C23—C22	109.3 (2)
O2—C12—C13	111.4 (2)	C12—O2—H2	109.5
C11—C12—C13	113.1 (2)	C14—O3—H3	109.5
O2—C12—H12	108.7	C23—O4—C21	108.4 (2)
C11—C12—H12	108.7	H1W—O6—H2W	114.3 (19)
C10—C1—C2—C3	-54.0 (4)	C7—C8—C14—O3	-64.4 (3)
C1—C2—C3—O1	-153.4 (3)	C9—C8—C14—O3	171.01 (18)
C1—C2—C3—C4	29.7 (4)	C7—C8—C14—C15	57.9 (3)
O1—C3—C4—C5	-177.1 (3)	C9—C8—C14—C15	-66.7 (3)
C2—C3—C4—C5	-0.2 (5)	C7—C8—C14—C13	178.7 (2)
C3—C4—C5—C6	170.4 (3)	C9—C8—C14—C13	54.1 (3)
C3—C4—C5—C10	-6.1 (5)	C19—C13—C14—O3	-48.0 (3)
C4—C5—C6—C7	131.9 (3)	C12—C13—C14—O3	-170.35 (19)
C10—C5—C6—C7	-51.3 (4)	C17—C13—C14—O3	76.7 (2)
C5—C6—C7—C8	53.6 (4)	C19—C13—C14—C15	-162.5 (2)
C6—C7—C8—C14	179.2 (2)	C12—C13—C14—C15	75.2 (2)
C6—C7—C8—C9	-55.2 (3)	C17—C13—C14—C15	-37.8 (2)
C7—C8—C9—C11	-178.5 (3)	C19—C13—C14—C8	70.3 (3)
C14—C8—C9—C11	-53.1 (3)	C12—C13—C14—C8	-52.1 (3)
C7—C8—C9—C10	54.5 (3)	C17—C13—C14—C8	-165.0 (2)
C14—C8—C9—C10	179.9 (2)	O3—C14—C15—C16	-75.9 (2)
C4—C5—C10—C1	-17.3 (4)	C8—C14—C15—C16	162.2 (2)

supplementary materials

C6—C5—C10—C1	166.1 (3)	C13—C14—C15—C16	36.3 (2)
C4—C5—C10—C18	102.6 (3)	C14—C15—C16—C17	-20.6 (3)
C6—C5—C10—C18	-74.1 (3)	C15—C16—C17—C20	125.5 (2)
C4—C5—C10—C9	-135.4 (3)	C15—C16—C17—C13	-3.0 (3)
C6—C5—C10—C9	47.9 (3)	C19—C13—C17—C20	22.6 (3)
C2—C1—C10—C5	46.7 (3)	C14—C13—C17—C20	-101.5 (2)
C2—C1—C10—C18	-71.5 (3)	C12—C13—C17—C20	144.6 (2)
C2—C1—C10—C9	165.6 (2)	C19—C13—C17—C16	148.9 (2)
C11—C9—C10—C5	-175.0 (2)	C14—C13—C17—C16	24.8 (2)
C8—C9—C10—C5	-49.3 (3)	C12—C13—C17—C16	-89.1 (2)
C11—C9—C10—C1	66.1 (3)	C16—C17—C20—C22	124.3 (3)
C8—C9—C10—C1	-168.16 (19)	C13—C17—C20—C22	-113.1 (3)
C11—C9—C10—C18	-55.6 (3)	C16—C17—C20—C21	-52.4 (4)
C8—C9—C10—C18	70.1 (3)	C13—C17—C20—C21	70.1 (4)
C8—C9—C11—C12	55.3 (3)	C22—C20—C21—O4	-0.7 (3)
C10—C9—C11—C12	-176.6 (2)	C17—C20—C21—O4	176.6 (2)
C9—C11—C12—O2	-179.6 (2)	C21—C20—C22—C23	0.4 (3)
C9—C11—C12—C13	-57.3 (3)	C17—C20—C22—C23	-176.9 (2)
O2—C12—C13—C19	48.2 (3)	C20—C22—C23—O5	179.4 (3)
C11—C12—C13—C19	-71.1 (3)	C20—C22—C23—O4	0.1 (3)
O2—C12—C13—C14	172.8 (2)	O5—C23—O4—C21	-179.9 (3)
C11—C12—C13—C14	53.5 (3)	C22—C23—O4—C21	-0.5 (3)
O2—C12—C13—C17	-76.6 (2)	C20—C21—O4—C23	0.7 (3)
C11—C12—C13—C17	164.1 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2—H2 \cdots O6 ⁱ	0.82	1.99	2.808 (3)	173
O3—H3 \cdots O2 ⁱⁱ	0.82	2.10	2.900 (3)	164
O6—H1W \cdots O1 ⁱⁱⁱ	0.86 (1)	1.94 (1)	2.801 (3)	174 (4)
O6—H2W \cdots O5 ^{iv}	0.86 (1)	1.91 (2)	2.741 (4)	162 (5)

Symmetry codes: (i) $x-1, y, z-1$; (ii) $x+1, y, z$; (iii) $x+1, y+1, z$; (iv) $x, y-1, z+1$.

Fig. 1




Fig. 2

